



First record of *Vampyrodes caraccioli* (Thomas, 1889) (Chiroptera, Phyllostomidae) in the state of Amazonas and its updated distribution in Brazil

Gerson P. Lopes^{1*}, Tamilly C. M. Santos^{1,2} and Paúl M. Velazco³

1 Institute for Sustainable Development Mamirauá, Research Group on Ecology of Terrestrial Vertebrates. Estrada do Bexiga, 2584, ZIP: 69553-255, Tefé, AM, Brazil

2 Vale Technology Institute, Sustainable Development. Rua Boaventura da Silva, 955, Nazaré. ZIP: 66055-090, Belém, PA, Brazil

3 American Museum of Natural History, Division of Paleontology, Central Park West at 79th Street, New York, NY 10024, USA

* Corresponding author. E-mail: gersonlps@hotmail.com

Abstract: The present note reports the first record of *Vampyrodes caraccioli* in the Brazilian state of Amazonas, and the central Amazon region, based on 44 specimens caught in the Auati-Paraná Extractive Reserve, north bank of the Solimões/Amazonas River. Our record fills a distribution gap of this species in the Brazilian Amazon and illustrates that the current knowledge on the distribution of Amazonian bats is far from complete.

Key words: Great Striped-Faced Bat; Amazon; geographical distribution

The genus *Vampyrodes* Thomas, 1900 currently includes two species, *V. caraccioli* (Thomas, 1889) and *V. major* Allen, 1908 (Velazco and Simmons 2011). *Vampyrodes caraccioli* can be easily distinguished from *V. major* by the following morphological characteristics: U-shaped uropatagium posterior margin with dense and short hair (> 2 mm) (V-shaped with dense and long hair in *V. major* [> 2 mm]); metacarpal III longer than metacarpal V (metacarpal III shorter than metacarpal V in *V. major*); two genal vibrissae present (three genal vibrissae in *V. major*); slender rostrum (broad and robust rostrum in *V. major*); parietal foramina well separated from the nuchal crest (parietal foramina are closer to the nuchal crest in *V. major*); groove between the occipital condyle and paracondylar process weakly developed (well developed in *V. major*); paraoccipital processes well developed (moderately developed in *V. major*); upper inner incisors broad (slender in *V. major*); M2 parastyle absent or poorly developed (well developed in *V. major*); M2 postentoconule absent or poorly developed (well developed in *V. major*); lower incisors are robust (slender in *V. major*); and absence of a cuspule on the m1 paracristid (cuspule present

on the m1 paracristid in *V. major*) (Velazco and Simmons 2011). In most of the external and craniodental measurements *V. caraccioli* is smaller than *V. major*, but there is some overlap in measurements where smaller individuals of *V. major* have similar measurements than larger *V. caraccioli* individuals (Velazco and Simmons 2011). *Vampyrodes major* is distributed from southern Mexico (Chiapas and Oaxaca) southwards, passing through Belize, Guatemala, Honduras, Nicaragua, Costa Rica, Panama, western Colombia and Ecuador (Willis et al. 1990; Gardner 2008; Velazco and Simmons 2011). *Vampyrodes caraccioli* occurs in eastern Colombia, eastern Ecuador, Peru, Bolivia, Venezuela, Trinidad and Tobago, French Guiana, Guyana, Suriname and Brazil (Willis et al. 1990; Velazco and Simmons 2011). Although it has an extensive distribution, *V. caraccioli* seems to be rare at lower latitudes and prefers moist and wet forests (Handley 1966; Gardner 2008; Velazco et al. 2010; Carvalho et al. 2014).

In Brazil, *V. caraccioli* has confirmed records (vouchered) from nine states: Acre (Taddei et al. 1990; Nogueira et al. 1999), Amapá (Martins et al. 2006, 2011; Martins 2012), Bahia (Faria et al. 2006), Mato Grosso (Sousa 2011), Mato Grosso do Sul (Cáceres et al. 2008), Pará (Thomas 1920; Kalko and Handley 2001; Peters et al. 2006; Tavares et al. 2012), Paraná (Scultroni 2009; Carvalho et al. 2014), Rio de Janeiro (Bezerra et al. 2004; Peracchi and Nogueira 2008; Souza et al. 2015), and São Paulo (Velazco et al. 2010) (Table 1). Velazco et al. (2010) and Carvalho et al. (2014) while reporting new Brazilian records of *V. caraccioli* from the states of São Paulo and Paraná respectively, overlooked some prior reports of the species (e.g., Aguiar 2000; Cáceres et al. 2008; Sousa 2011; Tavares 2013). Although *V. caraccioli* has been reported from the state of Goiás, this can be disputed

Table 1. Locality records of *Vampyrodes caraccioli* in Brazil. The code number refers to the points shown in Figure 1.

Code	Latitude	Longitude	State	Locality	Reference
1	-02.021611	-066.319444	Amazonas	Reserva Extrativista Auati–Paraná	This study
1	-01.998889	-066.204583	Amazonas	Reserva Extrativista Auati–Paraná	This study
2	-07.6666	-072.6666	Acre	Seringal Lagoinha, Rio Juruá	Taddei et al. (1990)
2	-07.4500	-073.6833	Acre	Parque Nacional da Serra do Divisor	Nogueira et al. (1999)
3	01.284	-051.589	Amapá	Floresta Nacional do Amapá	Martins et al. (2006)
3	01.387	-051.928	Amapá	Parque Nacional Montanhas do Tumucumaque	Martins et al. (2011)
3	01.387	-052.741	Amapá	Parque Nacional Montanhas do Tumucumaque	Martins et al. (2011)
3	01.3019	-051.5881	Amapá	Floresta Nacional do Amapá	Martins et al. (2011)
3	00.133	-051.067	Amapá	Área de Proteção Ambiental do Curiaú	Martins (2012)
3	00.492	-052.693	Amapá	Itapeuara (Rio Jari)	Martins (2012)
3	02.193	-054.588	Amapá	Parque Nacional Montanhas do Tumucumaque	Martins (2012)
3	01.845	-052.741	Amapá	Parque Nacional Montanhas do Tumucumaque	Martins (2012)
3	00.31	-052.44	Amapá	Parque Nacional Montanhas do Tumucumaque	Martins (2012)
3	00.579	-052.319	Amapá	Reserva de Desenvolvimento Sustentável do Rio Iratapuru	Martins (2012)
4	-015.300	-039.069	Bahia	Una	Faria et al. 2006
5	-014.6372	-052.3636	Mato Grosso	Nova Xavantina	Sousa (2011)
5	-14.6700	-052.3644	Mato Grosso	Nova Xavantina	Sousa (2011)
5	-14.6800	-052.3253	Mato Grosso	Nova Xavantina	Sousa (2011)
5	-16.4708	-054.6356	Mato Grosso	Rondonópolis	Universidade Federal do Paraná – Coleção Mastozoológica – DZUP 1550; 1547
6	-19.6117	-056.9514	Mato Grosso do Sul	Corumbá	Cáceres et al. (2008)
7	-01.450	-048.483	Pará	Belém, Área de Pesquisa Ecológica do Guamá	Thomas (1920)
7	-01.45	-048.50	Pará	Belém, Área de Pesquisa Ecológica do Guamá	Kalko and Handley (2001)
7	-01.4333	-048.4667	Pará	Belém, Área de Pesquisa Ecológica do Guamá	Kalko and Handley (2001)
7	-07.6833	-051.8667	Pará	Ourilândia do Norte, Área Indígena Kayapó	Peters et al. (2006)
7	-04.866	-050.750	Pará	Floresta Nacional de Carajás	Tavares et al. (2012)
8	-25.350	-048.767	Paraná	Reserva Natural Morro da Mina	Scultroni (2009)
8	-25.177	-048.308	Paraná	Reserva Natural Salto Morato	Carvalho et al. (2014)
9	-22.900	-043.233	Rio de Janeiro	Rio de Janeiro	Bezerra et al. (2004)
9	-22.942753	-044.040870	Rio de Janeiro	Mangaratiba	Peracchi and Nogueira (2008)
9	-23.066667	-043.883333	Rio de Janeiro	Marambaia	Lourenço et al. (2010)
9	-22.431	-042.756	Rio de Janeiro	Reserva Ecológica de Guapiaçu	Souza et al. (2015)
10	-23.726	-045.753	São Paulo	Parque Estadual da Serra do Mar	Velazco et al. (2010)

due to the lack of voucher material (Aguiar 2000; Silva-Júnior et al. 2007). In a similar way, the record of *V. caraccioli* from the state of Minas Gerais, presented by Tavares (2013), is not valid (V. Tavares pers. comm.). Carvalho et al. (2014, Figure 1), based on the study of Velazco et al. (2010, Figure 1), mistakenly indicated that the species was recorded in the state of Amazonas, Brazil. However, in the study by Velazco et al. (2010), the reported occurrences were in fact from Colombia and Peru on the border with the state of Amazonas, which may have been erroneously interpreted by Carvalho et al. (2014).

Here, we report for the first record of *V. caraccioli* from the state of Amazonas and provide an update on the distribution of the species in Brazil.

The specimens recorded here were collected during a bat inventory in the Auati–Paraná Extractive Reserve, state of Amazonas. Captures were performed using mist nets measuring 12 × 3 m, placed above the water column (depth of 1.5 m) at site 1 and set up at ground level at site 2. The nets were opened at 17:00 h, closed around 00:00 h, and checked every 30 minutes.

Sixteen specimens of *V. caraccioli*, all adult females, of which 11 were lactating and one was pregnant, were captured on 11 May 2015 in a secondary forest in a late stage of succession, in floodplain (01°59'56"S, 066°12'16.5"W) on the banks of the Auati–Paraná channel (site 1). In this location, together with the specimens of *V. caraccioli*, the following species were recorded: *Artibeus anderseni* Osgood, 1916; *Artibeus cinereus* (Gervais, 1856); *Artibeus planirostris* (Spix, 1823); *Carollia brevicauda* (Schinz, 1821); *Carollia perspicillata* (Linnaeus, 1758); *Chiroderma villosum* Peters, 1860; *Glossophaga soricina* (Pallas, 1766); *Myotis riparius* Handley, 1960; *Phyllostomus elongatus* (E. Geoffroy St.-Hilaire, 1810); *Platyrrhinus brachycephalus* (Rouk & Carter, 1972); *Rhynchoycteris naso* (Wied-Neuwied, 1820); *Saccopteryx bilineata* (Temminck, 1838); *Uroderma bilobatum* Peters, 1866; and *Uroderma magnirostrum* Davis, 1968.

Twenty-eight additional specimens of *V. caraccioli*, of which one was a male with a scrotal sac visible, but most were lactating females and only female showed no evidence of pregnancy or lactation, were captured on 16 May 2015 on the edge of the secondary forest, dominated




Figure 1. Known records of *Vampyromes caraccioli* in Brazil. Numbers refer to entries in Table 1.

by fruit trees, in an upland rain forest ($02^{\circ}01'07.8''S$, $066^{\circ}19'10''W$) on the shores of Lake Inambé (site 2). The following species were also captured in this locality: *Artibeus obscurus* (Schinz, 1821); *Artibeus planirostris*; *Artibeus gnomus* Handley, 1987; *Carollia perspicillata*; *Chiroderma villosum*; *Diphylla ecaudata* Spix, 1823; and *Platyrrhinus brachycephalus*.

Data collection was conducted with the authorization of the Chico Mendes Institute for Biodiversity Conservation, under the license of System of Authorization and Information on Biodiversity 42.111-2. Ten specimens were collected and sacrificed following the protocols defined by the American Society of Mammalogy (Sikes et al. 2011). The voucher specimens were fixed in 4%



Figure 2. Photographs of a female specimen of *V. caraccioli* (IDSM 1182) collected in the Auati-Paraná Extractive Reserve. (A) live specimen, (B) dorsal views of the skull and mandible, (C) dorsolateral view of the skull, and (D) ventral view of the skull. Photos by Gerson P. Lopes. Scale bar = 2 mm.

formalin and later preserved in 70% ethanol and are deposited in the Mammal Collection of the Institute for Sustainable Development Mamirauá, Amazonas, Brazil (IDSM) under the collection numbers IDMS 1181–1190.

The identification of specimens was based on the characters provided by Velazco and Simmons (2011) which are as follows: light to dark brown on the back, with lighter fur on the venter; presence of a distinct dorsal white stripe that runs from the top of the head to the base of the uropatagium; four distinct facial white stripes, with the suborbital stripes ranging from the corner of its mouth to the base of the ear, and the suprabital stripes ranging from the noseleaf to above the ear (Figure 2); absence of a groove on the mesial portion of second upper premolar; presence of two upper molars and three lower molars; absence of lingual cingula on the metacones of M1 and M2; absence of a lingual cingulum on M1; and absence of a cusp on the mesial portion of the protocone on M1.

Our specimens represent the first record of *V. caraccioli* from the state of Amazonas and central Amazon. These records fill a gap in the distribution of this species in the Brazilian Amazon, whose closest record is more than 940

km to the southwest, at Seringal Lagoinha, state of Acre.

The bat fauna is among the richest among all of the mammal groups distributed in the Amazon (Voss and Emmons 1996). Despite being the biome with the largest number of species in Brazil, only 23.9% of the Brazilian Amazon is represented by at least one record of bats, and only 5.1% of its area is considered minimally sampled (Bernard et al. 2011a, 2011b; Martins et al. 2011). Fragmented records of bats in the Amazon show that large gaps still need to be filled, including portions of northern and southern Amazonas state that are distant from the state capital at Manaus (Bernard et al. 2011b). Moreover, even in intensively sampled sites, such as the Biological Dynamics of Forest Fragments Project in Manaus (Sampaio et al. 2003), *V. caraccioli* has not been recorded.

We emphasize that although *V. caraccioli* is not commonly captured in Brazil, all specimens obtained here were collected within two evenings. This high number of specimens may be explained by the abundance of various fruiting trees in the study sites in proximity to the mist nets (e.g., *Cecropia* sp., *Ficus* sp., *Piper* sp., *Pouteria* sp., and *Vismia* sp.). In the other 18 days of the

expedition in the primary forest, no other *Vampyrodes* specimen was captured. The records of *V. caraccioli* in the Cerrado biome (Aguiar 2000; Silva-Júnior et al. 2007; Sousa 2011) also contradict the belief that this species is a frugivore strictly associated with tropical rain forests (Gardner 1977, 2008).

Although the Amazon harbors 87% of Brazil's bat fauna, basic knowledge on the distribution of Amazon bats is far from complete, with intensive surveys at only a few sites and many areas without any surveys whatsoever (Bernard et al. 2011b). In fact, the Amazon is the region least sampled for bats, which emphasizes the need of further research to better understand basic knowledge on bat distribution (Bernard et al. 2011b).

ACKNOWLEDGEMENTS

We thank the Ministry of Science, Technology and Innovation (Brazil) for their financial support; Caetano Franco for producing the map; and Kerry Kline for critical comments on an early draft of this manuscript. We thank Mr. Weliton Pereira and Mr. Jânio da Silva for their help in the field.

LITERATURE CITED

- Aguiar, L.S.M. 2000. Comunidades de Morcegos do Cerrado no Brasil Central [Doctoral thesis]. Brasília: Universidade de Brasília. 101 pp.
- Allen, G.M. 1908. Notes on Chiroptera. Bulletin of the Museum of Comparative Zoology 52: 25–62. <http://biodiversitylibrary.org/page/4336753>
- Bernard, E., V.C. Tavares and E. Sampaio. 2011a. Compilação atualizada das espécies de morcegos (Chiroptera) para a Amazônia Brasileira. Biota Neotropica 11(1): 1–13. doi: [10.1590/S1676-06032011000100003](https://doi.org/10.1590/S1676-06032011000100003)
- Bernard, E., L.M.S. Aguiar and R.B. Machado 2011b. Discovering the Brazilian bat fauna: a task for two centuries? Mammal Review 41(1): 23–39. doi: [10.1111/j.1365-2907.2010.00164.x](https://doi.org/10.1111/j.1365-2907.2010.00164.x)
- Bezerra, A.M.R., M. Baptista, S.M. Franco and J.A. Oliveira. 2004. A coleção de mamíferos preservados em meio líquido do Museu Nacional. Publicações Avulsas do Museu Nacional 101: 1–11.
- Cáceres, N.C., A.C. Carmignotto, E. Fischer and C.F. Santos. 2008. Mammals from Mato Grosso do Sul, Brazil. Check List 4(3): 321–335. doi: [10.15560/4.3.321](https://doi.org/10.15560/4.3.321)
- Carvalho, F., V. Mottin, J.M.D. Miranda and F.C. Passos 2014. First record of *Vampyrodes caraccioli* (Thomas, 1889) (Chiroptera: Phyllostomidae) for the state of Paraná, and range extension to southern region of Brazil. Check List 10(5): 1189–1194. doi: [10.15560/10.5.1189](https://doi.org/10.15560/10.5.1189)
- Davis, W.B. 1968. Review of the genus *Uroderma* (Chiroptera). Journal of Mammalogy 49(4): 676–698. doi: [10.2307/1378728](https://doi.org/10.2307/1378728)
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from the Atlantic rainforest of southern Bahia, Brazil. Biota Neotropica 6(2): 1–13. doi: [10.1590/S1676-06032006000200022](https://doi.org/10.1590/S1676-06032006000200022)
- Gardner, A.L. 1977. Feeding habits; pp. 355–357, in: R.J. Baker, J.K. Jones, Jr. and D.C. Carter (eds.). Biology of bats of the New World family Phyllostomatidae, Part II. Special Publications of the Museum Texas Tech University 13.
- Gardner, A.L. 2008. Genus *Vampyrodes* O. Thomas, 1900; pp. 355–357, in: A.L. Gardner (ed.). Mammals of South America, Volume 1, marsupials, xenarthrans, shrews, and bats. Chicago: The University of Chicago Press.
- Geoffroy St.-Hilaire, É. 1810. Sur les phyllostomes et les mégadermes, deux genres de la famille des chauve-souris. Annales Muséum National d'Histoire Naturelle 15: 157–198.
- Gervais, P. 1856. Deuxième mémoire. Documents zoologiques pour servir à la monographie des chéiroptères Sud-Américains; pp. 25–88, in: P. Gervais (ed.). Mammifères. In: Animaux nouveaux ou rares recueillis pendant l'expédition dans les parties centrales de l'Amérique du Sud, de Rio de Janeiro à Lima, et de Lima au Para; exécutée par ordre du gouvernement français pendant les années 1843 à 1847, sous la direction du comte Francis de Castelnau (F. de Castelnau ed.). Paris: P. Bertrand.
- Handley, C.O., Jr. 1960. Descriptions of new bats from Panama. Proceedings of the United States National Museum 112: 459–79. doi: [10.5479/si.00963801.112-3442.459](https://doi.org/10.5479/si.00963801.112-3442.459)
- Handley, C.O., Jr. 1966. Checklist of the mammals of Panama; pp. 753–795, in: R.L. Wenzel and V.J. Tipton (eds.). Ectoparasites of Panama. Chicago: Field Museum of Natural History.
- Handley, C.O., Jr. 1987. New species of mammals from northern South America: fruit-eating bats, genus *Artibeus* Leach; pp. 163–172, in: B.D. Patterson and R.M. Timm (eds.). Studies in Neotropical mammalogy, essays in honor of Philip Hershkovitz. Fieldiana Zoology 39. <http://biodiversitylibrary.org/page/4465287>
- Kalko, E.K.V. and C.O. Jr. Handley. 2001. Neotropical bats in the canopy: diversity, community structure, and implications for conservation. Plant Ecology 153: 319–333. doi: [10.1023/A:1017590007861](https://doi.org/10.1023/A:1017590007861)
- Linnaeus, C. 1758. Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata. Holmiae: Laurentii Salvii. 1: 1–824 pp. doi: [10.5962/bhl.title.542](https://doi.org/10.5962/bhl.title.542)
- Lourenço, E.C., L.M. Costa and C.E.L. Esbérard. 2010. Bat diversity of Ilha da Marombaia, southern Rio de Janeiro state, Brazil (Chiroptera, Mammalia). Brazilian Journal of Biology 70(3): 511–519. doi: [10.1590/S1519-69842010000300007](https://doi.org/10.1590/S1519-69842010000300007)
- Martins, A.C.M., E. Bernard and R. Gregorin, 2006. Rapid biological surveys of bats (Mammalia, Chiroptera) in three conservation units in Amapá, Brazil. Revista Brasileira de Zoologia 23(4): 1175–1184. doi: [10.1590/S0101-81752006000400026](https://doi.org/10.1590/S0101-81752006000400026)
- Martins, A.C.M., E. Bernard, R. Gregorin and W.A.S. Silva. 2011. Filling data gaps on the diversity and distribution of Amazonian bats (Chiroptera): the case of Amapá, easternmost Brazil. Zoologia 28(2): 177–185. doi: [10.1590/S1984-46702011000200004](https://doi.org/10.1590/S1984-46702011000200004)
- Martins, A.C.M. 2012. Morcegos do Amapá e a resposta da família Phyllostomidae às variáveis de composição de paisagens pouco fragmentadas na Amazônia [Master's thesis]. Brasília: Universidade de Brasília. 164 pp.
- Nogueira, M.R., A. Pol and A.L. Peracchi. 1999. New records of bats from Brazil with a list of additional species for the chiropteran fauna of the state of Acre, western Amazon basin. Mammalia 63(3): 363–368.
- Osgood, W.H. 1916. Mammals of the Collins-Day South American Expedition. Field Museum of Natural History, Zoology Series 10: 199–216. doi: [10.5962/bhl.title.2632](https://doi.org/10.5962/bhl.title.2632)
- Pallas, P.S. 1766. Miscellanea zoologica quibus novae imprimis atque obscurae animalium species describuntur et observationibus iconibusque illustrantur. Hague Comitum: P. van Cleef xii+224 pp. doi: [10.5962/bhl.title.69851](https://doi.org/10.5962/bhl.title.69851)
- Peracchi, A.L. and M.R. Nogueira. 2008. Quirópteros da Mata Atlântica do Sudeste do Brasil; pp. 279–298, in: S.M. Pacheco, R.V. Marques and C.E.L. Esbérard (eds.). Morcegos no Brasil: biologia, sistemática, ecologia e conservação. Porto Alegre: Editora Armazém Digital.
- Peters, S.L., J.R. Malcolm and B. Zimmerman. 2006. Effects of selective logging on bat communities in the southeastern Amazon. Conservation Biology 20(5): 1410–1421. doi: [10.1111/j.1523-173X.2006.00833.x](https://doi.org/10.1111/j.1523-173X.2006.00833.x)

1739.2006.00526.x

- Peters, W. 1860. Eine neue Gattung von Flederthieren, *Chiroderma villosum*, aus Brasilien. Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin 1860: 747–755. <http://biodiversitylibrary.org/page/36275992>
- Peters, W. 1866. Über die brasilianischen, von Spix beschriebenen Flederthiere. Monatsberichte der Königlich preussischen Akademie der Wissenschaften zu Berlin 1866: 568–588. <http://biodiversitylibrary.org/page/35169029>
- Rouk, C.S. and D.C. Carter. 1972. A new species of *Vampyrops* (Chiroptera: Phyllostomatidae) from South America. Occasional Papers of the Museum, Texas Tech University 1: 1–7.
- Sampaio, E.M., E.K.V. Kalko, E. Bernard, B. Rodríguez-Herrera and C.O. Handley. 2003. A biodiversity assessment of bats (Chiroptera) in a tropical lowland rainforest of Central Amazonia, including methodological and conservation considerations. Studies on Neotropical Fauna and Environment 38(1): 17–31. doi: 10.1076/snfe.38.1.17.14035
- Schinz, H.R. 1821. Das Thierreich eingetheilt nach dem Bau der Thiere als Grundlage ihrer Naturgeschichte und der vergleichenden Anatomie von dem Herrn Ritter von Cuvier. Erster band. Säugetiere und Vögel. Stuttgart/Tübingen: J. G. Cotta'schen Buchhandlung. 1: xxxviii + 894 pp.
- Scultori C. 2009. Comunidade de morcegos, interação com flores e estratificação vertical em Mata Atlântica no sul do Brasil [Master's thesis]. Campinas: Universidade Estadual de Campinas. 158 pp.
- Sikes, R.S., W.L. Gannon and the Animal Care and Use Committee of the American Society of Mammalogists. 2011. Guidelines of the American Society of Mammalogists for the use of wild animals in research. Journal of Mammalogy 92(1): 235–253. doi: 10.1644/10-MAMM-F-355.1
- Silva-Júnior, N.J., H.L.R. Silva, M.C. Costa, A.B. Marcus, M.L.S. Tonial, R.S. Ribeiro, L.A. Moreira and A.M. Pessoa. 2007. Avaliação preliminar da fauna silvestre do vale do rio Caiapó, Goiás: implicações para a conservação da biodiversidade regional. Estudos 34: 1057–1094. <http://seer.ucg.br/index.php/estudos/article/view/256>
- Sousa R.F. 2011. Biologia, ecologia e aspectos citogenéticos de morcegos (Chiroptera, Mammalia) em remanescentes florestais de Cerrado no leste mato-grossense [Master's thesis]. Nova Xavantina: Universidade do Estado de Mato Grosso. 96 pp.
- Souza, R.F., R.L.M. Novaes, A.C. Siqueira, C. Sauwen, G. Jacob, C.E.L. Santos, S. Felix, E. Ribeiro, C. Sant'Anna, D. Vrcibradic, L.S. Avilla, I. Sbragia and R.T. Santori. 2015. Bats (Mammalia, Chiroptera) in a remnant of Atlantic Forest, Rio de Janeiro, southeastern Brazil. Neotropical Biology and Conservation 10(1): 9–14. doi: 10.4013/nbc.2015.101.02
- Spix, J.B. von. 1823. Simiarum et vespertilionum Brasiliensium species novae, ou, Histoire naturelle des espèces nouvelles de singes et de chauves-souris observées et recueillies pendant le voyage dans l'intérieur du Brésil exécuté par ordre de S. M. le Roi de Bavière dans les années 1817, 1818, 1819, 1820. Monachii: Francisci Seraphici Hübschmanni, vii+72 pp.
- Taddei, V.A., I.M. Rezende and D. Camora. 1990. Notas sobre uma coleção de morcegos de Cruzeiro do Sul, Rio Juruá, estado do Acre (Mammalia, Chiroptera). Boletim do Museu Paraense Emílio Goeldi, Série Zoologia 6(1): 75–88.
- Tavares, V.C., C.F.S. Palmuti, R. Gregorin and T.T. Dornas. 2012. Morcegos; pp. 1–229, in: F.D. Martins, A.F. Castilho, J. Campos, F.M. Hatano and G.R. Samir (eds.). Fauna da Floresta Nacional de Carajás - Estudos Sobre Vertebrados Terrestres. Belo Horizonte: Nitro. <http://www.icmbio.gov.br/portal/images/Carajas.pdf>
- Tavares, V.C. 2013. Phyllostomid bat wings from Atlantic Forest bat ensembles: an ecomorphological study. Chiroptera Neotropical 19(3): 57–70. <https://chiroptera.unb.br/index.php/cn/article/view/219/168>
- Temminck, C.J. 1838. Over de geslachten *Taphozous*, *Emballonura*, *Urocyptus* en *Diclidurus*. Tijdschrift voor Natuurlijke Geschiedenis en Physiologie, Amsterdam 5: 1–34. <http://biodiversitylibrary.org/page/13474775>
- Thomas, O. 1889. Description of a new stenodermatous bat from Trinidad. Annals and Magazine of Natural History (6) 4: 167–170. <http://biodiversitylibrary.org/page/19301659>
- Thomas, O. 1900. Descriptions of new Neotropical mammals. Annals and Magazine of Natural History (7) 5: 269–274. <http://biodiversitylibrary.org/page/24340185>
- Thomas, O. 1920. On mammals from the lower Amazons in the Goeldi Museum, Para. Annals and Magazine of Natural History (9) 6: 266–283. <http://biodiversitylibrary.org/page/15687549>
- Velazco, P.M., C.C. Aires, A.P. Carmignotto and A.M.R. Bezerra. 2010. Mammalia, Chiroptera, Phyllostomidae, *Vampyrodes caraccioli* (Thomas, 1889): range extension and revised distribution map. Check List 6(1): 49–51. <http://www.checklist.org.br/getpdf?NGD141-09>
- Velazco, P.M. and N.B. Simmons. 2011. Systematics and taxonomy of great striped-faced bats of the genus *Vampyrodes* Thomas, 1900 (Chiroptera: Phyllostomidae). American Museum Novitates 3710: 1–35. <http://hdl.handle.net/2246/6104>
- Voss, R.S. and L.H. Emmons. 1996. Mammalian diversity in Neotropical lowland rainforests: a preliminary assessment. Bulletin of the American Museum of Natural History 230: 1–115. <http://hdl.handle.net/2246/1671>
- Wied-Neuwied, M.P. zu. 1820. Reise nach Brasilien in den Jahren 1815 bis 1817. Frankfurt a. M.: Heinrich Ludwig Bröner. 1: xxxvi+380 pp. doi: 10.5962/bhl.title.85967
- Willis, K.B., M.R. Willig and J.K. Jones. 1990. *Vampyrodes caraccioli*. Mammalian Species 359: 1–4. <http://www.science.smith.edu/msi/pdf/i0076-3519-359-01-0001.pdf>

Author contributions: GPL and TCMS participated in the capture, collection and identification of the specimens, as well as the drafting, discussion and writing of the article. PMV participated in the drafting, discussion and writing of the article.

Received: 1 December 2015

Accepted: 23 May 2016

Academic editor: Sergio Solari