

# *Peropteryx trinitatis* Miller, 1899 (Chiroptera, Emballonuridae): first record in central Brazil and revised distribution map

Adriano Silva dos Santos, Daiana Cardoso Silva and Karina de Cassia Faria\*

Universidade do Estado de Mato Grosso, Faculdade de Ciências Agrárias, Biológicas e Sociais Aplicadas. Av. Prof. Dr. Renato Figueiro Varella, s/n, CEP 78690-000, Nova Xavantina, MT, Brazil

\* Corresponding author. E-mail: [karinafaria@unemat.br](mailto:karinafaria@unemat.br)

**Abstract:** Reports of *Peropteryx trinitatis* in Brazil are restricted to the northern and northeastern regions. In this study, we extend its known geographic range to include central Brazil, with the first report of *P. trinitatis* in the Cerrado of the state of Mato Grosso. This is the southernmost known record for this species, 900 km from the closest record.

**Key words:** Dog-like Bat; Cerrado; new record; Mato Grosso

*Peropteryx* is composed of five species, *P. kappleri* Peters, 1867, *P. leucoptera* Peters, 1867, *P. macrotis* (Wagner, 1843), *P. pallidoptera* Lim et al. 2010, and *P. trinitatis* Miller, 1899 (Solari and Martinez-Arias 2014). *Peropteryx trinitatis* was once considered a subspecies of *P. macrotis* due to both their extensive morphological similarity and to an alleged lack of sympatry (Jones and Hood 1993; Koopman 1993, 1994); however, dental and external morphological comparisons and confirmation that both taxa may occur in sympatry have verified its classification as distinct species (Handley 1976; Brosset and Charles-Dominique 1990; Simmons and Voss 1998).

*Peropteryx trinitatis* is an insectivorous bat, weighing from 3 to 6 g and with a forearm length ranging from 36 to 43 mm (Hood and Gardner 2008; Lim et al. 2010; Reis et al. 2013). The species is sexually dimorphic, with females usually larger than males (Lim et al. 2010). It has large and wide ears which become slightly narrower at the tip; a conical muzzle, and dark brown to reddish brown fur (Miller 1899; Simmons and Voss 1998). *Peropteryx trinitatis* is known to occur in Aruba, Venezuela, French Guiana, Guyana, Trinidad and Tobago and Brazil; its type locality is Port of Spain, Trinidad (Miller 1889) (Figure 1; Table 1). In these countries, *P. trinitatis* has been found in tropical and subtropical moist broadleaf forests, tropical and subtropical grasslands, savannas


and shrublands, and in deserts and xeric shrublands (Figure 1).

There have been few captures of the *P. trinitatis* in Brazil. It has thus far been reported only in the north and northeast regions of the country covering the Brazilian biomes Amazon (states of Amazonas, Amapá and Pará), Caatinga (state of Bahia) and Cerrado savanna (state of Maranhão) (Table 1). Here we report a new record for the Central-West Region of Brazil, in the Cerrado savanna.

We captured two adult male *P. trinitatis* in Mário Viana Municipal Park (14°43'22" S, 052°21'39" W), a protected area (492 ha) located in the eastern portion of the Cerrado biome, in the city of Nova Xavantina, state of Mato Grosso. The predominant vegetation type in the municipal park is Cerrado *sensu strictu*, but there are patches of woodland, rocky Cerrado, and a gallery forest (Silva et al. 2008). The climate, according to the Köppen classification is tropical humid (*Aw*), the average annual temperature is 24°C (Vianello and Alves 2000), and mean rainfall is around 1,500 mm, with a dry season between April and September and a rainy season between October and March (Pirani et al. 2009).

The first specimen was captured on 10 May 2014 and the second specimen was captured on 11 April 2015. The captures were performed with the use of 6 × 3 m mist nets set up during the day at the opening to a roost site. The opening was characterized as a gap in a rocky wall within the gallery forest. The permit to collect the specimens was granted by the Chico Mendes Institute for Biodiversity Conservation (ICMBio), Ministry of Environment (Process #18276-1). The specimens were identified following Vizotto and Taddei (1973), Lim and Engstrom (2001), and Simmons and Voss (1998), and deposited in the scientific collection of the Universidade do Estado do Mato Grosso (UNEMAT), campus of Nova Xavantina, under the collection numbers RM 460 and RM 498 (Figure 2).

The dorsum of our *P. trinitatis* specimens were dark



**Figure 1.** Known geographic distribution of *Peropteryx trinitatis*. The black dots represent previous reports of the species, the star represents the type locality and the red dot represents the first report of the species in central Brazil, in Nova Xavantina, state of Mato Grosso. Biomes classified according to Olson et al. (2001; available at <http://www.worldwildlife.org/publications/terrestrial-ecoregions-of-the-world>).

brown in color, while the venters were lighter brown and a reddish tone (Figure 2A). According to Brosset and Charles-Dominique (1990), *P. trinitatis* has a shorter and more round skull than *P. macrotis*. The shape of the skull of the specimens captured in Mato Grosso was similar to those illustrated by these authors (Figure 2B). The anterior upper premolars of our specimens of *P. trinitatis* were peg-like and without well-defined anterior

and posterior cusps (Figure 2C), consistent with the description of Simmons and Voss (1998). External and cranial measurements (in mm) were found to be similar to the published morphometric data for *P. trinitatis* with small variations in measurements of thumb, foot, calcaneus, and greatest mandible length (Table 2). The dental formula was  $I1/3, C1/1, P2/2, M3/3 = 32$ .

*Peropteryx trinitatis* is categorized by the IUCN

**Table 1.** Locations where *Peropteryx trinitatis* is known to occur.

| Country             | State/department             | Locality                                     | Latitude | Longitude | Reference                            |
|---------------------|------------------------------|----------------------------------------------|----------|-----------|--------------------------------------|
| Aruba               |                              | 25 kilometer north of the coast of Venezuela | 12.488   | -69.970   | Bekker (1996)                        |
| Brazil              | Amazonas                     | Amanã Sustainable Development Reserve        | -02.621  | -64.616   | Marques et al. (2016)                |
| Brazil              | Amapá                        | APA Curiaú, Macapá                           | 00.116   | -51.050   | Silva et al. (2013)                  |
| Brazil              | Bahia                        | Queimadas                                    | -11.110  | -39.961   | Hood and Gardner (2008)              |
| Brazil              | Maranhão                     | Alto Parnaíba                                | -09.067  | -46.008   | Hood and Gardner (2008)              |
| Brazil              | Mato Grosso                  | Mário Viana Municipal Park, Nova Xavantina   | -14.722  | -52.360   | This study                           |
| Brazil              | Pará                         | Kararaô Cave                                 | -03.141  | -51.818   | Zortéa et al. (2015)                 |
| Brazil              | Pará                         | Utinga, Belém                                | -01.958  | -49.032   | Hood and Gardner (2008)              |
| French Guiana       |                              | Cayenne                                      | 04.825   | -52.382   | Brosset and Charles-Dominique (1990) |
| Guyana              | Upper Takutu–Upper Essequibo | 12 miles E of Dadanawa                       | 02.945   | -58.920   | Hood and Gardner (2008)              |
| Trinidad and Tobago | Trinidad                     | Saint George County, Port of Spain           | 10.047   | -61.883   | Miller (1899)                        |
| Venezuela           | Amazonas                     | San Juan                                     | 06.868   | -66.029   | Handley (1976)                       |
| Venezuela           | Apure                        | 8 km NW of Puerto Páez                       | 07.930   | -67.092   | Handley (1976)                       |
| Venezuela           | Bolívar                      | Hato La Florida                              | 07.468   | -65.846   | Lim et al. (2010)                    |
| Venezuela           | Bolívar                      | Rio Caura                                    | 03.595   | -65.584   | Ochoa et al. (2008)                  |
| Venezuela           | Carabobo                     | Rio Guárico, Belén                           | 10.167   | -67.667   | Bisbal and Rodríguez (2010)          |
| Venezuela           | Falcón                       | 13 km NNE of Mirimire                        | 11.071   | -69.566   | Handley (1976)                       |
| Venezuela           | Miranda                      | 4 km SW of Birongo                           | 09.891   | -66.029   | Handley (1976)                       |
| Venezuela           | Nueva Esparta                | Margarita Island                             | 11.030   | -64.290   | Rawlins (2011)                       |

**Table 2.** External and cranial measurements of *Peropteryx trinitatis*.

| Sex                         | Nova Xavantina, Brazil <sup>1</sup> |        | Port of Spain, Trinidad <sup>2</sup> | Cayenne, French Guiana <sup>3</sup> |        | Venezuela <sup>4</sup> |         |
|-----------------------------|-------------------------------------|--------|--------------------------------------|-------------------------------------|--------|------------------------|---------|
|                             | Females                             |        | Female                               | Male                                | Female | Males                  | Females |
| Measurements (mm)           | RM 460                              | RM 498 | N° 7496                              | N° 305                              | N° 595 | –                      | –       |
| Forearm                     | 39.2                                | 39.0   | 40.0                                 | 39.5                                | 40.4   | 38.3                   | 41.5    |
| Thumb                       | 4.1                                 | 4.6    | 7.0                                  | –                                   | –      | –                      | –       |
| Ear                         | 12.3                                | 10.6   | 11.0                                 | –                                   | –      | 12.9                   | 12.8    |
| Tibia                       | 17.3                                | 16.1   | 18.0                                 | –                                   | –      | –                      | –       |
| Foot                        | 5.3                                 | 6.1    | 6.4                                  | –                                   | –      | 6.9                    | 7.6     |
| Calcaneus                   | 10.4                                | 11.5   | 17.0                                 | –                                   | –      | –                      | –       |
| Greatest length of skull    | 14.0                                | 14.5   | 13.8                                 | 13.2                                | 13.2   | 12.6                   | 13.4    |
| Condylocanine length        | 12.6                                | 13.2   | –                                    | –                                   | –      | 11.4                   | 12.0    |
| Condylobasal length         | 13.1                                | 13.9   | 12.0                                 | –                                   | –      | –                      | –       |
| Maxillary toothrow length   | 5.0                                 | 5.3    | 5.4                                  | 4.8                                 | 4.7    | 4.6                    | 5.0     |
| Zygomatic breadth           | 7.9                                 | 8.0    | 8.0                                  | 7.8                                 | 7.7    | 7.4                    | 7.9     |
| Braincase breadth           | 6.3                                 | 6.4    | 6.4                                  | –                                   | –      | 6.0                    | 6.2     |
| Mastoid breadth             | 7.0                                 | 6.9    | 7.4                                  | –                                   | –      | 6.8                    | 7.1     |
| Postorbital breadth         | 2.3                                 | 2.8    | 2.6                                  | 2.9                                 | 3.0    | 2.8                    | 2.5     |
| Width across molars         | 5.7                                 | 6.2    | –                                    | 6.0                                 | 6.2    | –                      | –       |
| Width across canines        | 3.5                                 | 3.3    | –                                    | –                                   | –      | –                      | –       |
| Greatest of length mandible | 8.5                                 | 8.8    | 9.4                                  | 9.4                                 | 9.3    | –                      | –       |
| Weight (g)                  | 5.0                                 | 4.9    | –                                    | 4.0                                 | 4.4    | 3.6                    | 4.1     |

Present study<sup>1</sup>, Miller–Jr. (1899)<sup>2</sup>, Brosset and Charles–Dominique (1990)<sup>3</sup> and Lim et al. (2010)<sup>4</sup>.

(International Union for Conservation of Nature) (Sampaio 2016) and by Brazilian Ministry of the Environment (MMA 2014) as Data Deficient because of the lack of knowledge on its current range, and also because of the need for more information on its biology and ecology. The species is considered to be uncommon throughout its geographic range (Emmons and Feer 1990).


This capture is the first report of the species in the state of Mato Grosso and central Brazil. It expands the known distribution range of this species about 900 km

south from the closest record in the Brazilian state of Maranhão (Table 1). This finding emphasizes the need for more studies in areas that are currently lacking information on species richness, such as eastern Mato Grosso state.

## ACKNOWLEDGEMENTS

We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and the Universidade do Estado de Mato Grosso (UNEMAT) for financial support, Dr. Marlon Zórtéa for assistance in identifying





**Figure 2.** *Peropteryx trinitatis* (RM 498) captured in Nova Xavantina, Mato Grosso State, Brazil. **A.** Adult male. **B.** Dorsal, ventral and lateral views of skull and lateral and ventral views of mandible. Scale bar = 4.4 mm. **C.** Close-up lateral view of its upper right dentition (the arrow indicates the anterior upper premolar). Scale bar = 1.2 mm.

the specimens, and the two anonymous reviewers for their contributions.

## LITERATURE CITED

- Bekker, J.P. 1996. The mammals of Aruba. Nijmegen: Vereniging voor Zoogdierkunde en Zoogdierbescherming. 87 pp.
- Bisbal, F.J.E. and A.N. Rodríguez. 2010. Mamíferos de la cuenca del río Guárico, estados Aragua, Carabobo y Guárico, Venezuela. Memoria de la Fundación La Salle de Ciencias Naturales 172: 69–89. [http://www.fundacionlasalle.org.ve/userfiles/7\\_Mem2009%20Tom\(69\)172%20p%2069-89.pdf](http://www.fundacionlasalle.org.ve/userfiles/7_Mem2009%20Tom(69)172%20p%2069-89.pdf)
- Brosset, A. and P. Charles-Dominique. 1990. The bats from French Guiana: a taxonomic, faunistic and ecological approach. Mammalia 54: 509–560. doi: [10.1515/mamm.1990.54.4.509](https://doi.org/10.1515/mamm.1990.54.4.509)
- Emmons, L.H. and F. Feer. 1990. Neotropical rainforest mammals: a field guide. Chicago/London: The University of Chicago Press. 281 pp.
- Handley, C.O., Jr. 1976. Mammals of the Smithsonian Venezuelan Project. Brigham Young University Science Bulletin, Biological series 20(5): 1–89. [https://ojs.lib.byu.edu/spc/index.php/BYU\\_SciBullBioS/article/view/30679/29161](https://ojs.lib.byu.edu/spc/index.php/BYU_SciBullBioS/article/view/30679/29161)
- Hood, C. and A.L. Gardner. 2008. Family Emballonuridae Gervais, 1856; pp. 188–207, in: A.L. Gardner (ed.). Mammals of South America. Volume I: marsupials, xenarthrans, shrews, and bats. Chicago: The University of Chicago Press.
- Jones, J. K., Jr. and C. S. Hood. 1993. Synopsis of South American bats of the family Emballonuridae. Occasional Papers, The Museum, Texas Tech University 155: 1–32.
- Koopman, K.F. 1993. Order Chiroptera: pp. 137–242, in: D.E. Wilson and D.M. Reeder (eds). Mammal species of the world, a taxonomic and geographic reference. 2nd edition. Washington, DC: The Smithsonian Institution Press.
- Koopman, K. F. 1994. Chiroptera: systematics. Handbuch der Zoologie, volume 8, part 60, Mammalia. Berlin: Walter de Gruyter. 217 pp.
- Lim, B. K. and M. D. Engstrom. 2001. Species diversity of bats (Mammalia: Chiroptera) in Iwokrama Forest, Guyana, and the Guianan subregion: implications for conservation. Biodiversity and Conservation 10: 613–657. doi: [10.1023/A:1016660123189](https://doi.org/10.1023/A:1016660123189)
- Lim, B.K., M.D. Engstrom, F.A. Reid, N.B. Simmons, R.S. Voss and D.W. Fleck 2010. A new species of *Peropteryx* (Chiroptera: Emballonuridae) from western Amazonia with comments on phylogenetic relationships within the genus. American Museum Novitates 3686: 1–20. doi: [10.1206/691.1](https://doi.org/10.1206/691.1)
- Marques, J. T. et al. 2016. Patterns in the use of rainforest vertical space by Neotropical aerial insectivorous bats: all the action is up in the canopy. Ecography 39: 476–486. doi: [10.1111/ecog.01453](https://doi.org/10.1111/ecog.01453)
- Miller, G. S. 1899. Descriptions of three new free-tailed bats. Bulletin

- American Museum (Natural History) 12: 173–181. <http://hdl.handle.net/2246/776>
- MMA (Ministério do Meio Ambiente). 2014. Fauna brasileira. Lista das espécies quase ameaçadas e com dados insuficientes. Accessed at <http://www.icmbio.gov.br/portal/faunabrasileira/lista-de-especies-dados-insuficientes>, 1 September 2016.
- Ochoa G., J., E. García, S. Caura and J. Sánchez H. 2008. Mamíferos de la cuenca del río Caura, Venezuela: listado taxonómico y distribución conocida. Memoria de la Fundación La Salle de Ciencias Naturales 170: 5–80. <http://www.fundacionlasalle.org.ve/userfiles/Memorial70%205-80.pdf>
- Olson, D.M., E. Dinerstein, E.D. Wikramanayake, N.D. Burgess, G.V.N. Powell, E.C. Underwood, J.A. D'Amico, I. Itoua, H.E. Strand, J.C. Morrison, C.J. Loucks, T.F. Allnutt, T.H. Ricketts, Y. Kura, J.F. Lamoreux, W.W. Wettengel, P. Hedao and K.R. Kassem. 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience* 51: 933. doi: [10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)
- Pirani, F.R., M. Sanchez and F. Pedroni. 2009. Fenologia de uma comunidade arbórea em Cerrado sentido restrito, Barra do Garças, MT. *Acta Botanica Brasilica* 23: 1096–1109. doi: [10.1590/S0102-33062009000400019](https://doi.org/10.1590/S0102-33062009000400019)
- Rawlins, A.M.G. 2011. Dinámica de uso de cuevas por murciélagos cavernícolas de zonas áridas y semiáridas del norte de Venezuela e Islas Vecinas [master's thesis]. Altos de Pipe: Instituto Venezolano de Investigaciones Científicas. 147 pp.
- Reis, N.R., M.N. Fregonezi, A.L. Peracchi and O.A. Shibatta. 2013. *Morcegos do Brasil: guia de campo*. Rio de Janeiro: Technical Books. 252 pp.
- Sampaio, E., B. Lim and S. Peters. 2016. *Peropteryx trinitatis*. The IUCN Red List of threatened species 2016: e.T136790A22035534. Accessed at <http://www.iucnredlist.org/details/136790/0>, 20 September 2016.
- Silva, F.A.M., E.D. Assad and B.A. Evangelista. 2008. Caracterização climática do bioma Cerrado; pp. 61–88, in: S.M. Sano, S.P. Almeida and J.F. Ribeiro (eds.). *Cerrado: ecologia e flora*. Brasília: Embrapa Cerrados.
- Silva, C.R., A.C.M. Martins, I.J. Castro, E. Bernard, E.M. Cardoso, D.S. Lima, R. Gregorin, R.V. Rossi, A.R. Percequillo and K.C. Castro. 2013. Mammals of Amapá state, eastern Brazilian Amazonia: a revised taxonomic list with comments on species distributions. *Mammalia* 77(4): 409–424. doi: [10.1515/mammalia-2012-0121](https://doi.org/10.1515/mammalia-2012-0121)
- Simmons, N.B. and R.S. Voss. 1998. The mammals of Paracou, French Guiana: a Neotropical lowland rainforest fauna. Part 1, bats. *Bulletin American Museum (Natural History)* 237: 2–219. <http://digitallibrary.amnh.org/handle/2246/1634>
- Solari, S. and V. Martínez-Arias. 2014. Cambios recientes en la sistemática y taxonomía de murciélagos Neotropicales (Mammalia: Chiroptera). *Therya* 5: 167–196. doi: [10.12933/therya-14-180](https://doi.org/10.12933/therya-14-180)
- Vianelo, R.L. and A.R. Alves. 2000. *Metereologia básica e aplicações*. Viçosa: UFV. 448p.
- Vizotto, L.D. and V.A. Taddei. 1973. Chave para determinação de quirópteros brasileiros. *Revista da Faculdade de Filosofia, Ciências e Letras de São José do Rio Preto, Boletim de Ciências*: 1–72.
- Zortéa, M., N.A. Bastosb and T.C. Aciolib. 2015. The bat fauna of the Kararaô and Kararaô Novo caves in the area under the influence of the Belo Monte hydroelectric dam, in Pará, Brazil. *Brazilian Journal of Biology* 75: S168–S173. doi: [10.1590/1519-6984.00414BM](https://doi.org/10.1590/1519-6984.00414BM)

**Author contributions:** ASS and DCS collected the data; ASS, DCS and KCF wrote the text.

**Received:** 16 June 2016

**Accepted:** 29 September 2016

**Academic editor:** Guilherme S.T. Garbino