

# New record and distribution extension of the endangered freshwater fish *Cnesterodon hypselurus* (Cyprinodontiformes: Poeciliidae) in the upper Paraná River basin, Brazil

João Fernando Marques da Silva<sup>1\*</sup>, Fernando C. Jerep<sup>2,3</sup> and Sirlei Terezinha Bennemann<sup>1</sup>

1 Universidade Estadual de Londrina (UEL), Centro de Ciências Biológicas, Departamento de Biologia Animal e Vegetal, Laboratório de Ecologia Trófica de Peixes. Rod. Celso Garcia Cid, Pr. 445 Km 380, CEP 86051-980, Londrina, PR, Brazil

2 Universidade Estadual de Londrina (UEL), Centro de Ciências Biológicas, Departamento de Biologia Animal e Vegetal, Museu de Zoologia. Rod. Celso Garcia Cid, Pr. 445 Km 380, CEP 86051-980, Londrina, PR, Brazil

3 Research Associate, Division of Fishes, Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

\* Corresponding author. E-mail: [jfm.bio@gmail.com](mailto:jfm.bio@gmail.com)

**Abstract:** *Cnesterodon hypselurus* is an endangered species from the Upper Paraná River basin, with distribution restricted to three streams with diminished habitat quality. An updated distribution including new locations in the Tibagi River basin is presented, increasing its area of occurrence and providing a new low-elevation record of 728 m. The recognition of these populations suggests that *C. hypselurus* might have a larger distribution in the east-central Paraná State region, changing to vulnerable the previously endangered status of the species.

**Key words:** endemism; habitat degradation; Tibagi River; Itararé River

*Cnesterodon* Garman, 1895 is a monophyletic genus of the family Poeciliidae, of the live-bearing Cyprinodontiformes. An extensive revision of the genus was made by Lucinda (2005) and Lucinda and Reis (2005), recognizing eight valid species, with two more recently described: *Cnesterodon holopteros* Lucinda, Litz & Recuero, 2006 and *Cnesterodon pirai* Aguilera, Mirande & Azpeli-cueta, 2009. These 10 species inhabit coastal and inland drainages in Argentina, southern Brazil, Paraguay and Uruguay, restricted to headwaters and high elevation regions of these basins (Lucinda 2003, 2005; Lucinda et al. 2006).

*Cnesterodon hypselurus* Lucinda & Garavello, 2001:129 is endemic to streams in Paraná State, Brazil, and, according to the literature (Lucinda and Garavello 2001; Shibatta et al. 2002; Lucinda 2003, 2005; Lucinda and Reis 2005; Langeani et al. 2007; Meredith et al. 2010;

Pollux et al. 2014), its distribution is restricted to the Cilada and Lambari Rivers, both tributaries of the Capivari River, part of the Itararé River basin, and a nameless tributary of the Guaricanga River, of the Tibagi River basin. Both the Tibagi and Itararé River basins are part of the the Paranapanema drainage, flowing from South to North, approximately 550 km and 300 km in length respectively. Shibatta et al. (2002) proposed that *C. hypselurus* was endangered in the Tibagi River basin, however only recently the species was included in the Brazilian list of endangered species (MMA 2014). According to the IUCN Red List Categories and Criteria, it was classified as EN (B1ab(iii)), indicating that *C. hypselurus* is facing a very high risk of extinction, with a range of less than 5,000 km<sup>2</sup>, and known from no more than five locations, with continuing decline of habitat quality (IUCN 2001; ICMBIO 2014). Herein we report new locations for *C. hypselurus* in the Tibagi River basin, increasing the known localities for this species and changing its threat category according to the IUCN parameters (IUCN 2001).

Field studies were conducted between February and November 2014, in first and second order stretches of eight streams in the middle Tibagi River basin, property of Klabin S.A. cellulose and paper company, east-central Paraná State. This area has been classified as possessing extremely high biological importance and high priority for conservation (MMA 2007). A total of 38 specimens of *Cnesterodon hypselurus* (Figures 1 and 2) were sampled in one of the eight streams, in a first order stretch of the Arroio do Gica stream (24°12'32.78" S, 050°36'35.65" W, 728 m), tributary of the Tibagi River, Telêmaco Borba (Figure 3).




**Figure 1.** Specimens of *Cnesterodon hypselurus* showing coloration in life in male (above) and female (below). Specimens not preserved. Photos by F.C. Jerep.



**Figures 2.** Coloration of alcohol preserved specimens of *Cnesterodon hypselurus*. Mature male, MZUEL 12699, 24.1 mm SL (above) and female, MZUEL 12640, 34.5 mm SL (below). Photos by J.F.M. Silva.

Specimens were collected in pool habitats (according to Rincón 1999) using a seine net (2 mm mesh) and sieves (2 mm mesh) for one hour, under SISBIO permit #42829-1. Fishes were anesthetized with clove oil (Anderson et al. 1997; Lucena et al. 2013), fixed in 10% formalin for 48 hours, washed in running water and preserved in 70% ethanol (Uieda and Castro 1999). Voucher specimens analyzed in this study are deposited




**Figure 3.** Updated geographic distribution of *Cnesterodon hypselurus*. Additional sites in red represent seven streams sampled during the present study. Source: adapted from Miranda (2005), SUDERHSA (2007) and IBGE (2013).

at the Museu de Zoologia da Universidade Estadual de Londrina (MZUEL) under the following catalogue numbers: MZUEL 12640, 12699, 12752 and 12798.

Two other species were collected in the same stream: *Astyanax paranae* Eigenmann, 1914 and *Trichomycterus davisii* (Haseman, 1911). *Astyanax paranae* inhabits the same pool habitat where *C. hypselurus* was collected, while *T. davisii* was collected strictly in riffle habitat.

The specimens were identified as *Cnesterodon hypselurus* following the identification key and diagnosis proposed by Lucinda (2005) and comparisons with the paratypes (MZUEL 1600), specimens of congeners (Lucinda et al. 2006; Aguilera et al. 2009), and specimens identified by P.H.F. Lucinda (MZUEL 1727 and MZUEL 1728). The updated distribution map of the species (Figure 3; Table 1) was based on data from literature (Lucinda and Garavello 2001; Shibatta et al. 2002; Lucinda 2003, 2005; Lucinda and Reis 2005; Langeani et al. 2007; Meredith et al. 2010; Pollux et al. 2014) and from the SpeciesLink database (CRIA 2015). Vouchers from new localities found in SpeciesLink were examined to verify identification.

Sites where specimens were collected were included in a GIS base map containing limits of the Paraná State Watersheds (SUDERHSA 2007), Digital Elevation Model (Miranda 2005) and hydrography (IBGE 2013). The new extent of occurrence was calculated by the area of a



**Figure 4.** Distal tip of the gonopodium of a mature male of *Cnesterodon hypselurus*, MZUEL 12699, 25.2 mm SL. R3, R4a, R4p, R5a, R5p are sections of the anal-fin rays, modified in gonopodial rays. Left side, lateral view. Photo by F.C. Jerep.

minimum convex polygon, which is the smallest polygon with no internal angle exceeding 180° and containing all sites of occurrence (IUCN 2001).

According to Lucinda (2005), *Cnesterodon hypselurus* can be identified by two autapomorphies, the presence of a dark longitudinal band along the flank in preserved material (Figure 2), and the bony style of ray 3 of the gonopodium of fully developed males curly-bracket shaped, with a constriction located at two-thirds of its length (Figure 4). In addition, the combination of the dark brown blotches along body sides forming bars and covering more than four scales in a transverse row, the presence of a dark band along flank and depth of the caudal peduncle in adult males (15.5–17.8% SL) differentiate this species from its congeners. A variation in the number of segments at the tip of ray 4a of the gonopodium of mature males was observed, with four to six segments before the claw-shaped bony tip. The specimens also presented some variation in the coloration pattern (Figure 2) in comparison to the original description and specimens from other localities. The longitudinal dark stripe may be slightly fainter in some specimens, and the dark vertical bars might cover up to three scales of a transverse row instead of four, as mentioned in the original description (Lucinda and Garavello 2001). Notwithstanding, the variation herein observed is similar to the color pattern of the examined type material and photographs (Lucinda and Garavello 2001, fig. 5; Lucinda 2005, fig. 7).

The pool habitat where the specimens were collected at Arroio do Gica stream is 3.5–4.0 m wide, 40–59 cm deep, and 0.04–0.06 m/s current speed. The bottom of the stream was composed of rocks, leaf litter and non-consolidated clay (Figure 5). Riparian forest covered the right margin while the left margin was bordered by commercial cultivation of *Pinus* sp. that was harvested during the studies. The water temperature ranged from 15.4 to 19.8°C, pH 6.3 to 7.7, conductivity 15.2 to 25.1 µS/cm and dissolved oxygen 7.6 to 8.8 mg/L (80.5–88.1%).

**Table 1.** Known localities of *Cnesterodon hypselurus* in Paraná State and vouchers. Elevation in meters above sea level; N = number of specimens.

| Locality                                | Drainage | Municipality   | Geographic coordinates          | Altitude | Catalog number (N)  |
|---|----------|----------------|---------------------------------|----------|---|
| Cilada River <sup>7,8</sup>             | Itararé  | Jaguariaíva    | 24°19'58.03" S, 049°47'30.22" W | 1,078 m  | MCP12593 (31) <sup>1,4,5</sup> , MCP22743 (1) <sup>1,3,4,6</sup> , MCP22744 (9) <sup>1,4,5</sup> , MHNCI5890 (10) <sup>1,4</sup> , UMMZ215219 (20) <sup>1,4</sup> , USNM356395 (2) <sup>1,4</sup> , MZUEL1728 (122) <sup>1,4</sup> , MZUSP85935.0 (100) <sup>9</sup> , UFRGS18402 (32) <sup>9</sup> |
| Lambari River                           | Itararé  | Jaguariaíva    | 24°22'24.28" S, 049°50'23.47" W | 1,136 m  | MCP12602 (7) <sup>1,4</sup> , MCP22745 (5) <sup>1,4</sup> , MHNCI5892 (3) <sup>1,4</sup>  |
| Cinco Reis River                        | Itararé  | Jaguariaíva    | 24°17'27.70" S, 049°44'54.30" W | 941 m    | MZUSP85936.0 (3) <sup>9</sup>   |
| Guaricanga River tributary <sup>2</sup> | Tibagi   | Piraí do Sul   | 24°25'60.00" S, 050°01'60.00" W | 1,110 m  | MZUEL1600 (14) <sup>1,4</sup> , MZUEL1727 (10) <sup>1,4</sup> , MCP23326 (10) <sup>4</sup>  |
| Arroio do Gica stream                   | Tibagi   | Telêmaco Borba | 24°12'32.78" S, 050°36'35.65" W | 728 m    | MZUEL12640 (14), MZUEL12699 (14), MZUEL12752 (6), MZUEL12798 (4)  |
| Lajeado Maria Leme stream               | Tibagi   | Castro         | 24°45'34.00" S, 050°11'60.00" W | 1,102 m  | NUP15659 (10) <sup>9</sup>  |
| Pitangui River tributary                | Tibagi   | Carambeí       | 24°52'16.00" S, 050°14'10.00" W | 861 m    | NUP15828 (6) <sup>9</sup>   |
| Verde River tributary                   | Tibagi   | Ponta Grossa   | 25°03'04.00" S, 050°06'11.02" W | 904 m    | DZSJRP20376 (168) <sup>9</sup>  |

<sup>1</sup>Lucinda and Garavello 2001; <sup>2</sup>Shibatta et al. 2002; <sup>3</sup>Lucinda 2003; <sup>4</sup>Lucinda 2005; <sup>5</sup>Lucinda and Reis 2005; <sup>6</sup>Langeani et al. 2007; <sup>7</sup>Meredith et al. 2010; <sup>8</sup>Pollux et al. 2014; <sup>9</sup>CRIA 2015.



**Figure 5.** Photograph of the new record locality of *Cnesterodon hypselurus* in Arroio do Gica stream, Telêmaco Borba, PR, Brazil. Photo by S.T. Benemann.

The stream habitat where the species has been previously recorded were not described in the literature, although *C. hypselurus* lives in clear waters with little flow in the Tibagi basin (Shibatta et al. 2002). The other streams sampled during the study did not have pools with similar characteristics as found in the Arroio do Gica, indicating a possible habitat requirement of this species. The elevation of the previous known locations ranged from 861 to 1,136 m above sea level, but Arroio do Gica stream is at 728 m above sea level, making it the lowest site reported for the species.

New records from the SpeciesLink database indicated four additional locations for the species (CRIA 2015), for a total of eight different streams (Figure 3, Table 1). The updated extent of occurrence for *C. hypselurus* was found to be 3,989 km<sup>2</sup>. Although the updated area of distribution still fits the criteria for EN(B1) (extent of occurrence less than 5,000 km<sup>2</sup>), the occurrence in eight different locations invalidates the criteria EN(B1a) (species known to exist at no more than five locations). Based on the abovementioned, the category of the species changes from endangered (EN) to vulnerable (VU) according to the VU(B1ab(iii)) criteria: extent of occurrence restricted to less than 20,000 km<sup>2</sup> and known to exist at no more than ten locations with continuing decline of habitat quality.

The vulnerable category still indicates that the species is facing a high risk of extinction. Additional data about reduction in population size, number of mature individuals in the population and probability of extinction in the wild in five generations are still needed for the species, and can only be achieved through standardized sampling in the locations for an extended period of time.

The present study increases the known distribution of *Cnesterodon hypselurus* to eight localities; changing

the conservation status of the species according to the IUCN (2001) parameters to VU(B1ab(iii)). The population recorded herein is relatively small and should be regularly monitored. On the other hand, the discovery of *C. hypselurus* in this stream suggests that there may be additional undiscovered populations in the Tibagi, Cinzas and Itararé River basins. We encourage targeted inventories efforts in streams with similar habitat characteristics as observed in Arroio do Gica stream throughout this region, as well as a phylogeographic study to determine the processes involved in the geographic distribution of this species.

## ACKNOWLEDGEMENTS

We thank all the members of the Museu de Zoologia (UEL) who helped us with the fieldwork: A. Garcia, A. Souza, E. Santana, L. Jarduli, R. Nascimento and T. Lizarelli. We are especially grateful to Klabin S.A. and Sergio Adão Filipaki for the support given during field activities. JFMS is supported by a scholarship from CAPES/FA (11/2013) Doctorate Program. FCJ is supported by CNPq (453850-2014/6). We are thankful to Francisco Langeani (DZSJRP), Manoela Marinho (MZUSP), Weferson Júnio da Graça (UEM) and Vinícius Abilhoa (MHNCI) for providing photographs and specimens.

## LITERATURE CITED

- Aguilera G., J.M. Mirande and M.D.L.M. Azpelicueta. 2009. A new species of *Cnesterodon* (Cyprinodontiformes: Poeciliidae) from a small tributary of arroyo Cuñá-Pirú, río Paraná basin, Misiones, Argentina. *Zootaxa* 2195: 34–42.
- Anderson, W.G., R.S. McKinley and M. Colavecchia. 1997. The use of clove oil as an anesthetic for rainbow trout and its effects on swimming performance. *North American Journal of Fisheries Management* 17(2): 301–307. doi: 10.1577/1548-8675(1997)017<0301:TUOC OA>2.3.CO;2
- CRIA. [2015]. SpeciesLink network. Centro de Referência em Informação Ambiental. Accessed at <http://www.splink.org.br>, 29 July 2015.
- IBGE. 2013. Base cartográfica contínua do Brasil na escala 1:250.000. Version 1.0. Instituto Brasileiro de Geografia e Estatística. Accessed at [ftp://geofp.ibge.gov.br/mapeamento\\_sistemico/base\\_vetorial\\_continua\\_escala\\_250mil](ftp://geofp.ibge.gov.br/mapeamento_sistemico/base_vetorial_continua_escala_250mil), 29 April 2015.
- ICMBIO. 2014. Espécies ameaçadas, lista 2014. Instituto Chico Mendes de Conservação da Biodiversidade. Accessed at <http://www.icmbio.gov.br/portal/biodiversidade/fauna-brasileira/lista-de-especies.html>, 29 April 2015.
- IUCN. 2001. IUCN red list categories and criteria. Version 3.1. International Union for Conservation of Nature. Accessed at <http://www.iucnredlist.org/technical-documents/categories-and-criteria/2001-categories-criteria>, 29 April 2015.
- Langeani, F., R.M.C. Castro, O.T. Oyakawa, O.A. Shibatta, C.S. Pavanelli and L. Casatti. 2007. Diversidade da ictiofauna do Alto Rio Paraná: composição atual e perspectivas futuras. *Biota Neotropica* 7(3): 181–197. doi: 10.1590/S1676-06032007000300020
- Lucena, C.A.S., B.B. Calegari, E.H.L. Pereira and E. Dallegrove. 2013. O uso de óleo de cravo na eutanásia de peixes. *Boletim Sociedade Brasileira de Ictiologia* 105: 20–24. <http://www.sbi.bio.br/sbi/pdfs/boletins/BoletimSBI105.pdf>

- Lucinda, P.H.F. 2003. Family Poeciliidae; pp. 558–584, in: R.E. Reis, S.O. Kullander and C.J. Ferraris Jr. (eds.). Check list of the freshwater fishes of South and Central America. Porto Alegre: EDIPUCRS.
- Lucinda, P.H.F. 2005. Systematics of the genus *Cnesterodon* Garman, 1895 (Cyprinodontiformes: Poeciliidae: Poeciliinae). *Neotropical Ichthyology* 3(2): 259–270. doi: [10.1590/S1679-62252005000200003](https://doi.org/10.1590/S1679-62252005000200003)
- Lucinda, P.H.F. and J.C. Garavello. 2001. Two new species of *Cnesterodon* Garman, 1895 (Cyprinodontiformes: Poeciliidae) from the upper rio Paraná drainage. *Comunicações do Museu de Ciências e Tecnologia da PUCRS, Série Zoologia* 13(2): 119–138.
- Lucinda, P.H.F. and R.E. Reis. 2005. Systematics of the subfamily Poeciliinae Bonaparte (Cyprinodontiformes: Poeciliidae), with an emphasis on the tribe Cnesterodontini Hubbs. *Neotropical Ichthyology* 3(1): 1–60. doi: [10.1590/S1679-62252005000100001](https://doi.org/10.1590/S1679-62252005000100001)
- Lucinda, P.H.F., T. Litz and R. Recuero. 2006. *Cnesterodon holopteros* (Cyprinodontiformes: Poeciliidae: Poeciliinae), a new species from the Republic of Uruguay. *Zootaxa* 1350: 21–31.
- Meredith, R.W., M.N. Pires, D.N. Reznick and M.S. Springer. 2010. Molecular phylogenetic relationships and the evolution of the placenta in *Poecilia* (*Micropoecilia*) (Poeciliidae: Cyprinodontiformes). *Molecular Phylogenetics and Evolution* 55(2): 631–639. doi: [10.1016/j.ympev.2009.11.006](https://doi.org/10.1016/j.ympev.2009.11.006)
- Miranda, E. E. 2005. Brasil em relevo. Embrapa Monitoramento por Satélite. Accessed at <http://www.relevobr.cnpm.embrapa.br>, 29 April 2015.
- MMA (Ministério do Meio Ambiente). 2007. Áreas prioritárias para a conservação, uso sustentável e repartição de benefícios da biodiversidade brasileira: atualização — Portaria MMA no. 09 de 23 de janeiro de 2007. Brasília: MMA/SBF. 300 pp.
- MMA (Ministério do Meio Ambiente). 2014. Lista nacional oficial de espécies da fauna ameaçadas de extinção — peixes e invertebrados aquáticos. Portaria no. 445 de 17 de dezembro de 2014. Diário Oficial da União 245: 126–130.
- Pollux, B.J.A., R.W. Meredith, M.S. Springer, T. Garland and D.N. Reznick. 2014. The evolution of the placenta drives a shift in sexual selection in livebearing fish. *Nature* 513(7517): 233–236. doi: [10.1038/nature13451](https://doi.org/10.1038/nature13451)
- Rincón, P.A. 1999. Uso do micro-habitat em peixes de riachos: métodos e perspectivas; pp. 23–90, in: E.P. Caramaschi, R. Mazzoni and P.R. Peres-Neto (eds.). *Ecologia de Peixes de Riachos. Série Oecologia Brasiliensis, volume VI*. Rio de Janeiro: PPGE-UFRJ.
- Shibatta, O.A., M.L. Orsi, S.T. Bennemann and A.T. Silva-Souza. 2002. Diversidade e distribuição de peixes na bacia do rio Tibagi; pp. 403–423, in: M.E. Medri, E. Bianchini, O.A. Shibatta and J.A. Pimenta (eds.). *A bacia do rio Tibagi*. Londrina: M.E. Medri.
- SUDERHSA. 2007. Bacias hidrográficas do Paraná. Superintendência de Desenvolvimento de Recursos Hídricos e Saneamento Ambiental. Accessed at [http://www.aguasparana.pr.gov.br/arquivos/File/DADOS%20ESPACIAIS/Bacias\\_Hidrograficas\\_Parana.zip](http://www.aguasparana.pr.gov.br/arquivos/File/DADOS%20ESPACIAIS/Bacias_Hidrograficas_Parana.zip), 29 April 2015.
- Uieda, V.S. and R.M.C. Castro. 1999. Coleta e fixação de peixes de riacho; pp. 1–22, in: E.P. Caramaschi, R. Mazzoni and P.R. Peres-Neto (eds.). *Ecologia de Peixes de Riachos. Série Oecologia Brasiliensis, volume VI*. Rio de Janeiro: PPGE-UFRJ.

**Author contributions:** JFMS collected the data and wrote the text; FCJ and STB also contributed to the text writing; JFMS and FCJ identified specimens and assessed morphological variation; and FCJ, STB and JFMS collected specimens and prepared the photographs.

**Received:** 11 June 2015

**Accepted:** 5 December 2015

**Academic editor:** Marina Loeb