

# Faunal richness and the checklist of Indian mosquitoes (Diptera: Culicidae)

D. R. Bhattacharyya<sup>1</sup>, A. R. Rajavel<sup>2</sup>, R. Natarajan<sup>2</sup>, P. K. Mohapatra<sup>1</sup>, P. Jambulingam<sup>2</sup>, J. Mahanta<sup>1</sup> and A. Prakash<sup>1,3\*</sup>

1 Regional Medical Research Centre, NE (Indian Council of Medical Research), Post Box No. 105, Dibrugarh – 786 001, Assam, India.

2 Vector Control Research Centre (Indian Council of Medical Research), Indiranagar, Puducherry – 605 006, Tamilnadu, India.

3 National Institute for Research in Environmental Health (Indian Council of Medical Research), Kamla Nehru Hospital Building, Bhopal – 462 001, Madhya Pradesh, India.

\* Corresponding Author. E-mail: [anilprakashin@yahoo.co.in](mailto:anilprakashin@yahoo.co.in)

**ABSTRACT:** A review of published studies revealed that the Indian mosquito fauna comprises 393 species in 49 genera and 41 subgenera. The subfamily Anophelinae contains 61 species in one genus followed by Culicinae with 332 species in 11 tribes and 48 genera. The tribe Aedini (subfamily Culicinae) contains the highest number of species (176 species in 33 genera and two groups of *incertae sedis*; i.e., “*Aedes*” *sensu auctorum* and “*Ochlerotatus*” *sensu auctorum*). With the recent taxonomic changes in tribe Aedini, the Indian mosquito genera have gone up from 22 to 49. Changes to the Indian Aedini fauna subsequent to the reclassification of tribe Aedini are discussed. A total of 31 species are currently recognized in India for transmitting various mosquito-borne agents of human diseases. A checklist for the Indian mosquito species is presented and the need for a comprehensive study is emphasized.

DOI: 10.15560/10.6.1342

## INTRODUCTION

Mosquitoes, belonging to family Culicidae and order Diptera, are a large group of insects present throughout the temperate and tropical regions and even beyond the Arctic Circle of the world (Harbach 2007). The Oriental Region, which includes India, is regarded as one of the richest biogeographic regions for mosquitoes of the world, along with the Neotropics (Gaston and Hudson 1994). At present, a total of 3,540 recognized mosquito species, divided between two subfamilies and 112 genera, are recorded in the world (Harbach 2014). A three to five-fold increase of the present numbers can be foreseen with more discoveries and naming of sibling species mainly in *Anopheles*, as a result of application of DNA-based methods (Harbach 2007). India is ranked fifth in terms of mosquito biodiversity after Brazil, Indonesia, Malaysia and Thailand (Foley *et al.* 2007). After the monumental works of Christophers (1933) and Barraud (1934), not many comprehensive biosystematics studies of Indian Culicidae have been undertaken. Since that time, many changes have taken place in the mosquito taxonomic literature. However, in the absence of any up-to-date monograph on Indian Culicidae, entomologists in India are still compelled to use these two books. In this context, as mentioned by Reuben *et al.* (1993), it needs to be emphasized that publications of the Southeast Asian Mosquito Project are very useful for the identification of Indian mosquito species. Berlin (1972) rightly proposed for a systematic study of Indian Culicidae because of increasing mosquito-borne diseases, changing ecology and advances in mosquito systematics. The online systematic catalog of Culicidae listed 356 species in India (Gaffigan *et al.* 2014). Although fewer in

numbers, since the 1980s some faunistic surveys carried out in various parts of India detected many species which are new additions to the mosquito fauna of India. It seems that some of these records did not reach the mosquito catalog. As such, an up-to-date check list of mosquitoes of India is still not available, and the actual number of species present in India, has to be approximated. Moreover, with the advancement of molecular biology, many researchers from non-entomology backgrounds are now working with mosquitoes, and there is confusion about the informal taxonomic series, groups and subgroups to which mosquito species belong. Confusion seems to have increased after the changes made to the classification of tribe Aedini (Reinert *et al.* 2004, 2006, 2008, 2009) as journals are using two systems of classification. Reclassification of the tribe Aedini brought major changes raising the number of genera in this tribe from 11 to 81 along with changes to the spelling of names of some species in accordance with provisions of the *International Code of Zoological Nomenclature*. Hence, a checklist seems essential for the entomologists and public health personnel working in India to reflect changes to nomenclature to promote familiarity with the new names. Considering this, an up-to-date check list of the mosquito species of India based on the current classification is provided. The present status of the mosquito fauna of India and need for future work are also discussed.

## MATERIALS AND METHODS

Information was collected from two online resource databases, namely systematic catalog of Culicidae and mosquito taxonomic inventory and published materials

mainly since the 1980s (Bhattacharyya *et al.* 2000, 2002, 2003, 2004a, 2004b, 2005, 2007, 2008, 2009; Khan *et al.* 1998; Rajavel and Natarajan 2006, 2011; Rajavel *et al.* 1998, 2004, 2005a, 2005b, 2005c, 2005d, 2011; Rajput and Singh 1987a, 1987b, 1987c, 1989, 1990; Reuben *et al.* 1993; Tewari *et al.* 1987; Tewari and Hiriyan 1991, 1992, 1995). The systematic catalog of Culicidae provides information on country occurrence records of mosquito species. The mosquito taxonomic inventory contains up-to-date information on world Culicidae. The internal classification of genera *Anopheles*, *Culex*, *Verrallina* and *Uranotaenia* are available in the website of mosquito taxonomic inventory, which is followed here for the Indian mosquito fauna belonging to these genera. Recent abbreviations of generic and subgeneric names provided by Reinert (2009a) are followed here. The authorship of generic and species names is provided in the check list (Appendix 1).

## RESULTS AND DISCUSSION

Records indicate that the Indian mosquito fauna includes 393 species divided among 49 genera and 41 subgenera. Subfamily Anophelinae contains 61 species in one genus followed by subfamily Culicinae with 332 species in 11 tribes and 48 genera. Tribe Aedini of subfamily Culicinae contains the highest number of species (176 species in 33 genera and two groups of *incertae sedis*, i.e., "*Aedes*" *sensu auctorum*, "*Ochlerotatus*" *sensu auctorum*) (Table 1). The check list of the Indian Culicidae is presented as Appendix 1. In India, 31 species are currently recognized for transmitting various mosquito-borne pathogens; these are listed in Table 2.

**Subfamily Anophelinae:** Subfamily Anophelinae has three genera globally. Indian species are confined to genus *Anopheles*, with 61 formally named species divided between subgenera *Anopheles* and *Cellia*. However, the number of species will increase with the naming or recognition of synonymous names of sibling species in several groups or complexes. In India, subgenus *Anopheles* contains 26 species. In this subgenus, the assemblage of species in two most important groups (Hyrceanus and Barbirostris Groups) is still uncertain in India. The Hyrceanus Group is a highly complex group, which includes 30 closely related species distributed widely in the Oriental and Palaearctic regions, with some species playing important role in transmission of malaria and filariasis (Ma and Xu 2005). This Group was also incriminated as a vector of Japanese encephalitis in India and neighboring countries. Only seven species of this Group were recorded from India in comparison to 25 species recorded in China (Ma *et al.* 1998, 2000a, 2000b). Morphological identification of these species in adult stage seems to be very difficult or impossible unless accompanied by associated immature skins (Harrison and Scanlon 1975). Many anopheline surveys carried out in India recorded *An. nigerrimus* of Hyrceanus Group to be the most predominant, seemingly because their identification was based on adult stage (Malhotra *et al.* 1987; Nagpal and Sharma 1987). However, surveys carried out in north-east India and Western Ghats of South India, where species identification was done using adult as well as associated larval and pupal characters, recorded *An. peditaeniatus* to be the most

prevalent and *An. nigerrimus* as rather uncommon species (Khan *et al.* 1998; Tewari *et al.* 1987). Immature surveys carried out in upper Brahmaputra valley detected *An. crawfordi*, *An. paraliae*, *An. peditaeniatus* and *An. sinensis* of Hyrceanus Group from the north-east India, but did not collect any specimen of *An. nigerrimus* (Khan *et al.* 1998). A few specimens identified as *An. paraliae* in these surveys were collected as immatures from a forest fringe area of Assam. Earlier considered as a subspecies of *An. lesteri*, *An. paraliae* was later elevated to species status by Harrison *et al.* (1990), but more recently it has been synonymized with *lesteri* by Taai *et al.* (2013), hence it is listed as *lesteri* here. However, distribution of this species is restricted to low elevation coastal areas of Malaysia, Brunei, Vietnam and Thailand (Harrison and Scanlon 1975). This species is also easily misidentified with *An. pursati*. Hence, record of this species in India needs further investigation, incorporating DNA-based analysis similar to China, where, ITS2 marker of r-DNA was used to differentiate member species of *An. hyrceanus* group (Ma *et al.* 1998, 2000a, 2000b). Recently, ITS2 sequencing of some specimens of Hyrceanus Group confirmed the presence of *An. crawfordi*, *An. peditaeniatus* and *An. sinensis* in the north-east India (Regional Medical Research Centre, Dibrugarh, Assam, India; unpublished data). Similarly Barbirostris Group which includes 11 species globally also has uncertain distribution records for most of the species, particularly in Indonesia and Indian sub-region, mainly because of identification difficulties (Harrison and Scanlon 1975). Four species were recognized in this group from India. *An. hodgkini* was detected from Assam during larval collections (Khan *et al.* 1998).

**Subgenus *Cellia*** containing 35 species encompasses all important vectors of human malaria in India. Recently, two new species have been described from this subgenus from India. *An. pseudosundaicus* belonging to Pyrethophorus Series was described from coastal areas of Kerala (Tyagi *et al.* 2009). Another species belonging to Maculatus Group of the Neocellia Series named as *An. krishnai* was described (Sathe and Jagtap 2012). However, Harbach (2014) noted the lack of credible evidence to support the recognition of this proposed species. Hence, *krishnai* is regarded as *nomen nudum* and is not included in our list of Indian *Anopheles*. *Anopheles dthali* and *Anopheles superpictus* have been known only from extreme north-west (Baluchistan) which is not part of India. They have also not been recorded subsequently in India and therefore not included in the present checklist.

Several anopheline taxa recognized earlier to be medically important are now found to be complexes or groups of morphologically indistinguishable species (Rattanaarithikul *et al.* 2006). Some important complexes and groups recorded in India are, *An. annularis* (species A and B), *An. culicifacies* (species A, B, C, D and E), *An. dirus* (species X), *An. fluviatilis* (species S, T, U and V), *An. subpictus* (species A, B, C, and D) and the *An. sundaicus* (cytotype D) complexes (Walton *et al.* 1999; WHO 2007). With the advances in mosquito taxonomy several species belonging to such groups or complexes are now formally named. *An. maculatus*, recognized as one such species group, includes nine formally named species (Harbach 2014; Somboon *et al.* 2011). A recent study revealed the

**TABLE 1** Member species of Culicidae in India.

SUBFAMILY	TRIBE	GENUS	SUBGENUS	NUMBER OF SPECIES
Anophelinae		<i>Anopheles</i>	<i>Anopheles</i>	26
			<i>Cellia</i>	35
Culicinae	Aedeomyiini	<i>Aedeomyia</i>	<i>Aedeomyia</i>	1
	Aedini	" <i>Aedes</i> " <i>sensu auctorum</i>		4
		<i>Aedimorphus</i>		15
		<i>Armigeres</i>	<i>Armigeres</i>	9
			<i>Leicesteria</i>	11
		<i>Ayurakitia</i>		1
		<i>Bruceharrisonius</i>		4
		<i>Christophersomyia</i>		4
		<i>Canraeodes</i>		2
		<i>Collessius</i>	<i>Collessius</i>	3
			<i>Alloeomyia</i>	1
		<i>Danielsia</i>		2
		<i>Dendroskusea</i>		5
		<i>Downsiomyia</i>		6
		<i>Edwardsaedes</i>		1
		<i>Finlaya</i>		2
		<i>Fredwardsius</i>		1
		<i>Gilesius</i>		1
		<i>Himalaius</i>		2
		<i>Heizmannia</i>	<i>Heizmannia</i>	11
			<i>Mattinglyia</i>	2
		<i>Hopkinsius</i>	<i>Yamada</i>	1
		<i>Hulecoeteomyia</i>		5
		<i>Jihlienius</i>		1
		<i>Kenknightia</i>		2
		<i>Lorrainea</i>		2
		<i>Mucidus</i>	<i>Mucidus</i>	3
		<i>Neomelaniconion</i>		1
		<i>Ochlerotatus</i>		1
		" <i>Ochlerotatus</i> " <i>sensu auctorum</i>	" <i>Finlaya</i> " <i>sensu auctorum</i>	5
		<i>Paraedes</i>		4
		<i>Petermattinglyius</i>	<i>Petermattinglyius</i>	2
		<i>Phagomyia</i>		11
		<i>Rhinoskusea</i>		3
		<i>Scutomyia</i>		1
		<i>Stegomyia</i>	<i>Actinothrix</i>	2
			<i>Heteraspidion</i>	2
			<i>Huangmyia</i>	2
			<i>Stegomyia</i>	1
			<i>Xyele</i>	1
			<i>Without subgenus</i>	11
		<i>Tewarius</i>		3
		<i>Udaya</i>		2
		<i>Verrallina</i>	<i>Harbachius</i>	4
			<i>Neomacleaya</i>	16
			<i>Verrallina</i>	3
	Culicini	<i>Culex</i>	<i>Culex</i>	23
			<i>Culiciomyia</i>	9
			<i>Eumelanomyia</i>	10
			<i>Lophoceraomyia</i>	28
			<i>Maillotia</i>	1
			<i>Oculeomyia</i>	6
		<i>Lutzia</i>	<i>Metalutzia</i>	4
	Culisetini	<i>Culiseta</i>	<i>Allotheobaldia</i>	1
			<i>Culiseta</i>	2
	Ficalbiini	<i>Ficalbia</i>		1
		<i>Mimomyia</i>	<i>Etorleptomyia</i>	1
			<i>Mimomyia</i>	4
			<i>Ingramia</i>	1
	Hodgesini	<i>Hodgesia</i>		1
	Mansoniini	<i>Coquilletidia</i>	<i>Coquilletidia</i>	3
		<i>Mansonia</i>	<i>Mansonioides</i>	4
	Orthopodomyiini	<i>Orthopodomyia</i>		5
	Sebethini	<i>Malaya</i>		2
		<i>Topomyia</i>	<i>Suaymyia</i>	1
			<i>Topomyia</i>	3
		<i>Tripteroides</i>	<i>Rachionotomyia</i>	5
			<i>Tripteroides</i>	3
	Toxorhynchitini	<i>Toxorhynchites</i>	<i>Toxorhynchites</i>	9
	Uranotaeniini	<i>Uranotaenia</i>	<i>Pseudoficalbia</i>	15
			<i>Uranotaenia</i>	13
Total of each taxonomic level				
2	11	49 + 2 <i>sensu auctorum</i>	41 + 1 <i>sensu auctorum</i>	393

presence of six member species of the Maculatus Group in the north-east India, with detection of *An. rampae* constituting a new country record (Singh *et al.* 2012). Such studies may be extended to other areas where Maculatus Group is present. Similarly, species X of the Dirus Complex, reported earlier from Yunnan province of China, was also detected in Haflong area of Assam in the north-east India (Prakash *et al.* 2010).

**Subfamily Culicinae:** Subfamily Culicinae is comprised of 332 species divided into 11 tribes and 48 genera in India.

**Tribe Aedeomyiini:** A single genus *Aedeomyia* is included in this tribe. This genus is not listed in the mosquito catalog under the Indian mosquito fauna. However, Tyson (1970) reported Andaman Islands as one of the distribution localities for *Aedeomyia catastica*. This species was also reported from mainland India from Assam and Western Ghats (Khan *et al.* 1998; Reuben *et al.* 1993) and in the mangrove forests of Sunderbans, West Bengal (Rajavel *et al.* 2005a).

**Tribe Aedini:** In the earlier classification the tribe Aedini contained 11 genera (Knight and Stone 1977 and its supplements). Recent reclassification of this tribe has raised the number of genera to 81 (Reinert *et al.* 2004, 2006, 2008, 2009). Reinert (1999, 2000a) earlier resurrected *Verrallina* and *Ayurakitia* to generic rank. Revision of tribe Aedini resulted in the elevation of many former subgenera of *Aedes* to generic status (Reinert *et al.* 2004). *Downsiomyia* was elevated to genus level from synonymy with *Finlaya*. Further examination and revision of *Finlaya* and allied taxa and *Ochlerotatus* and allied taxa (Reinert *et al.* 2006, 2008), followed by comprehensive phylogenetic study of tribe Aedini (Reinert *et al.* 2009) resulted in recognition of more genera and rearrangement of taxa.

As per the current classification, Indian fauna of tribe Aedini includes 176 species in 33 genera and two groups of *incertae sedis* species ("*Aedes*" *sensu auctorum*-4 and "*Ochlerotatus*" *sensu auctorum*-5). Indian species belonging to subgenus *Finlaya* genus *Aedes* are now placed in 12 genera (*Bruceharrisonius*, *Collessius*, *Downsiomyia*, *Danielsia*, *Finlaya*, *Gilesius*, *Hopkinsius*, *Hulecoeteomyia*, *Himalaius*, *Jihlienius*, *Ochlerotatus* and *Phagomyia*). Similarly species of the subgenus *Diceromyia* of India are now placed in three genera *Dendroskusea*, *Petermattinglyius* and *Tewarius*. Illustrated keys of these new genera present in Thailand were provided by Rattanaarithikul *et al.* (2010). Further details are available in the website of mosquito taxonomic inventory. Kaur (2003) provided updated distribution maps of Indian Aedini fauna.

**Genus Aedimorphus:** Earlier recognized as a subgenus of genus *Aedes*, *Aedimorphus* is now elevated to generic rank (Reinert *et al.* 2009). Mostly Oriental in distribution, of the 67 species recognized in this genus, 15 species were recorded from India.

**Genus Armigeres:** Distributed mainly in the Oriental region and also in the Palaearctic and Australasian regions, the genus *Armigeres* is represented by 20 species in India, between two subgenera *Armigeres* and *Leicesteria*. Rajput and Singh (1987c) reported the detection of *Ar. dolichocephalus* from Manipur state. *Ar. joloensis*, a rare mosquito species, was detected from upper Assam

(Bhattacharyya *et al.* 2000). Recently a new species *Ar. mahantai* collected as immatures from endemic pitcher plant *Nepenthes khasiana* of Meghalaya state in the north-east India was described (Bhattacharyya *et al.* 2009). *Ar. pallithorax*, described from Yunnan, China was detected from Namsai, Arunachal Pradesh (Rajavel *et al.* 2011).

**Genus Ayurakitia:** This genus is represented by only two species detected from mountainous areas of western Thailand (Reinert 1972). Recently, occurrence of this genus was reported for the first time from India as larvae collected from *Pandanus* axils in Meghalaya, north-east India and reared to adults, were found to be *Ayurakitia peytoni* (Rajavel and Natarajan 2011).

**Genus Bruceharrisonius:** An earlier subgenus of genus *Ochlerotatus* (Reinert 2003) *Bruceharrisonius* was later elevated to generic rank (Reinert *et al.* 2006). This genus includes four species in India. Three species earlier in Aureostriatus Subgroup (Aureostriatus Group) and one species of Auronitens Subgroup (Alboannulatus Group) of subgenus *Finlaya* genus *Aedes* (Knight and Marks 1952) are now placed in genus *Bruceharrisonius*.

**Genus Christophersiomyia:** A former subgenus of genus *Aedes*, now elevated to generic rank (Reinert *et al.* 2004), *Christophersiomyia* includes five species, of which, four occur in India. *Cr. gombakensis*, reported for the first time from Western Ghats, is a new country record (Reuben *et al.* 1993).

**Genus Canraedes:** *Canraedes* was elevated to genus from former subgenus of *Aedes* (Reinert *et al.* 2009). Of the total ten species included in this genus from Oriental Region, only two species occur in India.

**Genus Collessius:** A newly recognized genus *Collessius* (Reinert *et al.* 2006) includes species formerly in Pseudotaeniatus Subgroup of Mediovittatus Group, subgenus *Finlaya* of genus *Aedes* (Knight and Marks 1952). In India this genus is represented by four species divided into two subgenera.

**Genus Danielsia:** *Danielsia* was elevated to generic rank from synonymy with *Finlaya* and included three taxa (Reinert 2009b). In India this genus is comprised of two species formerly in Alboteniatus Subgroup of subgenus *Finlaya*, genus *Aedes*.

**Genus Dendroskusea:** All five species included in genus *Dendroskusea* are found in India. These species were earlier placed in subgenus *Diceromyia* of genus *Aedes*.

**Genus Downsiomyia:** *Downsiomyia* as a genus was resurrected from synonymy with *Finlaya* (Reinert *et al.* 2006; Reinert and Harbach 2006). This genus is represented by six species in India which were earlier placed in the Niveus Subgroup of subgenus *Finlaya*. The species *Do. nivea* (*Ae. niveus*) has been incriminated as a vector of diurnally subperiodic *Wuchereria bancrofti* in the Nicobar islands (Tewari *et al.* 1995). Tewari and Hiriyan (1995) redescribed this species from Andaman and Nicobar Islands.

**Genus Edwardsaedes:** Comprising of three species, the genus *Edwardsaedes* is represented by only one species in India.

**Genus Finlaya:** In the earlier classification *Finlaya* was one of the largest subgenus of genus *Aedes* comprising of 42 species in India. However, in the current classification, the genus *Finlaya* includes species only from Kochi Group

of subgenus *Finlaya* (Reinert *et al.* 2004, 2006; Reinert and Harbach 2005). Two species of *Finlaya* are found in India. *Fl. flavipennis* was reported for the first time from mangrove forest ecosystem of Andaman and Nicobar Islands (Rajavel and Natarajan 2006).

Genus *Fredwardsius*: *Fredwardsius vittatus* is the only species included in this genus, which is present in India.

Genus *Gilesius*: Of the two species included in this genus, only one species occurs in India.

Genus *Heizmannia*: This genus is represented by 13 species in India, 11 species under subgenus *Heizmannia* and two species under subgenus *Mattinglyia*. Rajput and Singh (1987b) first reported the occurrence of *H. aureochaeta* in India from Manipur state. Similarly *H. chengi* was reported for the first time from Jeypore hill tracts of Orissa (Rajavel *et al.* 2005b).

Genus *Himalaius*: Two species representing this genus are found in India.

Genus *Hopkinsius*: *Hopkinsius* contains two subgenera, *Hopkinsius* and *Yamada*. Only one species belonging to subgenus *Yamada* occurs in India.

Genus *Hulecoeteomyia*: Reinert *et al.* (2006) resurrected *Hulecoeteomyia* to generic status from synonymy with *Finlaya*. Five species included in this genus from India were earlier placed in Chrysolineatus Subgroup of subgenus *Finlaya*.

Genus *Jihlienius*: Of the three species recognized in this genus, only one is found in India.

Genus *Kenknightia*: *Kenknightia* comprised of 12 species of which only two species represent this genus in India.

Genus *Lorrainea*: Presence of *Lo. amesii* and *Lo. fumidus* belonging to genus *Lorrainea* were first reported from India based on collections made in mangrove forests of Sundarbans, in West Bengal (Rajavel *et al.* 2005a), Bhitarkanika in Orissa (Rajavel *et al.* 2005c) and Andaman Nicobar Islands (Rajavel *et al.* 2004). Previously, species of this genus were reported from Thailand, Philippines and Solomon Islands.

Genus *Mucidus*: Reinert (2000b) earlier treated *Mucidus* as a subgenus of newly elevated genus *Ochlerotatus*. However, in the later classification *Mucidus* was elevated to generic rank (Reinert *et al.* 2004). Three species of *Mucidus* are known to exist in India. Confirmed occurrence of *Mu. laniger* was reported from two forest areas of Assam, north-east India (Bhattacharyya *et al.* 2005).

Genus *Neomelaniconion*: Mainly distributed in Afrotropical region, only one species *Neomelaniconion lineatopenne* represent this genus in the Oriental region including India.

Genus *Ochlerotatus*: This genus includes only one species in India.

Genus *Paraedes*: Subgenus of the genus *Aedes* in the earlier classification, *Paraedes* has been raised to genus level (Reinert *et al.* 2004) and is represented by four species in India. In this genus the confirmed presence of *Pr. ostentatio* was reported from a tropical forest zone of Dibrugarh district, Assam (Bhattacharyya *et al.* 2002). Barraud (1934) earlier reported this species from Malabar Coast of Kerela on the basis of female specimens. Reinert (1981) on resurrection of *Pr. chrysoscuta* from *Pr. ostentatio* considered Barraud's Indian record under

distribution of *Pr. chrysoscuta*, but pointed that Mattingly (1958) questioned the records from India. Until evidence is available otherwise, we retain Barraud's record as *Pr. chrysoscuta* and include it in the present checklist.

Genus *Petermattinglyius*: *Petermattinglyius* comprises of two species in India. These two species were earlier placed in subgenus *Diceromyia*. *Pe. franciscoi* was reported for the first time from mangrove forest ecosystem of Orissa (Rajavel *et al.* 2005c).

Genus *Phagomyia*: *Phagomyia* was elevated to generic status from synonymy with *Finlaya* and includes species formerly in Gubernatoris Subgroup of subgenus *Finlaya* (Reinert *et al.* 2006). *Phagomyia* includes 11 species in

**TABLE 2.** Vectors of mosquito-borne diseases in India.

<b>Malaria</b>
<i>Anopheles (Cellia) culicifacies s.l.</i>
<i>Anopheles (Cellia) baimaii</i>
<i>Anopheles (Cellia) fluviatilis s.l.</i>
<i>Anopheles (Cellia) minimus s.l.</i>
<i>Anopheles (Cellia) stephensi</i>
<i>Anopheles (Cellia) sundaicus s.l.</i>
<i>Anopheles (Cellia) annularis s.l.</i>
<i>Anopheles (Cellia) jeyporiensis</i>
<i>Anopheles (Cellia) philippinensis</i>
<i>Anopheles (Cellia) nivipes</i>
<i>Anopheles (Cellia) varuna</i>
<i>Anopheles (Cellia) maculatus s.l.</i>
(WHO 2007; Bhattacharyya <i>et al.</i> 2010; Rao 1984)
<b>Japanese encephalitis</b>
<i>Culex (Culex) vishnui</i>
<i>Culex (Culex) pseudovishnui</i>
<i>Culex (Culex) tritaeniorhynchus</i>
<i>Culex (Culex) fuscocephala</i>
<i>Culex (Culex) quinquefasciatus</i>
<i>Culex (Culex) gelidus</i>
<i>Culex (Culex) whitmorei</i>
<i>Culex (Oculeomyia) bitaeniorhynchus</i>
<i>Culex (Oculeomyia) infula</i>
<i>Culex (Oculeomyia) epidesmus</i>
<i>Anopheles (Anopheles) barbirostris s.l.</i>
<i>Anopheles (Anopheles) peditaeniatus</i>
<i>Anopheles (Cellia) subpictus s.l.</i>
<i>Mansonia (Mansonioides) annulifera</i>
<i>Mansonia (Mansonioides) indiana</i>
<i>Mansonia (Mansonioides) uniformis</i>
(Kanojia 2007)
<b>Dengue</b>
<i>Stegomyia (Stegomyia) aegypti</i>
<i>Stegomyia albopicta</i>
(Kaul <i>et al.</i> 1998; Das <i>et al.</i> 2004)
<b>Chikungunya</b>
<i>Stegomyia (Stegomyia) aegypti</i>
<i>Stegomyia albopicta</i>
(Mourya <i>et al.</i> 2001)
<b>West Nile</b>
<i>Culex (Culex) vishnui</i>
<i>Culex (Culex) quinquefasciatus</i>
(Paramasivan <i>et al.</i> 2003)
<b>Filariasis</b>
<i>Culex (Culex) quinquefasciatus</i>
<i>Mansonia (Mansonioides) annulifera</i>
<i>Mansonia (Mansonioides) uniformis</i>
<i>Downsiomyia nivea</i>
(Agarwal and Sashindran 2006; Tewari <i>et al.</i> 1995)

India. *Ph. feegradei* was new addition to the mosquito fauna of India from the mangrove forest ecosystem of Orissa (Rajavel et al. 2005c).

Genus *Rhinoskusea*: *Rhinoskusea* comprises of three species in India. In this genus one new species *Rh. portonovoensis* was described from mangrove forest of the east coast (Tewari and Hiriyan 1991). The detection of *Rh. wardi* from mangrove forests of Andaman and Nicobar Islands is a new country record (Rajavel and Natarajan 2006). Reinert (2000b) earlier treated *Rhinoskusea* as a subgenus of genus *Ochlerotatus*. However, in the later classification *Rhinoskusea* was elevated to generic status (Reinert et al. 2004).

Genus *Scutomyia*: *Scutomyia* includes only one species in India.

Genus *Stegomyia*: The medically important genus *Stegomyia* includes 19 species in India. In this genus *St. krombeini*, earlier placed in Scutellaris Group of genus *Aedes* and subgenus *Stegomyia* (Huang 1979), was detected first from south India (Tewari et al. 1987) and later from north-east India (Bhattacharyya et al. 2008). Huang (1975) reported this species to be very common in Sri Lanka, which probably remained undetected as it was mistaken for another common species, *St. albopicta*. Hence careful observations during entomological surveys are necessary to differentiate *St. krombeini* from *St. albopicta*, especially in rural and forest areas of India. The records of *St. flavopicta* in Assam, western Himalayas and Coorg (Barraud 1934) was not considered by Huang (1972) who restricted its distribution to Japan and Korea. Its inclusion by Kaur (2003) in Indian species is based on Barraud (1934). While *St. pseudoalbopictus*, *St. subalbopictus* and *St. novalbopictus* have been recorded in collections made in several parts of the country, *St. flavopicta* has not been obtained and hence it is not included in the present checklist. At present of the eight new subgenera recognized for this genus, only eight Indian species are placed in five subgenera. Remaining 11 species under this genus are still without subgeneric placement.

Genus *Tewari*: Of the four species included in this genus, three are found in India. Reinert (2006) placed *Te. agastyai*, *Te. reubenae* and *Te. nummatus* of subgenus *Diceromyia* to a new genus *Tewari*. *Te. agastyai* and *Te. reubenae* were described from Western Ghats (Tewari and Hiriyan 1992).

Genus *Udaya*: *Udaya* includes three species from the Oriental region, of which two are found in India.

Genus *Verrallina*: Divided into three subgenera this genus contains 23 species in India. Subgenus *Neomacleaya* contains most of the species (16) followed by *Harbachius* (four) and *Verrallina* (three). A new species *Ve. assamensis* was described from Assam (Bhattacharyya et al. 2004a) and *Ve. consonensis* was recorded for the first time from Andaman and Nicobar Islands (Rajavel and Natarajan 2006). *Ve. ceylonica* was recently collected from Kerala state (Vector Control Research Centre, Puducherry, India unpublished report).

*Incertae sedis* species: Nine species belonging to two groups "*Aedes*" *sensu auctorum* and "*Ochlerotatus*" *sensu auctorum*, regarded to be of uncertain taxonomic position, are now under this head. Though, not considered as their formal status, the genus and subgenus of these species

are now kept as provided by the authors. Four new species of undetermined subgenus (*Aedes kolhapuriensis*, *Aedes sangitee*, *Aedes panchgangee* and *Aedes sangiti*) were described from Kolhapur district of Maharashtra (Girhe and Sathe 2001; Sathe and Girhe 2001). However, descriptions of these species were inadequate and mainly based on adult female characters. Comparisons with closely related species were not available in the descriptions. Hence, collections of all life stages may be required for comparing with other related species to establish their validity and placing them in relevant genera and subgenera. Genus *Aedes* is now restricted mainly in the Palaearctic and Nearctic Regions. We retained these species in the checklist as they were already in the list of mosquito catalog and mosquito taxonomic inventory. Similarly, five species retained in genus "*Ochlerotatus*" and subgenus "*Finlaya*" (*Oc. auronitens*, *Oc. oreophilus*, *Oc. sintoni*, *Oc. suffusus* and *Oc. versicolor*) are yet to be placed in relevant genera in the current classification.

Tribe Culicini: Culicini includes 81 species in two genera and seven subgenera in India.

Genus *Culex*: Genus *Culex* is represented by 77 species in six subgenera in India with *Lophoceraomyia* being the predominant subgenus with 28 species, followed by subgenus *Culex* with 23 species. This genus contains many important vectors of Japanese encephalitis (JE) virus and microfilariiae in India. In subgenus *Culex*, Vishnui Subgroup contains three most important vectors (*Cx. tritaeniorhynchus*, *Cx. vishnui* and *Cx. pseudovishnui*) of JE in India. Presence of another three members in this subgroup i.e. *Cx. alienus*, *Cx. perplexus* and *Cx. whitei*, which can only be identified distinctively from other members on the basis of larval, pupal and male phallosome characters (Sirivanakarn 1976), can create diagnostic problem in JE vector surveillance in areas of their occurrence. Barraud (1934) earlier collected larvae of *Cx. whitei* from Haflong area of Assam. Few larvae of *Cx. alienus* which is regarded as one of the uncommon members by Sirivanakarn (1976) within this subgroup were collected from Assam. *Cx. perplexus* earlier known only from Andaman Islands was also reported from mangrove forests of Orissa (Rajavel et al. 2005c).

In subgenus *Culiciomyia*, Rajput and Singh (1987a) first reported *Cx. harrisoni* from Senapati district of Manipur. *Cx. spathifurca* was reported for the first time during a mosquito faunistic study in a mangrove forest ecosystem of Tamil Nadu (Rajavel et al. 1998). *Cx. scanloni* known earlier from Indonesia, Malaysia, Thailand and Vietnam was detected from Nagarhole National Park, Karnataka (Rajavel et al. 2011).

Subgenus *Eumelanomyia* is now represented by 10 species in India. One species *Cx. hinglungensis* was reported first time from Manipur state of India (Rajput and Singh 1989).

In subgenus *Lophoceraomyia* one new species *Cx. singhbhumensis* was described from Orissa (Natarajan and Rajavel 2009). Several other species were also recorded from various parts of India such as *Cx. lasiopalpis* and *Cx. pholeter* from south India (Reuben et al. 1993), *Cx. quadripalpis* from the northeast India (Bhattacharyya et al. 2003) and *Cx. pilifemoralis* and *Cx. wilfredi* from Jeypore hills, Orissa (Rajavel et al. 2005b), are new addition to

the mosquito fauna of India. Similarly *Cx. aculeatus*, *Cx. paraculeatus* and *Cx. gracicornis* from Assam, *Cx. cubitatus* from Andaman and Nicobar Islands and *Cx. inculus* and *Cx. demissus* from Orissa are new country records (Rajavel et al. 2011). *Cx. wardi* previously known only from Sri Lanka was recently collected from Kerela state (Vector Control Research Centre, Puducherry, unpublished report).

The subgenus *Maillotia* is represented by only one species in India.

Tanaka (2004) resurrected *Oculeomyia* from synonymy as a new subgenus of genus *Culex*, which includes species earlier placed in the *Bitaniorhynchus* Subgroup of subgenus *Culex*. This subgenus includes 6 species in India. Under this subgenus two females of *Cx. luzonensis* resting on vegetation were collected from Alwar, Rajasthan (Rajavel et al. 2011).

Genus *Lutzia*: It contains four species all in subgenus *Metalutzia*. A new species *Lt. agranensis* was described by Singh and Prakash (2008).

Tribe Culisetini: This tribe includes a single genus. Three species under two subgenera were recorded from India.

Tribe Ficalbiini: The tribe includes two genera i.e. *Ficalbia* and *Mimomyia*. One species of *Ficalbia* and six species of *Mimomyia* are found in India.

Tribe Hodgesiini: Tribe Hodgesiini includes a single genus and represented by only one species in India.

Tribe Mansoniini: Two genera are included in this tribe. The genus *Coquillettidia* includes three species and genus *Mansonia* is represented by four species in India.

Tribe Orthopodomyiini: Only one genus is included in this tribe which is represented by five species in India.

Tribe Sabethini: Sabethini includes 14 genera of which only three (*Malaya*, *Topomyia* and *Tripteroides*) are found in India. Genus *Malaya* is represented by two species and *Tripteroides* by eight species in India. Our experience indicates that genus *Tripteroides* requires further studies in India, especially in bamboo forested areas of the north-east India. Oriental in distribution (Thurman 1959), the genus *Topomyia* was represented by only one species in India, till Bhattacharyya et al. (2007) added three more species from Arunachal Pradesh. Further collections in high rainfall forested areas may increase the number of species in this genus.

Tribe Toxorhynchitini: Only one genus is included in this tribe. The genus *Toxorhynchites* includes nine species in India. *Tx. tyagii* is a new species described from Nilgiri hills, Western Ghats, southern India (Krishnamoorthy et al. 2013).

Tribe Uranotaeniini: *Uranotaenia* is the only genus in this tribe represented by 28 species (15 in subgenus *Pseudoficalbia* and 13 in subgenus *Uranotaenia*) in India. *Ur. micans* was first reported from India from Manipur state (Rajput and Singh 1990). Subsequently, *Ur. ohamai* was recorded from Western Ghats, South India (Reuben et al. 1993). *Ur. dibrugarhensis* is a new species described from Dibrugarh district of Assam (Bhattacharyya et al. 2004b). *Ur. rutherfordi* reported only from Sri Lanka was recently collected from Kerela state (Vector Control Research Centre, Puducherry, unpublished report).

At least 31 species are recognized in India for transmitting various pathogens to humans (Table 2).

Mosquito taxonomy provides essential inputs for vector control. In Vietnam, non-vector species *An. varuna* was misidentified as *An. minimus* and targeted as vector (Bortel et al 2001). *An. fluviatilis*, a recognized primary vector of malaria in the north-east India (Mohapatra et al. 1998) is now found to be a seasonal variant of *An. minimus* (Singh et al. 2010). In respect of malaria vectors, Manguin et al. (2008) suggested that vector control in a region is dependent on the numbers and complexity of the primary and secondary vectors and hence, requires study of the whole anopheline fauna in order to integrate the diverse information about vector system to formulate appropriate and effective control strategy. Besides, sylvatic cycles of some unknown arboviruses capable of spreading to humans may thrive in our forest ecosystem. The history of Chikungunya virus available in the web resource database of Chikungunya Virus Net suggested this virus to be originally circulated in the sylvatic cycle between forest dwelling mosquitoes and non-human primates, has now spread to different parts of the world including India. These facts clearly signify the importance of mosquito fauna study.

Limited studies carried out in some parts of India since the 1980s detected several new species and many new country records. With the increasing mosquito borne diseases and changing ecology there is a need for the reassessment of Indian Culicidae fauna. Comprehensive surveys with the incorporation of DNA-based methods such as DNA barcoding of mosquitoes (Pradeep Kumer et al. 2007) need to be undertaken on a large scale.

**ACKNOWLEDGEMENTS:** Researchers working in the Malariology department of Regional Medical Research Centre (RMRC), Dibrugarh are gratefully acknowledged for their assistance in the preparation of this checklist.

#### LITERATURE CITED

- Agarwal, V.K. and V.K. Sashindran. 2006. Lymphatic Filariasis in India: problems, challenges and new initiatives. *Medical Journal of Armed Forces India* 62: 359-362 (doi: [http://dx.doi.org/10.1016/S0377-1237\(06\)80109-7](http://dx.doi.org/10.1016/S0377-1237(06)80109-7)).
- Barraud, P.J. 1934. *The Fauna of British India, Including Ceylon and Burma. Diptera. Vol V. Family Culicidae. Tribes Megarhinini and Culicini*. London: Taylor and Francis. 463 pp. (<http://www.mosquitocatalog.org/files/pdfs/007000-28.pdf>).
- Berlin, O.G.W. 1972. A proposal for a systematic study of the culicine mosquitoes of India. *Mosquito Systematics* 4: 41-44 (<http://www.mosquitocatalog.org/files/pdfs/MS04N02P041.pdf>).
- Bhattacharyya, D.R., A. Prakash, S.C. Tewari and J. Mahanta. 2000. *Armigeres joloensis* (Diptera: Culicidae), a rare mosquito in upper Assam: first report from India. *Entomon* 25: 63-65.
- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra and J. Mahanta. 2002. A note on the occurrence of *Aedes (Paraedes) ostentatio* (Leicester) in Dibrugarh district, Assam, India (Diptera: Culicidae). *Oriental Insects* 36: 47-49.
- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra and J. Mahanta. 2003. Occurrence of *Culex (Lophoceraomyia) quadripalpis*, *Culex (Lophoceraomyia) mammilifer* and *Uranotaenia (Pseudoficalbia) novobscura* in Assam. *Journal of the American Mosquito Control Association* 19: 13-14 (<http://www.mosquitocatalog.org/files/pdfs/MQ0018.pdf>).
- Bhattacharyya, D.R., A. Prakash, S.C. Tewari, P.K. Mohapatra and J. Mahanta. 2004a. *Verrallina (Neomacleaya) assamensis* (Diptera: Culicidae), a new species from Assam, India. *Journal of the American Mosquito Control Association* 20: 115-120 ([http://www.biodiversitylibrary.org/content/part/JAMCA/JAMCA\\_V20\\_N2\\_P115-120.pdf](http://www.biodiversitylibrary.org/content/part/JAMCA/JAMCA_V20_N2_P115-120.pdf)).
- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra and J. Mahanta. 2004b. *Uranotaenia dibrugarhensis*, a new species in subgenus *Pseudoficalbia* (Diptera: Culicidae) from Assam, India. *Journal of the American Mosquito Control Association* 20: 1-5 ([http://www.biodiversitylibrary.org/content/part/JAMCA/JAMCA\\_V20\\_N1\\_P001-005.pdf](http://www.biodiversitylibrary.org/content/part/JAMCA/JAMCA_V20_N1_P001-005.pdf)).

- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra and J. Mahanta. 2005. Occurrence of *Ochlerotatus (Mucicus) laniger* (Wiedemann) (Diptera: Culicidae) in Assam. *Entomol* 30: 105–107.
- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra and J. Mahanta. 2007. Addition of three species of *Topomyia* mosquitoes (Diptera: Culicidae) to the mosquito fauna of north-east India. *Journal of the American Mosquito Control Association* 23: 76–77 (doi: 10.2987/8756-971X(2007)23[76:AOTSOT]2.0.CO;2).
- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra, D.K. Sarma and J. Mahanta. 2008. A note on the occurrence of *Aedes (Stegomyia) krombeini* (Diptera: Culicidae) in Assam, India. *Entomol* 33: 275–276.
- Bhattacharyya, D.R., A. Prakash, P.K. Mohapatra and D.K. Sarma. 2009. *Armigeres (Armigeres) mahantai*, a new mosquito species from India. *Journal of the American Mosquito Control Association* 25: 1–5 (doi: 10.2987/08-5785.1).
- Bhattacharyya, D.R., A. Prakash, N.P. Sarma, P.K. Mohapatra, S. Singh, D.K. Sarma, M.C. Kalita and J. Mahanta. 2010. Molecular evidence for the involvement of *Anopheles nivipes* (Diptera: Culicidae) in the transmission of *Plasmodium falciparum* in north-eastern India. *Annals of Tropical Medicine and Parasitology* 104: 331–336 (doi: 10.1179/136485910X12743554759948).
- Bortel, W.V., R.E. Harbach, H.D. Trung, P. Roelants, T. Backeljau and M. Coosemans. 2001. Confirmation of *Anopheles varuna* in Vietnam, previously misidentified and mistargeted as the malaria vector *Anopheles minimus*. *American Journal of Tropical Medicine and Hygiene* 65: 729–732 (<http://www.mosquitocatalog.org/files/pdfs/MQ0260.pdf>).
- Christophers, S.R. 1933. *The fauna of British India, including Ceylon and Burma. Diptera. Vol. V. Family Culicidae. Tribe Anophelini*. London: Taylor and Francis. 371 pp. (<http://www.mosquitocatalog.org/files/pdfs/021600-12.PDF>).
- Das, B.P., L. Kabilan, S.N. Sharma, S. Lal, K. Ragu and V.K. Saxena. 2004. Detection of Dengue virus in wild caught *Aedes albopictus* (Skuse) around Calicut Airport, Malapuram district, Kerala, India. *Dengue Bulletin* 28: 210–212 (<http://www.repository.searo.who.int/bitstream/123456789/15976/2/dbv28p210.pdf>).
- Foley, D.H., L.M. Rueda and R.C. Wilkerson. 2007. Insight into mosquito biogeography from country species records. *Journal of Medical Entomology* 44: 554–567 (<http://www.mosquitocatalog.org/files/pdfs/MQ0287.pdf>).
- Gaffigan, T.V., R.C. Wilkerson, J.E. Pecor, J.A. Stoffer and T. Anderson. 2014. *Systematic Catalog of Culicidae*. Accessible at (<http://www.mosquitocatalog.org>). Captured on 17 October 2014.
- Gaston, K.J. and E. Hudson. 1994. Regional patterns of diversity and estimates of global insect species richness. *Biodiversity and Conservation* 3: 493–500.
- Girhe, B.E. and T. V. Sathé. 2001. On a new species of the genus *Aedes* Meigen (Diptera: Culicidae) from India. *Journal of Advanced Zoology* 22: 46–47 (<http://www.mosquitocatalog.org/files/pdfs/050150-0.pdf>).
- Harbach, R.E. 2007. The Culicidae (Diptera): a review of taxonomy, classification and phylogeny. *Zootaxa* 1668: 591–638.
- Harbach, R.E. 2014. *Mosquito Taxonomic Inventory*. Accessible at (<http://mosquito-taxonomic-inventory.info>). Captured on 17 October 2014.
- Harrison, B.A. and J.E. Scanlon. 1975. Medical Entomology studies-II. The subgenus *Anopheles* in Thailand (Diptera: Culicidae). *Contributions of the American Entomological Institute (Ann Arbor)* 12: 1–307 (<http://www.mosquitocatalog.org/files/pdfs/wr113.pdf>).
- Harrison, B.A., R. Rattananarithikul, E.L. Peyton and K. Mongkolpanya. 1990. Taxonomic changes, revised occurrence records and notes on the Culicidae of Thailand and neighboring countries. *Mosquito Systematics* 22: 196–227 (<http://www.mosquitocatalog.org/files/pdfs/MS22N03P196.pdf>).
- History of Chikungunya Virus. 2014. *Chikungunya Virus Net. com*. Accessible at (<http://www.chikungunyavirusnet.com/history-of-chikungunya.html>). Captured on 17 October 2014.
- Huang, Y.M. 1972. Contributions to the mosquito fauna of southeast Asia. XIV. The subgenus *Stegomyia* of *Aedes* in southeast Asia 1- the scutellaris group of species. *Contributions of the American Entomological Institute (Ann Arbor)* 9: 1–109.
- Huang, Y.M. 1975. A new species of *Aedes (Stegomyia)* from Sri Lanka (Ceylon) (Diptera: Culicidae). *Mosquito Systematics* 7: 345–356 (<http://www.mosquitocatalog.org/files/pdfs/MS07N04P345.pdf>).
- Huang, Y.M. 1979. Medical Entomology studies-XI. The subgenus *Stegomyia* of *Aedes* in the Oriental region with keys to the species (Diptera: Culicidae). *Contributions of the American Entomological Institute (Ann Arbor)* 15: 1–79 (<http://www.mosquitocatalog.org/files/pdfs/WR156.pdf>).
- Kanojia, P.C. 2007. Ecological study on mosquito vectors of Japanese encephalitis virus in Bellary district, Karnataka. *Indian Journal of Medical Research* 126: 152–157. (<http://www.medind.nic.in/ibyt/t07/i8/ibyt07i8p152.pdf>).
- Kaul, S.M., R.S. Sharma, S.N. Sharma, M. Panigrahi, P.K. Phukan and S. Lal. 1998. Preventing dengue and DHF, the role of entomological surveillance. *Journal of Communicable Diseases* 38: 187–192.
- Kaur, R. 2003. An update on the distribution of mosquitoes of the tribe Aedini in India (Diptera: Culicidae). *Oriental Insects* 37: 439–455.
- Khan, S.A., R. Handique, S.C. Tewari, P. Dutta, K. Narain and J. Mahanta. 1998. Larval ecology and mosquito fauna of upper Brahmaputra valley, northeast India. *Indian Journal of Malariology* 35: 131–145.
- Knight, K.L. and E.N. Marks. 1952. An annotated checklist of the mosquitoes of subgenus *Finlaya* genus *Aedes*. *Proceedings of the United States National Museum* 101: 513–574.
- Knight, K. and A. Stone. 1977. *A Catalog of the Mosquitoes of the World (Diptera: Culicidae)*. Maryland: Thomas Say Foundation. 611 pp.
- Krishnamoorthy, R., A. Munirathinam, K. Dhananjeyan, J. Hiriyan, T. Mariappan, P. Samuel and A. Venkatesh. 2013. Description of a new species *Toxorhynchites (Toxorhynchites) tyagii* (Diptera: Culicidae), from Nilgiri hills, Western Ghats, southern India. *Zootaxa* 3701: 437–446 (doi: <http://dx.doi.org/10.11646/zootaxa.3701.4.4>).
- Ma, Y.J., F.Y. Qu and J.J. Xu. 1998. Sequence differences of rDNA-ITS2 and species-diagnostic PCR assay of *Anopheles sinensis* and *Anopheles anthropophagus* from China. *Journal of Medical Colleges of PLA* 13: 123–128.
- Ma, Y.J., F.Y. Qu., J.J. Xu. and G.H. Song. 2000a. Differences in sequences of ribosomal DNA second internal transcribed spacer among three members of the *Anopheles hyrcanus* complex from the republic of Korea. *Entomology Sinica* 7: 36–40.
- Ma, Y.J., Qu F.Y., Cao Y.C., and B.J. Yang. 2000b. On molecular identification and taxonomic status of *Anopheles lesteri* and *Anopheles anthropophagus* in China (Diptera: Culicidae). *Chinese Journal of Parasitology and Parasitic Diseases* 18: 325–328.
- Ma, Y. and J. Xu. 2005. The Hyrcanus group of *Anopheles (Anopheles)* in China (Diptera: Culicidae): species discrimination and phylogenetic relationship inferred by ribosomal DNA internal transcribed spacer 2 sequences. *Journal of Medical Entomology* 42: 610–619 (<http://www.mosquitocatalog.org/files/pdfs/MQ0115.pdf>).
- Malhotra, P.R., P.K. Sarkar, N.G. Das, S. Hazarika and V.M. John. 1987. Mosquito survey in Tirap and Subansiri districts of Arunachal Pradesh. *Indian Journal of Malariology* 24: 151–158.
- Manguin, S., C. Garros, I. Dufour, R.E. Harbach and M. Coosemans. 2008. Bionomics, taxonomy and distribution of the major malaria vector taxa of *Anopheles* subgenus *Cellia* in Southeast Asia: an updated view. *Infection, Genetics and Evolution* 8: 489–503 (doi: 10.1016/j.meegid.2007.11.004).
- Mattingly, P.F. 1958. *The Culicine mosquitoes of Indomalayan area. Part III. Genus Aedes Meigen, subgenera Parades Edwards, Rhinoskusea Edwards and Canraedes Edwards*. London: British Museum of Natural History. 61pp.
- Mourya, D.T., J.R. Thakare, M.D. Gokhale, A.M. Powers, S.L. Hundekar, P.C. Jayakumar, V.P. Bondre, Y.S. Shouche and V.S. Paddidri. 2001. Chikungunya virus isolation in mosquito in India. *Acta Virologica* 45: 305–309.
- Mohapatra, P.K., A. Prakash, D.R. Bhattacharyya and J. Mahanta. 1998. Malaria situation in north-eastern region of India. *ICMR Bulletin* 28: 21–30.
- Nagpal, B.N. and V.P. Sharma. 1987. Survey of mosquito fauna of northeastern region of India. *Indian Journal of Malariology* 24: 143–149.
- Natarajan, R. and A.R. Rajavel. 2009. Description of a new species of *Culex (Lophoceraomyia)* from Orissa, India. *Journal of the American Mosquito Control Association* 25: 403–408 (doi: 10.2987/09-5918.1).
- Paramasivan, R., A.C. Mishra and D.T. Mourya. 2003. West Nile Virus: Indian scenario. *Indian Journal of Medical Research* 118: 101–108.
- Pradeep Kumar, N., A.R. Rajavel, R. Natarajan and P. Jambulingam. 2007. DNA barcodes can distinguish species of Indian mosquitoes (Diptera: Culicidae). *Journal of Medical Entomology* 44: 1–7 (doi: 10.1603/0022-2585(2007)44[1:BCDSO]2.0.CO;2).
- Prakash, A., D.K. Sarma, D.R. Bhattacharyya, P.K. Mohapatra, K. Bhattacharjee, K. Das and J. Mahanta. 2010. Spatial distribution and r-DNA second internal transcribed spacer characterization of *Anopheles dirus* (Diptera: Culicidae) complex species in north-east India. *Acta Tropica* 114: 49–54 (doi: 10.1016/j.actatropica.2010.01.003).
- Rajavel, A.R. and R. Natarajan. 2006. Mosquitoes of the mangrove forests of India: Part 3-Andaman and Nicobar Islands, including an update on the mosquito fauna of the islands. *Journal of the American Mosquito Control Association* 22: 366–377 (doi: 10.2987/8756-971X(2006)22[366:MOTMFO]2.0.CO;2).
- Rajavel, A.R. and R. Natarajan. 2011. First record of genus *Ayurakitia* in India: occurrence of *Ayurakitia peytoni* in Meghalaya state. *Journal of the American Mosquito Control Association* 27: 79–80 (doi: 10.2987/10-6062.1).
- Rajavel, A.R., R. Natarajan and A. Munirathinam. 1998. Occurrence of *Culex (Culiciomyia) spathifurca* (Edwards) (Diptera: Culicidae). *Journal of the American Mosquito Control Association* 14: 346–347



- (<http://www.mosquitocatalog.org/files/pdfs/105750-0.pdf>).
- Rajavel, A.R., R. Natarajan and K. Vaidyanathan. 2004. *Aedes fumidus* and *Aedes amesii* in mangrove forests of India – two new country records. *Journal of the American Mosquito Control Association* 20: 315–316 (<http://www.mosquitocatalog.org/files/pdfs/MQ0153.pdf>).
- Rajavel, A.R., R. Natarajan and K. Vaidyanathan. 2005a. Mosquitoes of the mangrove forests of India: Part 2-Sundarbans, West Bengal. *Journal of the American Mosquito Control Association* 21: 136–138 (doi: 10.2987/8756-971X(2005)21[136:MOTMFO]2.0.CO;2).
- Rajavel, A.R., R. Natarajan and K. Vaidyanathan. 2005b. Mosquito collection in the Jeypore hill tracts of Orissa, with notes on three new country records, *Culex (Lophoceraomyia) pilifemoralis*, *Culex (Lophoceraomyia) wilfredi* and *Heizmannia (Heizmannia) chengi*. *Journal of the American Mosquito Control Association* 21: 121–127 (doi: 10.2987/8756-971X(2005)21[121:MCITJH]2.0.CO;2).
- Rajavel, A.R., R. Natarajan and K. Vaidyanathan. 2005c. Mosquitoes of the mangrove forests of India: Part 1-Bhitarkanika, Orissa. *Journal of the American Mosquito Control Association* 21: 131–135 (doi: 10.2987/8756-971X(2005)21[131:MOTMFO]2.0.CO;2).
- Rajavel, A.R., R. Natarajan, K. Vaidyanathan and V.P. Soniya. 2005d. A list of the mosquitoes housed in the mosquito museum at the Vector Control Research Centre, Pondicherry, India. *Journal of the American Mosquito Control Association* 21: 243–251 (doi: 10.2987/8756-971X(2005)21[243:ALOTMH]2.0.CO;2).
- Rajavel, A.R., R. Natarajan, K. Vaidyanathan and P. Jambulingam. 2011. Systematic list of the species added to the mosquito museum at the Vector Control Research Centre, Pondicherry, India. *Journal of the American Mosquito Control Association* 27: 8–14 (doi: 10.2987/8756-971X-27.1.8).
- Rajput, K.B. and T.K. Singh. 1987a. A report on the occurrence of *Culex (Culiciomyia) harrisoni* Sirivanakarn, 1977, in Manipur, India. *Entomon* 12: 373–374.
- Rajput, K.B. and T.K. Singh. 1987b. Occurrence of *Heizmannia aureochaeta* (Leicester) in Manipur, India (Diptera). *Akitu* (N.S.) 92: 10.
- Rajput, K.B. and T.K. Singh. 1987c. A report on the occurrence of *Armigeres (Leicesteria) dolichocephalus* (Leicester), 1908, in Manipur, India. *Mosquito Systematics* 19: 243 (<http://www.mosquitocatalog.org/files/pdfs/MS19N03P243.pdf>).
- Rajput, K.B. and T.K. Singh. 1989. A report on the occurrence of *Culex (Eumelanomyia) hinglungensis* Chu, 1957, in Manipur, India. *Mosquito Systematics* 21: 133–134 (<http://www.mosquitocatalog.org/files/pdfs/MS21N02P133.pdf>).
- Rajput, K.B. and T.K. Singh. 1990. Report on the occurrence of *Uranotaenia (Uranotaenia) micans* Leicester, 1908 in Manipur, India. *Mosquito Systematics* 22: 184 (<http://www.mosquitocatalog.org/files/pdfs/MS22N03P184.pdf>).
- Rao, T.R. 1984. *The Anophelines of India*. Second edition. Malaria Research Centre. New Delhi: Indian Council of Medical Research. 518 pp.
- Rattananarithkul, R., R.E. Harbach, B.A. Harrison, P. Panthusiri and R.E. Coleman. 2006. Illustrated keys to the mosquitoes of Thailand IV. *Anopheles*. *Southeast Asian Journal of Tropical Medicine and Public Health* 41 Suppl 2: 1–87 ([http://www.tm.mahidol.ac.th/seameo/journal\\_37\\_2\\_2006\\_spp.html](http://www.tm.mahidol.ac.th/seameo/journal_37_2_2006_spp.html)).
- Rattananarithkul, R., R.E. Harbach, B.A. Harrison, P. Panthusiri, R.E. Coleman and J.H. Richardson. 2010. Illustrated keys to the mosquitoes of Thailand VI Tribe Aedini. *Southeast Asian Journal of Tropical Medicine and Public Health* 41 Suppl 1: 1–38 (<http://www.tm.mahidol.ac.th/seameo/2010-41-1-suppl/page1-38.pdf>).
- Reinert, J.F. 1972. Contributions to the mosquito fauna of Southeast Asia. Genus *Aedes* Meigen, subgenus *Ayurakitia* Thurman. *Contributions of the American Entomological Institute (Ann Arbor.)* 9: 1–42 (<http://www.mosquitocatalog.org/files/pdfs/wr71.pdf>).
- Reinert, J.F. 1981. Medical Entomology Studies-XV. A revision of the subgenus *Paraedes* of the genus *Aedes* (Diptera: Culicidae). *Contributions of the American Entomological Institute (Ann Arbor)* 18: 1–91 (<http://www.mosquitocatalog.org/files/pdfs/wr173.pdf>).
- Reinert, J.F. 1999. Restoration of *Verrallina* to generic rank in tribe Aedini (Diptera: Culicidae) and descriptions of the genus and three included subgenera. *Contributions of the American Entomological Institute (Ann Arbor)* 31: 1–83 (<http://www.mosquitocatalog.org/files/pdfs/wr351.pdf>).
- Reinert, J.F. 2000a. Restoration of *Ayurakitia* to generic rank in tribe Aedini and a revised definition of the genus. *Journal of the American Mosquito Control Association* 16: 57–65 (<http://www.mosquitocatalog.org/files/pdfs/108000-43-1.pdf>).
- Reinert, J.F. 2000b. New classification of composite genus *Aedes* (Diptera: Culicidae: Aedini) elevation of subgenus *Ochlerotatus* to generic rank, reclassification of the other subgenera and notes on certain subgenera and species. *Journal of the American Mosquito Control Association* 16: 175–188 (<http://www.mosquitocatalog.org/files/pdfs/108000-45.pdf>).
- Reinert, J.F. 2003. Description of *Bruceharrisonius*, a new subgenus of *Ochlerotatus*, and a redescription of its type species *Oc. (Brh.) greenii*. *Journal of the American Mosquito Control Association* 19: 309–322.
- Reinert, J.F. 2006. *Tewarius* Reinert, a new genus of Aedini (Diptera: Culicidae). *Proceedings of the Entomological Society of Washington* 108: 639–645 (<http://biodiversitylibrary.org/page/30254064>).
- Reinert, J.F. 2009a. List of abbreviations for currently valid generic-level taxa in family Culicidae (Diptera). *European Mosquito Bulletin* 27: 68–76 ([http://e-m-b.org/sites/e-m-b.org/files/EMB\(27\)68-76.pdf](http://e-m-b.org/sites/e-m-b.org/files/EMB(27)68-76.pdf)).
- Reinert, J.F. 2009b. Descriptions of *Danielsia* Theobald, a genus recently resurrected from synonymy (Diptera: Culicidae: Aedini) and its type species *Dn. albotaeniata* Leicester. *Zootaxa* 1976: 25–38 (<http://www.mapress.com/zootaxa/2009/f/z01976p038.pdf>).
- Reinert, J.F. and R.E. Harbach. 2005. Descriptions of *Finlaya* Theobald a genus in tribe Aedini (Diptera: Culicidae) and its type species *Fl. kochi* Donitz. *Journal of the American Mosquito Control Association* 21: 154–170 (doi: 10.2987/8756-971X(2005)21[154:DOFTAG]2.0.CO;2).
- Reinert, J.F. and R.E. Harbach. 2006. Descriptions of *Downsiomyia* Vargas (Diptera: Culicidae: Aedini) and its type species *Do. nivea* (Ludlow). *Zootaxa* 1196: 33–61 (<http://www.mapress.com/zootaxa/2006f/z01196p061f.pdf>).
- Reinert, J.F., R.E. Harbach and I.J. Kitching. 2004. Phylogeny and classification of Aedini (Diptera: Culicidae) based on morphological characters of all life stages. *Zoological Journal of the Linnean Society* 142: 289–368 (doi: 10.1111/j.1096-3642.2004.00144.x).
- Reinert, J.F., R.E. Harbach and I.J. Kitching. 2006. Phylogeny and classification of *Finlaya* and allied taxa (Diptera: Culicidae) based on morphological characters of all life stages. *Zoological Journal of the Linnean Society* 148: 1–101 (doi: 10.1111/j.1096-3642.2006.00254.x).
- Reinert, J.F., R.E. Harbach and I.J. Kitching. 2008. Phylogeny and classification of *Ochlerotatus* and allied taxa (Diptera: Culicidae: Aedini) based on morphological data from all life stages. *Zoological Journal of the Linnean Society* 153: 29–114 (doi: 10.1111/j.1096-3642.2008.00382.x).
- Reinert, J.F., R.E. Harbach and I.J. Kitching. 2009. Phylogeny and classification of tribe Aedini (Diptera: Culicidae). *Zoological Journal of the Linnean Society* 157: 700–794 (doi: 10.1111/j.1096-3642.2009.00570.x).
- Reuben, R., S.C. Tewari and J. Hiriyani. 1993. Studies on the mosquito fauna of South India; pp. 47–50, in: M. Coetzee (ed.). *A Festschrift in honour of Botha de Meillon*. Johannesburg: South African Institute for Medical Research.
- Sathe, T.V. and B.E. Girhe. 2001. Biodiversity of mosquitoes in Kolhapur District, Maharashtra. *Rivista di Parassitologia* 18(3): 193–198 (<http://www.mosquitocatalog.org/files/pdfs/MQ0342.pdf>).
- Sathe, T.V. and M. Jagtap. 2012. Taxonomy and molecular phylogeny of new species of the genus *Anopheles (Cellia)* Theobald (Culicidae: Diptera) from India. *Journal of Entomological Research* 36(1): 71–75 (<http://www.indianjournals.com/ijor.aspx?target=ijor:jer&volume=36&issue=1&article=014>).
- Singh, S., A. Prakash, R.N.S. Yadav, P.K. Mohapatra, D.K. Sarma, J. Mahanta and D.R. Bhattacharyya. 2012. *Anopheles (Cellia) maculatus* group: its spatial distribution and molecular characterization of member species in north-east India. *Acta Tropica* 124(1): 62–70 (doi: 10.1016/j.actatropica.2012.06.011).
- Singh, G. and S. Prakash. 2008. A new species of *Lutzia* (Diptera: Culicidae) comparable with *halifaxii* Theobald, 1903 in a semi-arid zone of India; pp. 195–204, in: B.K. Tyagi (ed.). *Vector Borne Diseases: Epidemiology and Control*. Tamil Nadu: Centre for Research in Medical Entomology.
- Singh, O.P., N. Nanda, V. Dev, P. Bali, M. Sohail, A. Mehruunnisa, T. Adak and A.P. Dash. 2010. Molecular evidence of misidentification of *Anopheles minimus* as *Anopheles fluviatilis* in Assam (India). *Acta Tropica* 113(3): 241–244 (doi: 10.1016/j.actatropica.2009.11.002).
- Sirivanakarn, S. 1976. Medical entomology studies-III. A revision of the subgenus *Culex* in the Oriental Region (Diptera: Culicidae). *Contributions of the American Entomological Institute (Ann Arbor)* 13: 1–272 (<http://www.mosquitocatalog.org/files/pdfs/wr121.pdf>).
- Somboon, P., D. Thongwat and R.E. Harbach. 2011. *Anopheles (Cellia) rampae* n. sp., alias chromosomal form K of the Oriental Maculatus Group (Diptera: Culicidae) in Southeast Asia. *Zootaxa* 2810: 47–55 (<http://www.mapress.com/zootaxa/2011/f/z02810p055f.pdf>).
- Taai, K., V. Baimai, A. Saeung, S. Thongsahuan, G. Min, Y. Otsuka, M. Park, M. Fukuda, P. Somboon and W. Choochote. 2013. Genetic compatibility between *Anopheles lesteri* from Korea and *Anopheles paraliae* from Thailand. *Memorias do Instituto Oswaldo Cruz* 108: 312–320.
- Tanaka, K. 2004. Studies on the pupal mosquitoes of Japan (11) subgenera *Oculeomyia* (stat. nov.) and *Sirivanakarnius* (nov.) of the genus *Culex*, with a key of pupal mosquitoes from Ogasawara-gunto (Diptera: Culicidae). *Medical Entomology and Zoology* 55: 217–231.
- Tewari, S.C. and J. Hiriyani. 1991. Description of a new species of *Aedes (Rhinoskusea)* from South India. *Mosquito Systematics* 23: 123–131 (<http://www.mosquitocatalog.org/files/pdfs/MS23N02P123.pdf>).

- Tewari, S.C. and J. Hiriyan. 1992. Description of two new species of *Aedes* (*Diceromyia*) from south India. *Mosquito Systematics* 24: 154–174 (<http://www.mosquitocatalog.org/files/pdfs/MS24N02P154.pdf>).
- Tewari, S.C. and J. Hiriyan. 1995. Description of *Aedes* (*Finlaya*) *niveus* (Diptera: Culicidae) from Andaman and Nicobar, India. *Mosquito Systematics* 27: 167–176 (<http://www.mosquitocatalog.org/files/pdfs/MS27N03P167.pdf>).
- Tewari, S.C., J. Hiriyan and R. Reuben. 1987. Occurrence of *Aedes* (*Stegomyia*) *krombeini* Huang (Diptera: Culicidae) in India. *Current Science* 56: 736–737.
- Tewari, S.C., J. Hiriyan and R. Reuben. 1995. Epidemiology of subperiodic *Wuchereria bancrofti* infection in Nicobar Islands, India. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 89: 163–166.
- Thurman, E.B. 1959. A contribution to a revision of the Culicidae of northern Thailand. *University of Maryland Agricultural Experimental Station Bulletin A-100*: 182 pp. (<http://www.mosquitocatalog.org/files/pdfs/132400-2.pdf>)
- Tyagi, B.K., J. Hiriyan, S.C. Tewari, K. Ayanar, P. Samuel, N. Arunachalam, R. Paramasivan, K.J. Krishnamoorthy, S. Dhananjeyan, L. Victorjerald and R. Rajendran. 2009. Description of a new species *Anopheles pseudosundaicus* (Diptera: Culicidae) from Kerala, India. *Zootaxa* 2219: 49–60 (<http://www.mapress.com/zootaxa/2009/f/z02219p060f.pdf>).
- Tyson, W.H. 1970. Contributions to the mosquito fauna of Southeast Asia. VII. Genus *Aedomyia* Theobald in Southeast Asia. *Contributions of the American Entomological Institute (Ann Arbor)* 6: 1–27 (<http://www.mosquitocatalog.org/files/pdfs/WR44.pdf>).
- Walton, C., J.M. Handley, C. Kuvangkadilok, F.H. Collins, R.E. Harbach, V. Baimai and R.K. Butlin. 1999. Identification of five species of *Anopheles dirus* complex from Thailand, using allele specific polymerase chain reaction. *Medical and Veterinary Entomology* 13: 24–32.
- World Health Organization, 2007. *Anopheles species complexes in South and South East Asia*. SEARO Technical Publication No. 57. 102 pp ([http://apps.searo.who.int/pds\\_docs/B2406.pdf](http://apps.searo.who.int/pds_docs/B2406.pdf)).

RECEIVED: January 2014  
 ACCEPTED: October 2014  
 PUBLISHED ONLINE: December 2014  
 EDITORIAL RESPONSIBILITY: Marco Gottschalk

## APPENDIX 1. Systematic checklist of mosquitoes of India.

### Subfamily Anophelinae

Genus *Anopheles* Meigen, 1818

Subgenus *Anopheles* Meigen, 1818

ANGUSTICORN SECTION

Anopheles Series

Aitkenii Group

*Anopheles* (*Anopheles*) *aitkenii* James, 1903

*Anopheles* (*Anopheles*) *bengalensis* Puri, 1930

*Anopheles* (*Anopheles*) *insulaeflorum* (Swellengrebel & Swellengrebel de Graaf, 1920)

*Anopheles* (*Anopheles*) *pinjaurensis* Barraud, 1932

Culiciformis Group

*Anopheles* (*Anopheles*) *culiciformis* Cogill, 1903

*Anopheles* (*Anopheles*) *sintoni* Puri, 1929

Lindesayi Group

Gigas Complex

*Anopheles* (*Anopheles*) *baileyi* Edwards, 1929

*Anopheles* (*Anopheles*) *gigas* Giles, 1901

Lindesayi Complex

*Anopheles* (*Anopheles*) *lindesayi* Giles, 1900

*Anopheles* (*Anopheles*) *nilgiricus* Christophers, 1924

Plumbeus Group

*Anopheles* (*Anopheles*) *barianensis* James, 1911

Lophoscelomyia Series

Asiaticus Group

*Anopheles* (*Anopheles*) *annandalei* Prashad, 1918

Interruptus Subgroup

*Anopheles* (*Anopheles*) *interruptus* Puri, 1929

LATICORN SECTION

Myzorhynchus Series

Barbirostris Group

Barbirostris Subgroup

*Anopheles* (*Anopheles*) *barbirostris* van der Wulp, 1884

*Anopheles* (*Anopheles*) *hodgkini* Reid, 1962

Vanus Subgroup

*Anopheles* (*Anopheles*) *ahomi* Chowdhury, 1929

*Anopheles* (*Anopheles*) *barbumbrosus* Strickland & Chowdhury, 1927

Hyrcanus Group

*Anopheles* (*Anopheles*) *argyropus* (Swellengrebel, 1914)

*Anopheles* (*Anopheles*) *sinensis* Wiedemann, 1828

Lesteri Subgroup

*Anopheles* (*Anopheles*) *crawfordi* Reid, 1953

*Anopheles* (*Anopheles*) *lesteri* Baisas & Hu, 1936

*Anopheles* (*Anopheles*) *peditaeniatus* (Leicester, 1908)

Nigerrimus Subgroup

*Anopheles* (*Anopheles*) *nigerrimus* Giles, 1900

*Anopheles* (*Anopheles*) *nitidus* Harrison, Scanlon & Reid, 1973

Umbrosus Group

Letifer Subgroup

*Anopheles* (*Anopheles*) *roperi* Reid, 1950

Umbrosus Subgroup

*Anopheles* (*Anopheles*) *umbrosus* (Theobald, 1903)

Subgenus *Cellia* Theobald, 1902

Myzomyia Series

*Anopheles* (*Cellia*) *majidi* Young & Majid, 1928

Funestus Group

*Anopheles* (*Cellia*) *jeyporiensis* James, 1902

Aconitus Subgroup

## APPENDIX 1. Continued.

- Anopheles (Cellia) aconitus* Donitz, 1902  
*Anopheles (Cellia) varuna* Iyengar, 1924  
 Culicifacies Subgroup  
*Anopheles (Cellia) culicifacies* Giles, 1901 (species A, B, C, D and E)  
 Minimus Subgroup  
 Fluviatilis Complex  
*Anopheles (Cellia) fluviatilis* James, 1902 (species S, T, U and V)  
 Minimus Complex  
*Anopheles (Cellia) minimus* Theobald, 1901  
 Neocellia Series  
*Anopheles (Cellia) karwari* (James, 1903)  
*Anopheles (Cellia) moghulensis* Christophers, 1924  
*Anopheles (Cellia) pulcherrimus* Theobald, 1902  
*Anopheles (Cellia) stephensi* Liston, 1901  
*Anopheles (Cellia) theobaldi* Giles, 1901  
 Annularis Group  
*Anopheles (Cellia) annularis* van der Wulp, 1884 (species A and B)  
*Anopheles (Cellia) pallidus* Theobald, 1901  
*Anopheles (Cellia) philippinensis* Ludlow, 1902  
 Nivipes Complex  
*Anopheles (Cellia) nivipes* (Theobald, 1903)  
 Jamesii Group  
*Anopheles (Cellia) jamesii* Theobald, 1901  
*Anopheles (Cellia) pseudojamesi* Strickland & Chowdhury, 1927  
*Anopheles (Cellia) splendidus* Koidzumi, 1920  
 Maculatus group  
*Anopheles (Cellia) pseudowillmori* (Theobald, 1910)  
*Anopheles (Cellia) willmori* (James, 1903)  
 Maculatus Subgroup  
*Anopheles (Cellia) dravidicus* Christophers, 1924  
*Anopheles (Cellia) maculatus* Theobald, 1901  
 Sawadwongporni Subgroup  
*Anopheles (Cellia) sawadwongporni* Rattanakrithikul & Green, 1987  
*Anopheles (Cellia) rampae* Harbach & Somboon, 2011  
 Neomyzomyia Series  
 Kochi Group  
*Anopheles (Cellia) kochi* Donitz, 1901  
 Leucosphyrus Group  
 Hackeri Subgroup  
*Anopheles (Cellia) mirans* Sallum & Peyton, 2005  
 Leucosphyrus Subgroup  
 Dirus complex  
*Anopheles (Cellia) baimaii* Sallum & Peyton, 2005  
*Anopheles (Cellia) elegans* (James, 1903)  
 Tessellatus Group  
*Anopheles (Cellia) tessellatus* Theobald, 1901  
 Paramyzomyia Series  
 Cinereus Group  
*Anopheles (Cellia) turkhudi* Liston, 1901  
 Pyretophorus Series  
*Anopheles (Cellia) pseudosundaicus* (Tyagi, Hiriyan, Tewari, Ayanar, Samuel, Arunachalam, Paramsivam, Krishnamoorthy, Dhanajeyan, Leo & Rajendran, 2009)  
*Anopheles (Cellia) vagus* Donitz, 1902  
 Subpictus complex  
*Anopheles (Cellia) subpictus* Grassi, 1899 (species A, B, C and D)  
 Sundaicus Complex  
*Anopheles (Cellia) sundaicus* (Rodenwaldt, 1925) (cytotype D)
- Subfamily Culicinae**  
**Tribe Aedeomyiini**  
**Genus Aedeomyia** Theobald, 1901  
**Subgenus Aedeomyia** Theobald, 1901  
*Aedeomyia (Aedeomyia) catastica* Knab, 1909  
**Tribe Aedini**  
**Genus "Aedes" sensu auctorum**  
*Aedes kolhapuriensis* Sathe & Girhe, 2001  
*Aedes panchgangee* Sathe & Girhe, 2001  
*Aedes sangitee* Sathe & Girhe, 2001  
*Aedes sangiti* Girhe & Sathe, 2001  
**Genus Aedimorphus** Theobald, 1903  
*Aedimorphus alboscuteclatus* (Theobald, 1905)  
*Aedimorphus caecus* (Theobald, 1901)  
*Aedimorphus culicinus* (Edwards, 1922)  
*Aedimorphus jamesi* (Edwards, 1914)  
*Aedimorphus lowisii* (Theobald, 1910)  
*Aedimorphus nigrostriatus* (Barraud, 1927)  
*Aedimorphus pallidostratus* (Theobald, 1907)  
*Aedimorphus pampangensis* (Ludlow, 1905)  
*Aedimorphus pipersalatus* (Giles, 1902)  
*Aedimorphus punctifemoris* (Ludlow, 1921)  
*Aedimorphus stenoetrus* (Theobald, 1907)

## APPENDIX 1. Continued.

- Aedimorphus syntheticus* (Barraud, 1928)  
*Aedimorphus taeniorhynchoides* (Christophers, 1911)  
*Aedimorphus trimaculatus* (Theobald, 1905)  
*Aedimorphus vexans* (Meigen, 1830)
- Genus *Armigeres*** Theobald, 1901
- Subgenus *Armigeres*** Theobald, 1901
- Armigeres (Armigeres) aureolineatus* (Leicester, 1908)  
*Armigeres (Armigeres) durhami* (Edwards, 1917)  
*Armigeres (Armigeres) joloensis* (Ludlow, 1904)  
*Armigeres (Armigeres) kesseli* Ramalingam, 1987  
*Armigeres (Armigeres) kuchingensis* Edwards, 1915  
*Armigeres (Armigeres) mahantai* Bhattacharyya, Prakash, Mohapatra & Sarma, 2009  
*Armigeres (Armigeres) pallithorax* Dong, Zhou & Dong, 2004  
*Armigeres (Armigeres) subalbatus* (Coquillett, 1898)  
*Armigeres (Armigeres) theobaldi* Barraud, 1934
- Subgenus *Leicesteria*** Theobald, 1904
- Armigeres (Leicesteria) annulipalpis* (Theobald, 1910)  
*Armigeres (Leicesteria) annulitarsis* (Leicester, 1908)  
*Armigeres (Leicesteria) cingulatus* (Leicester, 1908)  
*Armigeres (Leicesteria) dentatus* Barraud, 1927  
*Armigeres (Leicesteria) digitatus* (Edwards, 1914)  
*Armigeres (Leicesteria) dolicocephalus* (Leicester, 1908)  
*Armigeres (Leicesteria) flavus* (Leicester, 1908)  
*Armigeres (Leicesteria) inchoatus* Barraud, 1927  
*Armigeres (Leicesteria) longipalpis* (Leicester, 1904)  
*Armigeres (Leicesteria) magnus* (Theobald, 1908)  
*Armigeres (Leicesteria) omissus* (Edwards, 1914)
- Genus *Ayurakitia*** Thurman, 1954
- Ayurakitia peytoni* (Reinert, 1972)
- Genus *Bruceharrisonius*** Reinert, 2003
- Bruceharrisonius aureostriatus* (Doleschall, 1857)  
*Bruceharrisonius christophersi* (Edwards, 1922)  
*Bruceharrisonius doonii* (Wattal, Bhatia & Kalra, 1958)  
*Bruceharrisonius greenii* (Theobald, 1903)
- Genus *Christophersiomyia*** Barraud, 1923
- Christophersiomyia annulirostris* (Theobald, 1905)  
*Christophersiomyia gombakensis* (Mattingly, 1959)  
*Christophersiomyia ibis* (Barraud, 1931)  
*Christophersiomyia thomsoni* (Theobald, 1905)
- Genus *Cancraedes*** Edwards, 1929
- Cancraedes cancricomus* (Edwards, 1922)  
*Cancraedes simplex* (Theobald, 1903)
- Genus *Collessius*** Reinert, Harbach & Kitching, 2006
- Subgenus *Collessius*** Reinert, Harbach & Kitching, 2006
- Collessius (Collessius) elsiae* (Barraud, 1923)  
*Collessius (Collessius) macdougalli* (Edwards, 1922)  
*Collessius (Collessius) shortti* (Barraud, 1923)
- Subgenus *Alloeomyia*** Reinert, Harbach & Kitching, 2008
- Collessius (Alloeomyia) pseudotaeniatus* (Giles, 1901)
- Genus *Danielsia*** Theobald, 1904
- Danielsia albotaeniata* (Leicester, 1904)  
*Danielsia lepchana* (Barraud, 1923)
- Genus *Dendroskusea*** Edwards, 1929
- Dendroskusea kanarensis* (Edwards, 1934)  
*Dendroskusea micropterus* (Giles, 1901)  
*Dendroskusea periskelata* (Giles, 1902)  
*Dendroskusea ramachandrai* (Reuben, 1967)  
*Dendroskusea reginae* (Edwards, 1922)
- Genus *Downsiomyia*** Vargas, 1950
- Downsiomyia albolateralis* (Theobald, 1908)  
*Downsiomyia albonivea* (Barraud, 1934)  
*Downsiomyia mohani* (Knight, 1969)  
*Downsiomyia nivea* (Ludlow, 1903)  
*Downsiomyia niveoides* (Barraud, 1934)  
*Downsiomyia novonivea* (Barraud, 1934)
- Genus *Edwardsaedes*** Belkin, 1962
- Edwardsaedes imprimens* (Walker, 1861)
- Genus *Finlaya*** Theobald, 1903
- Finlaya flavipennis* Giles, 1904  
*Finlaya poicilia* Theobald, 1903
- Genus *Fredwardsius*** Reinert, 2000
- Fredwardsius vittatus* (Bigot, 1861)
- Genus *Gilesius*** Reinert, Harbach & Kitching, 2006
- Gilesius pulchriventer* (Giles, 1901)
- Genus *Himalaius*** Reinert, Harbach & Kitching, 2006
- Himalaius gilli* (Barraud, 1924)  
*Himalaius simlensis* (Edwards, 1922)
- Genus *Heizmannia*** Ludlow, 1905
- Subgenus *Heizmannia*** Ludlow, 1905
- Heizmannia (Heizmannia) aureochaeta* (Leicester, 1908)

## APPENDIX 1. Continued.

- Heizmannia* (*Heizmannia*) *chandi* Edwards, 1922  
*Heizmannia* (*Heizmannia*) *chengi* Lien, 1968  
*Heizmannia* (*Heizmannia*) *complex* (Theobald, 1910)  
*Heizmannia* (*Heizmannia*) *covelli* Barraud, 1929  
*Heizmannia* (*Heizmannia*) *funerea* (Leicester, 1908)  
*Heizmannia* (*Heizmannia*) *greenii* (Theobald, 1905)  
*Heizmannia* (*Heizmannia*) *himalayensis* Edwards, 1922  
*Heizmannia* (*Heizmannia*) *indica* (Theobald, 1905)  
*Heizmannia* (*Heizmannia*) *reidi* Mattingly, 1957  
*Heizmannia* (*Heizmannia*) *viridis* Barraud, 1929
- Subgenus *Mattinglyia*** Lien, 1968  
*Heizmannia* (*Mattinglyia*) *discrepans* (Edwards, 1922)  
*Heizmannia* (*Mattinglyia*) *tripunctata* (Theobald, 1908)
- Genus *Hopkinsius*** Reinert, Harbach & Kitching, 2008
- Subgenus *Yamada*** Reinert, Harbach & Kitching, 1908  
*Hopkinsius* (*Yamada*) *albocinctus* (Barraud, 1924)
- Genus *Hulecoeteomyia*** Theobald, 1904  
*Hulecoeteomyia* *chrysolineata* (Theobald, 1907)  
*Hulecoeteomyia* *formosensis* (Yamada, 1921)  
*Hulecoeteomyia* *harveyi* (Barraud, 1923)  
*Hulecoeteomyia* *pallirostris* (Edwards, 1922)  
*Hulecoeteomyia* *saxicola* (Edwards, 1922)
- Genus *Jihlienius*** Reinert, Harbach & Kitching, 2006  
*Jihlienius* *unicinctus* Edwards, 1922
- Genus *Kenknightia*** Reinert, 1990  
*Kenknightia* *dissimilis* (Leicester, 1908)  
*Kenknightia* *karwari* (Barraud, 1924)
- Genus *Lorrainea*** Belkin, 1962  
*Lorrainea* *amesii* (Ludlow, 1903)  
*Lorrainea* *fumida* (Edwards, 1928)
- Genus *Mucidus*** Theobald, 1901
- Subgenus *Mucidus*** Theobald, 1901  
*Mucidus* (*Mucidus*) *laniger* (Wiedemann, 1820)  
*Mucidus* (*Mucidus*) *quasiferinus* (Mattingly, 1961)  
*Mucidus* (*Mucidus*) *scatophagoides* Theobald, 1901
- Genus *Neomelaniconion*** Newstead, 1907  
*Neomelaniconion* *lineatopenne* (Ludlow, 1905)
- Genus *Ochlerotatus*** Lynch Arribalzaga, 1891  
 Unassigned to Subgenus  
*Ochlerotatus* *pullatus* (Coquillett, 1904)
- Genus "*Ochlerotatus*" sensu auctorum**
- Subgenus '*Finlaya*' sensu auctorum**  
*Ochlerotatus* (*Finlaya*) *auronitens* Edwards, 1922  
*Ochlerotatus* (*Finlaya*) *oreophilus* Edwards, 1916  
*Ochlerotatus* (*Finlaya*) *sintoni* (Barraud, 1924)  
*Ochlerotatus* (*Finlaya*) *suffusus* (Edwards, 1922)  
*Ochlerotatus* (*Finlaya*) *versicolor* (Barraud, 1924)
- Genus *Paraedes*** Edwards, 1934  
*Paraedes* *barraudi* Edwards, 1934  
*Paraedes* *chrysoseuta* (Theobald, 1910)  
*Paraedes* *menoni* Mattingly, 1958  
*Paraedes* *ostentatio* Leicester, 1908
- Genus *Petermattinglyius*** Reinert, Harbach & Kitching, 2009
- Subgenus *Petermattinglyius*** Reinert, Harbach & Kitching, 2009  
*Petermattinglyius* (*Petermattinglyius*) *franciscoi* (Mattingly, 1959)  
*Petermattinglyius* (*Petermattinglyius*) *iyengari* (Edwards, 1923)
- Genus *Phagomyia*** Theobald, 1905  
*Phagomyia* *assamensis* (Theobald, 1908)  
*Phagomyia* *cacharana* (Barraud, 1923)  
*Phagomyia* *cogilli* (Edwards, 1922)  
*Phagomyia* *deccana* (Barraud, 1923)  
*Phagomyia* *feegradei* (Barraud, 1934)  
*Phagomyia* *gubernatoris* (Giles, 1901)  
*Phagomyia* *inquinata* Edwards, 1922  
*Phagomyia* *khazani* (Edwards, 1922)  
*Phagomyia* *lophoventralis* (Theobald, 1910)  
*Phagomyia* *prominens* (Barraud, 1923)  
*Phagomyia* *stevensoni* (Barraud, 1923)
- Genus *Rhinoskusea*** Edwards, 1929  
*Rhinoskusea* *longirostris* (Leicester, 1908)  
*Rhinoskusea* *portonovoensis* (Tewari & Hiriyan, 1992)  
*Rhinoskusea* *wardi* (Reinert, 1976)
- Genus *Scutomyia*** Theobald, 1904  
*Scutomyia* *albolineata* Theobald, 1904
- Genus *Stegomyia*** Theobald, 1901
- Subgenus *Actinothrix*** Reinert, Harbach & Kitching, 2009  
*Stegomyia* (*Actinothrix*) *edwardsi* Barraud, 1923  
*Stegomyia* (*Actinothrix*) *seampi* (Huang, 1974)
- Subgenus *Heteraspidion*** Reinert, Harbach & Kitching, 2009  
*Stegomyia* (*Heteraspidion*) *annandalei* Theobald, 1910  
*Stegomyia* (*Heteraspidion*) *craggy* Barraud, 1923

## APPENDIX 1. Continued.

**Subgenus *Huangmyia*** Reinert, Harbach & Kitching, 2009  
*Stegomyia (Huangmyia) mediopunctata* Theobald, 1905  
*Stegomyia (Huangmyia) perplexa* Leicester, 1908

**Subgenus *Stegomyia*** Theobald, 1901  
*Stegomyia (Stegomyia) aegypti* (Linnaeus, 1762)

**Subgenus *Xyele*** Reinert, Harbach & Kitching, 2006  
*Stegomyia (Xyele) desmotes* Giles, 1904

Species without subgeneric placement  
*Stegomyia w-alba* Theobald, 1905  
*Stegomyia gardnerii* Ludlow, 1905  
*Stegomyia albopicta* (Skuse, 1895)  
*Stegomyia novalbopicta* (Barraud, 1931)  
*Stegomyia patriciae* (Mattingly, 1954)  
*Stegomyia pseudalbopicta* Borel, 1928  
*Stegomyia subalbopicta* (Barraud, 1931)  
*Stegomyia unilineata* (Theobald, 1906)  
*Stegomyia krombeini* (Huang, 1975)  
*Stegomyia malayensis* (Colless, 1962)  
*Stegomyia scutellaris* (Walker, 1858)

**Genus *Udaya*** Thurman, 1954  
*Udaya argyrurus* (Edwards, 1934)  
*Udaya subsimilis* (Barraud, 1927)

**Genus *Tewarius*** Reinert, 2006  
*Tewarius agastyai* (Tewari & Hiriyan, 1992)  
*Tewarius nummatus* (Edwards, 1923)  
*Tewarius reubena* (Tewari & Hiriyan, 1992)

**Genus *Verrallina*** Theobald, 1903

**Subgenus *Harbachius*** Reinert, 1999  
*Verrallina (Harbachius) abdita* (Barraud, 1931)  
*Verrallina (Harbachius) consonensis* (Reinert, 1973)  
*Verrallina (Harbachius) uniformis* (Theobald, 1910)  
*Verrallina (Harbachius) yusafi* (Barraud, 1931)

**Subgenus *Neomacleaya*** Theobald, 1907  
*Verrallina (Neomacleaya) agrestis* (Barraud, 1931)  
*Verrallina (Neomacleaya) andamanensis* (Edwards, 1922)  
*Verrallina (Neomacleaya) assamensis* Bhattacharyya, Tewari, Prakash, Mohapatra & Mahanta, 2004  
*Verrallina (Neomacleaya) atria* (Barraud, 1928)  
*Verrallina (Neomacleaya) cauta* (Barraud, 1928)  
*Verrallina (Neomacleaya) ceylonica* (Edwards, 1917)  
*Verrallina (Neomacleaya) clavata* (Barraud, 1931)  
*Verrallina (Neomacleaya) comata* (Barraud, 1931)  
*Verrallina (Neomacleaya) indica* (Theobald, 1907)  
*Verrallina (Neomacleaya) pseudodiurna* (Theobald, 1910)  
*Verrallina (Neomacleaya) pseudomediofasciata* (Theobald, 1910)  
*Verrallina (Neomacleaya) rami* (Barraud, 1928)  
*Verrallina (Neomacleaya) seculata* (Menon, 1950)  
*Verrallina (Neomacleaya) unca* (Theobald, 1901)  
*Verrallina (Neomacleaya) vallistris* (Barraud, 1928)  
*Verrallina (Neomacleaya) yerburyi* (Edwards, 1917)

**Subgenus *Verrallina*** Theobald, 1903  
*Verrallina (Verrallina) butleri* (Theobald, 1901)  
*Verrallina (Verrallina) dux* (Dyar & Shannon, 1925)  
*Verrallina (Verrallina) lugubris* (Barraud, 1928)

## Tribe Culicini

**Genus *Culex*** Linnaeus, 1758

**Subgenus *Culex*** Linnaeus, 1758

Piapiens Group

*Culex (Culex) nilgiricus* Edwards, 1916

Piapiens Subgroup

*Culex (Culex) quinquefasciatus* Say, 1823

Theileri Subgroup

*Culex (Culex) theileri* Theobald, 1903

Trifilatus Subgroup

*Culex (Culex) hutchinsoni* Barraud, 1924

*Culex (Culex) vagans* Wiedemann, 1828

Univittatus Subgroup

*Culex (Culex) fuscocephala* Theobald, 1907

*Culex (Culex) perexiguus* Theobald, 1903

Sitiens Group

Barraudi Subgroup

*Culex (Culex) barraudi* Edwards, 1922

*Culex (Culex) edwardsi* Barraud, 1923

Gelidus Subgroup

*Culex (Culex) gelidus* Theobald, 1901

Mimeticus Subgroup

*Culex (Culex) jacksoni* Edwards, 1934

*Culex (Culex) mimeticus* Noe, 1899

*Culex (Culex) mimuloides* Barraud, 1924

*Culex (Culex) mimulus* Edwards, 1915

*Culex (Culex) murrelli* Lien, 1968

## APPENDIX 1. Continued.

- Sitiens Subgroup  
*Culex (Culex) sitiens* Wiedemann, 1828  
*Culex (Culex) whitmorei* (Giles, 1904)
- Vishnui Subgroup  
*Culex (Culex) alienus* Colless, 1957  
*Culex (Culex) pseudovishnui* Colless, 1957  
*Culex (Culex) perplexus* Leicester, 1908  
*Culex (Culex) tritaeniorhynchus* Giles, 1901  
*Culex (Culex) vishnui* Theobald, 1901  
*Culex (Culex) whitei* Barraud, 1923
- Subgenus *Culiciomyia*** Theobald, 1907
- Fragilis Group  
*Culex (Culiciomyia) fragilis* Ludlow, 1903  
*Culex (Culiciomyia) nigropunctatus* Edwards, 1926  
*Culex (Culiciomyia) pallidothorax* Theobald, 1905  
*Culex (Culiciomyia) ramakrishnii* Wattal & Kalra, 1965  
*Culex (Culiciomyia) scanloni* Bram, 1967  
*Culex (Culiciomyia) spathifurca* (Edwards, 1915)  
*Culex (Culiciomyia) viridiventer* Giles, 1901
- Shebbearei Group  
*Culex (Culiciomyia) bailyi* Barraud, 1934  
*Culex (Culiciomyia) shebbearei* Barraud, 1924
- Subgenus *Eumelanomyia*** Theobald, 1909
- Mochthogenes Group
- Foliatus Subgroup  
*Culex (Eumelanomyia) foliatus* Brug, 1932
- Hinglungensis Subgroup  
*Culex (Eumelanomyia) castrensis* Edwards, 1922  
*Culex (Eumelanomyia) hinglungensis* Chu, 1957
- Iphis Subgroup  
*Culex (Eumelanomyia) iphis* Barraud, 1924
- Khazani Subgroup  
*Culex (Eumelanomyia) khazani* Edwards, 1922
- Malayi Subgroup  
*Culex (Eumelanomyia) malayi* (Leicester, 1908)
- Pluvialis Subgroup  
*Culex (Eumelanomyia) pluvialis* Barraud, 1924
- Tenuipalpis Subgroup  
*Culex (Eumelanomyia) tenuipalpis* Barraud, 1924  
*Culex (Eumelanomyia) mohani* Sirivanakarn, 1977
- Protomelanoconion Group  
*Culex (Eumelanomyia) brevipalpis* (Giles, 1902)
- Subgenus *Lophoceraomyia*** Theobald, 1905
- Fraudatrix Group
- Fraudatrix Subgroup
- Cinctellus complex  
*Culex (Lophoceraomyia) cinctellus* Edwards, 1922
- Quadripalpis complex  
*Culex (Lophoceraomyia) aculeatus* Colless, 1965  
*Culex (Lophoceraomyia) paraculeatus* Sirivanakarn, 1977  
*Culex (Lophoceraomyia) quadripalpis* (Edwards, 1914)
- Rubithoracis complex  
*Culex (Lophoceraomyia) rubithoracis* (Leicester, 1908)
- Seniori complex  
*Culex (Lophoceraomyia) seniori* Barraud, 1934
- Variatus complex  
*Culex (Lophoceraomyia) cubitatus* Colless, 1965  
*Culex (Lophoceraomyia) gracicornis* Sirivanakarn, 1977  
*Culex (Lophoceraomyia) variatus* (Leicester, 1908)  
*Culex (Lophoceraomyia) macdonaldi* Colless, 1965
- Inculus complex  
*Culex (Lophoceraomyia) inculus* Colless, 1965
- Minutissimus Subgroup  
*Culex (Lophoceraomyia) minutissimus* (Theobald, 1907)  
*Culex (Lophoceraomyia) infantulus* Edwards, 1922
- Mammilifer Group
- Mammilifer Subgroup
- Flavicornis complex  
*Culex (Lophoceraomyia) flavicornis* Barraud, 1924  
*Culex (Lophoceraomyia) lasiopalpis* Sirivanakarn, 1977  
*Culex (Lophoceraomyia) raghavanii* Rahman, Chowdhury & Kalra, 1968
- Singhbhumensis complex  
*Culex (Lophoceraomyia) singhbhumensis* Natarajan & Rajavel, 2009
- Mammilifer complex  
*Culex (Lophoceraomyia) mammilifer* (Leicester, 1908)  
*Culex (Lophoceraomyia) demissus* Colless, 1965  
*Culex (Lophoceraomyia) wardi* Sirivanakarn, 1977
- Traubi complex  
*Culex (Lophoceraomyia) uniformis* (Theobald, 1905)

## APPENDIX 1. Continued.

- Minor complex  
*Culex (Lophoceraomyia) bengalensis* Barraud, 1934  
*Culex (Lophoceraomyia) bicornutus* (Theobald, 1910)  
*Culex (Lophoceraomyia) minor* (Leicester, 1908)
- Peytoni complex  
*Culex (Lophoceraomyia) peytoni* Bram & Rattanarithikul, 1967
- Pholeter complex  
*Culex (Lophoceraomyia) pholeter* Bram & Rattanarithikul, 1967
- Wilfredi Group  
*Culex (Lophoceraomyia) pilifemoralis* Wang & Feng, 1964  
*Culex (Lophoceraomyia) wilfredi* Colless, 1965
- Subgenus *Maillotia*** Theobald, 1907  
 Hortensis Group  
*Culex (Maillotia) hortensis* Ficalbi, 1889
- Subgenus *Oculeomyia*** Theobald, 1907  
 Bitaeniorhynchus Complex  
*Culex (Oculeomyia) bitaeniorhynchus* Giles, 1901  
*Culex (Oculeomyia) infula* Theobald, 1901  
*Culex (Oculeomyia) luzonensis* Sirivanakarn, 1976
- Sinensis Complex  
*Culex (Oculeomyia) cornutus* Edwards, 1922  
*Culex (Oculeomyia) epidesmus* (Theobald, 1910)  
*Culex (Oculeomyia) sinensis* Theobald, 1903
- Genus *Lutzia*** Theobald, 1903  
**Subgenus *Metalutzia*** Tanaka, 2000  
*Lutzia (Metalutzia) agranensis* Singh & Prakash, 2008  
*Lutzia (Metalutzia) fuscana* Wiedemann, 1820  
*Lutzia (Metalutzia) halifaxii* Theobald, 1903  
*Lutzia (Metalutzia) vorax* Edwards, 1921
- Tribe Culisetini**  
**Genus *Culiseta*** Felt, 1904  
**Subgenus *Allotheobaldia*** Brolemann, 1919  
*Culiseta (Allotheobaldia) longiareolata* (Macquart, 1838)  
**Subgenus *Culiseta*** Felt, 1904  
*Culiseta (Culiseta) alaskaensis* (Ludlow, 1906)  
*Culiseta (Culiseta) niveitaeniata* (Theobald, 1907)
- Tribe Ficalbiini**  
**Genus *Ficalbia*** Theobald, 1903  
*Ficalbia minima* (Theobald, 1901)
- Genus *Mimomyia*** Theobald, 1903  
**Subgenus *Etorleptiomyia*** Theobald, 1904  
*Mimomyia (Etorleptiomyia) luzonensis* (Ludlow, 1905)
- Subgenus *Mimomyia*** Theobald, 1903  
*Mimomyia (Mimomyia) aurea* (Leicester, 1908)  
*Mimomyia (Mimomyia) chamberlaini* Ludlow, 1904  
*Mimomyia (Mimomyia) hybrida* (Leicester, 1908)  
*Mimomyia (Mimomyia) intermedia* (Barraud, 1929)
- Subgenus *Ingramia*** Edwards, 1912  
*Mimomyia (Ingramia) fusca* (Leicester, 1908)
- Tribe Hodgesiini**  
**Genus *Hodgesia*** Belkin, 1962  
*Hodgesia bailyi* Barraud, 1929
- Tribe Mansoniini**  
**Genus *Coquillettidia*** Dyar, 1905  
**Subgenus *Coquillettidia*** Dyar, 1905  
*Coquillettidia (Coquillettidia) crassipes* (van der Wulp, 1881)  
*Coquillettidia (Coquillettidia) novochracea* (Barraud, 1927)  
*Coquillettidia (Coquillettidia) ochracea* (Theobald, 1903)
- Genus *Mansonia*** Blanchard, 1901  
**Subgenus *Mansonioides*** Theobald, 1907  
*Mansonia (Mansonioides) annulifera* (Theobald, 1901)  
*Mansonia (Mansonioides) dives* (Schiner, 1868)  
*Mansonia (Mansonioides) indiana* Edwards, 1930  
*Mansonia (Mansonioides) uniformis* (Theobald, 1901)
- Tribe Orthopodomyiini**  
**Genus *Orthopodomyia*** Belkin, Heinemann & Page, 1970  
*Orthopodomyia albipes* Leicester, 1904  
*Orthopodomyia anopheloides* (Giles, 1903)  
*Orthopodomyia andamanensis* Barraud, 1934  
*Orthopodomyia flavicosta* Barraud, 1927  
*Orthopodomyia flavithorax* Barraud, 1927
- Tribe Sabethini**  
**Genus *Malaya*** Leicester, 1908  
*Malaya genurostris* Leicester, 1908  
*Malaya jacobsoni* (Edwards, 1930)
- Genus *Topomyia*** Leicester, 1908  
**Subgenus *Suaymyia*** Thurman, 1959  
*Topomyia (Suaymyia) cristata* Thurman, 1959
- Subgenus *Topomyia*** Leicester, 1908  
*Topomyia (Topomyia) bifurcata* Dong, Wang & Lu, 1995



## APPENDIX 1. Continued.

*Topomyia (Topomyia) aureoventer* (Theobald, 1910)*Topomyia (Topomyia) hirtusa* Gong, 1989**Genus *Tripteroides*** Giles, 1904**Subgenus *Rachionotomyia*** Theobald, 1905*Tripteroides (Rachionotomyia) affinis* (Edwards, 1913)*Tripteroides (Rachionotomyia) aranoides* (Theobald, 1901)*Tripteroides (Rachionotomyia) coonorensis* Mattingly, 1981*Tripteroides (Rachionotomyia) edwardsi* (Barraud, 1929)*Tripteroides (Rachionotomyia) serratus* (Barraud, 1929)**Subgenus *Tripteroides*** Giles, 1904*Tripteroides (Tripteroides) indicus* (Barraud, 1929)*Tripteroides (Tripteroides) similis* (Leicester, 1908)*Tripteroides (Tripteroides) tarsalis* Delfinado & Hodges, 1968**Tribe Toxorhynchitini****Genus *Toxorhynchites*** Theobald, 1901**Subgenus *Toxorhynchites*** Theobald, 1901*Toxorhynchites (Toxorhynchites) albipes* (Edwards, 1922)*Toxorhynchites (Toxorhynchites) edwardsi* (Barraud, 1924)*Toxorhynchites (Toxorhynchites) gravelyi* (Edwards, 1921)*Toxorhynchites (Toxorhynchites) kempi* (Edwards, 1921)*Toxorhynchites (Toxorhynchites) klossi* (Edwards, 1921)*Toxorhynchites (Toxorhynchites) metallicus* Leicester, 1904*Toxorhynchites (Toxorhynchites) minimus* (Theobald, 1905)*Toxorhynchites (Toxorhynchites) splendens* (Wiedemann, 1819)*Toxorhynchites (Toxorhynchites) tyagii* Krishnamoorthy *et al.*, 2013**Tribe Uranotaeniini****Genus *Uranotaenia*** Lynch Arribalzaga, 1891**Subgenus *Pseudoficalbia*** Theobald, 1912

## Section A

## Bicolor series

*Uranotaenia (Pseudoficalbia) atra* Theobald, 1905*Uranotaenia (Pseudoficalbia) bicolor* Leicester, 1908*Uranotaenia (Pseudoficalbia) lutescens* Leicester, 1908*Uranotaenia (Pseudoficalbia) obscura* Edwards, 1915

## Bimaculata Series

*Uranotaenia (Pseudoficalbia) nivipleura* Leicester, 1908*Uranotaenia (Pseudoficalbia) novobscura* Barraud, 1934

## Recondita Series

*Uranotaenia (Pseudoficalbia) dibrugarhensis* Bhattacharyya, Prakash, Mohapatra & Mahanta, 2004*Uranotaenia (Pseudoficalbia) husaini* Qutubuddin, 1947*Uranotaenia (Pseudoficalbia) luteola* Edwards, 1934*Uranotaenia (Pseudoficalbia) maculipleura* Leicester, 1908*Uranotaenia (Pseudoficalbia) mattinglyi* Qutubuddin, 1951*Uranotaenia (Pseudoficalbia) recondita* Edwards, 1922*Uranotaenia (Pseudoficalbia) ohamai* Tanaka, Mizusawa & Saugstad, 1975*Uranotaenia (Pseudoficalbia) stricklandi* Barraud, 1926

## Section B

## Maxima Series

*Uranotaenia (Pseudoficalbia) maxima* Leicester, 1908**Subgenus *Uranotaenia*** Lynch Arribalzaga*Uranotaenia (Uranotaenia) alboannulata* (Theobald, 1905)*Uranotaenia (Uranotaenia) annandalei* Barraud, 1926*Uranotaenia (Uranotaenia) campestris* Leicester, 1908*Uranotaenia (Uranotaenia) christophersi* Barraud, 1926*Uranotaenia (Uranotaenia) edwardsi* Barraud, 1926*Uranotaenia (Uranotaenia) hebes* Barraud, 1931*Uranotaenia (Uranotaenia) lateralis* Ludlow, 1905*Uranotaenia (Uranotaenia) longirostris* Leicester, 1908*Uranotaenia (Uranotaenia) micans* Leicester, 1908*Uranotaenia (Uranotaenia) macfarlanei* Edwards, 1914*Uranotaenia (Uranotaenia) orientalis* Barraud, 1926*Uranotaenia (Uranotaenia) rutherfordi* Edwards, 1922*Uranotaenia (Uranotaenia) testacea* Theobald, 1905