

# *Diclidurus albus* Wied-Neuwied, 1820 (Chiroptera: Emballonuridae): First record of the species in the state of Paraíba, Brazil

Alvino Pedrosa Ferreira<sup>1\*</sup>, Daniela de Carvalho Melo<sup>1</sup> and Alan Loures-Ribeiro<sup>2</sup>

1 Universidade Federal da Paraíba, Programa de Pós-graduação em Ciências Biológicas. Cidade Universitária - Castelo Branco, 58059-900, João Pessoa, PB, Brazil.

2 Universidade Federal da Paraíba, Departamento de Sistemática e Ecologia. Cidade Universitária - Castelo Branco, 58059-900, João Pessoa, PB, Brazil.

\* Corresponding author. E-mail: [alvinopf@yahoo.com.br](mailto:alvinopf@yahoo.com.br)

**ABSTRACT:** *Diclidurus albus* Wied-Neuwied, 1820 has few known records in Brazil. This study reports the easternmost record of the species and the first for the state of Paraíba, Brazil. One adult female specimen of *D. albus* was captured in the forest canopy of Reserva Biológica Guaribas – SEMA 3. This record extends the distribution of *D. albus* to the north in the Brazilian Atlantic Forest.

The bat family Emballonuridae includes 13 genera, among these is *Diclidurus* Wied-Neuwied, 1820 (Kitchener 1989; Koopman 1993; Simmons 2005; Reis *et al.* 2007). The genus includes four Neotropical species that are popularly known as ghost bats because of their whitish color (Ceballos and Medellín 1988; Jones and Hood 1993): *Diclidurus albus* Wied-Neuwied, 1820; *D. ingens* Hernández-Camacho, 1955; *D. isabella* (Thomas, 1920); and *D. scutatus* Peters, 1869. The species from this genus are considered rare and little is known about their biology, ecology, and distribution (Eisenberg and Redford 1999; Emmons and Feer 1999; Hood and Gardner 2008), mainly because of the low rate of capture, as they fly high in open areas and in the forest canopy (Ceballos and Medellín 1988; Linares 1998; Lim *et al.* 1999; Ochoa G. *et al.* 2008).

*Diclidurus albus* is the only species found west of the Andes, with the widest known distribution of the genus occurring from Mexico to the northwest of Peru, Brazil and the island of Trinidad in an altitudinal range from sea level to 1700 m.a.s.l. (Ceballos and Medellín 1988; Simmons 2005; Hood and Gardner 2008; Moscoso and Tirira 2009). The type locality of *D. albus* is Canavieiras, Pardo River, Bahia, Brazil (Vieira 1942). Two subspecies are currently recognized for this species: *Diclidurus albus albus*, distributed in Guyana, Surinam, Brazil and Peru; and *Diclidurus albus virgo*, occurring from Mexico south to Ecuador, Colombia, Venezuela and Trinidad (Simmons 2005; Hood and Gardner 2008).

*Diclidurus albus* is an aerial insectivorous bat (Kalko 1995). The species feeds above the canopy using low calls with very long pulse intervals during fast flight and starts call alternations when approaching either prey or obstacles (Jung *et al.* 2007). Deciduous and moist tropical forests and vegetation with large leaves, like palms (especially *Cocos nucifera*) or banana, are the preferences of the species (Cuervo *et al.* 1986; Ceballos and Medellín 1988; Muñoz 2001; Sodr e and Uieda 2006; Hood and Gardner 2008). Some individuals have been recorded

in open urban areas, like human constructions and near mangroves, rivers and the sea (Moscoso and Tirira 2009). The reproductive period of *D. albus* in Mexico seems to extend from January to June and the copulation is thought to occur in January or February (Ceballos and Medellín 1988). During reproduction, groups of up to four individuals, generally including one male and three females, may gather in shelters (Ceballos and Medellín 1988).

There are few records of the species in Brazil, which are restricted to the states of Amazonas (Reis and Peracchi 1987, Bernard and Fenton 2002, Lim and Engstrom 2005, Castro-Arellano *et al.* 2007, Bernard *et al.* 2011, Peracchi *et al.* 2011, Martins *et al.* 2012), Par a (Peracchi *et al.* 2011), Amap a (Peracchi *et al.* 1984, Peracchi *et al.* 2011), Espirito Santo (Mendes *et al.* 2010, Peracchi *et al.* 2011), Bahia (Oliveira 2004, Vaz 2005, Faria *et al.* 2006, Peracchi *et al.* 2011), Rond nia (Peracchi *et al.* 2011) and Pernambuco (Feij o and Langguth, 2011) (Figure 1). Despite being reported in a neighboring state, the species has never been recorded in the state of Para ba.


Here, we report the easternmost record of this species and the first for the state of Para ba, Brazil (Figure 1). The registry occurred during a canopy ornithological mist-net study developed under permit number: 32654-1, authentication code: 74714939, from SISBIO system at Instituto Chico Mendes de Conserva o da Biodiversidade (ICMBio). One adult female specimen of *D. albus* was captured, photographed and then released (Figure 2). The capture occurred in the forest of the Reserva Biol gica Guaribas – SEMA 3 remnant (6°48'17.82" S 35°4'59.94" W, Figure 1), in the municipality of Rio Tinto on January 18, 2012 at 14:00 h. It was a cloudy, rainy day with 88 % relative humidity and 26.7 °C. The mist-net was installed at a height of 20 m above the ground on the forest canopy, next to a natural glade surrounded by trees with an average height of 25 m and a dense understory formed by secondary growth vegetation.

The identification of the specimen was possible because of its characteristic coloration and morphology, associated with the distribution of the other congeneric species (Hood and Gardner 2008; Dalponte and Aguiar 2009). The specimen had large black eyes, no wing sacs, two valved-chambered black uropatagial glands, white and rounded ears, as well as long and silky white hair without variation in the color between the base and the edge. Its skin had a whitish coloration, which was semi-transparent at the wings, with a visibly red innervation. Its external measurements (mm) are as follows: forearm length = 58.9; head and body length = 62.3; total length = 74.7; ear length = 10.6; wingspan = 225; weight = 13 g.

The forest remnant has a total area of 338 ha approximately 3 km away from a mangrove area and 13 km from the sea and is characterized by secondary semi-deciduous lowland forest, associated with savannahs and

transitional areas. Its climate is predominantly hot and humid, with average annual temperatures from 24 to 26°C, with the annual maximum of about 36°C (Barbosa et al. 2011). The vegetation is under a natural regeneration process after episodes of logging and other forms of human pressure which occurred decades ago (MMA/IBAMA 2003).

The capture of *D. albus* in a remnant of Atlantic Forest in the state of Paraíba may indicate the possible presence of other important bat species in the state, or at least more species that fly high and are underestimated in understory studies. We report the extension of the distribution of *D. albus* to about 150 km to the north of the biome. Furthermore, we recommend the conservation of the Atlantic Forest fragments of the region and the inclusion of the upper strata of these fragments in bat surveys.



**FIGURE 1.** Distribution of *Diclidurus albus* in Brazil (star = new record; circles = existing records). Codes are as follows: 1: Porto Velho, RO (Peracchi et al. 2011); 2: Manaus, AM (Reis and Peracchi 1987, Bernard and Fenton 2002, Lim and Engstrom 2005, Peracchi et al. 2011, Bernard et al. 2011); 3: Floresta Nacional do Tapajós, PA (Castro-Arellano et al. 2007); 4: Tracajatuba, AP (Peracchi et al. 1984, Peracchi et al. 2011); 5: Floresta Nacional do Carajás, PA (Martins et al. 2012); 6: Belém, PA (Peracchi et al. 2011); 7: Rio Tinto, PB (this study); 8: Pernambuco State (Feijó and Langguth 2011); 9: Itabuna, BA (Faria et al. 2006); 10: Buerarema, BA (Vaz 2005); 11: Canavieiras, BA (Peracchi et al. 2011); 12: Rio Jequitinhonha, BA (Oliveira 2004); 13: Espírito Santo State (Mendes et al. 2010, Peracchi et al. 2011).



FIGURE 2. Live photograph of an adult female of *Diclidurus albus* captured in the Atlantic Forest fragment of Reserva Biológica Guaribas – SEMA 3.

**ACKNOWLEDGMENTS:** The authors would like to thank Programa de Pós-graduação em Ciências Biológicas (PPGCB), Reestruturação e Expansão das Universidades Federais (REUNI) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for financial support; Reserva Biológica Guaribas for logistics support; and José Anderson Feijó for the confirmation of the identification of the species.

#### LITERATURE CITED

- Barbosa, M.R., W.W. Thomas, E.L.P. Zárate, R.B. Lima, M.F. Agra, I.B. Lima, M.C.R. Pessoa, A.R.L. Lourenço, G.C.D. Júnior, R.A.S. Pontes, E.C.O. Chagas, J.L. Viana, P.C.G. Neto, C.M.L.R. Araújo, A.A.M. Araújo, G.B. Freitas, J.R. Lima, F.O. Silva, L.A.F. Vieira, L.A. Pereira, R.M.T. Costa, R.C. Duré and MGV Sá. 2011. Checklist of the Vascular Plants of the Guaribas Biological Reserve, Paraíba, Brazil. *Revista Nordestina de Biologia* 20(2): 79-106.
- Bernard, E. and M.B. Fenton. 2002. Species diversity of bats (Mammalia: Chiroptera) in forest fragments, primary forests, and savannas in central Amazonia, Brazil. *Canadian Journal of Zoology* 80(6): 1124-1140.
- Bernard, E., V.C. Tavares and E. Sampaio. 2011. Compilação atualizada das espécies de morcegos (Chiroptera) para a Amazônia Brasileira. *Biota Neotropica* 11(1): 35-46.
- Castro-Arellano, I., S.J. Presley, L.N. Saldanha, M.R. Willig and J.M. Wunderle. 2007. Effects of reduced impact logging on bat biodiversity in terra firme forest of lowland Amazonia. *Biological Conservation* 138(1-2): 269-285.
- Ceballos, G. and R. Medellín. 1988. *Diclidurus albus*. *Mammalian Species* 316: 1-4.
- Cuervo, A., J. Hernandez and A. Cadena. 1986. Lista actualizada de los mamíferos de Colombia, anotaciones sobre su distribución. *Caldasia* 15(71-75): 471-501.
- Dalponete, J.C. and L.M.S. Aguiar. 2009. The first record of *Diclidurus ingens* Hernandez Camacho, 1955 (Emballonuridae) in Central Brazil. *Biota Neotropica* 9(4): 249-252.
- Eisenberg, J.F. and K.H. Redford. 1999. *Mammals of the Neotropics: The Central Neotropics volume 3: Ecuador, Peru, Bolivia, Brazil*. Chicago: The University of Chicago Press. 609 p.
- Emmons, L.H. and F. Feer. 1999. *Mamíferos de los bosques húmedos de América tropical, una guía de campo*. Santa Cruz de la Sierra: Editorial FAN. 298 p.
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from the Atlantic rainforest of southern Bahia, Brazil. *Biota Neotropica* 6(2): 2-13.
- Feijó, J.A. and A. Langguth. 2011. Lista de Quirópteros da Paraíba, Brasil com 25 novos registros. *Chiroptera Neotropical* 17(2): 1055-1062.
- Hood, C. and A.L. Gardner. 2008 [2007]. Family Emballonuridae Gervais, 1856; p. 188-207. In A.L. Gardner (ed.). *Mammals of South America, volume 1: marsupials, xenarthrans, shrews, and bats*. Chicago: The University of Chicago Press [Available from 31 March, 2008].
- Jones, J. and C. Hood. 1993. Synopsis of South American bats of the family Emballonuridae. *Occasional Papers of the Museum Texas Tech University* 155: 1-32.
- Jung, K., E.K.V. Kalko and V.O. Herversen. 2007. Echolocation calls in Central American emballonurid bats: signal design and call frequency alternation. *Journal of Zoology* 272(2): 125-137.
- Kalko, E.K.V. 1995. Echolocation signal design, foraging habitats and guild structure in six Neotropical sheath-tailed bats (Emballonuridae).

- Symposia of the Zoological Society of London* 67: 259-273.
- Kitchener, D.J. 1989. Emballonuridae; p.1-15 In D.W. Walton and B.J. Richardson (ed.). *Fauna of Australia: Mammalia*. Volume 1b. Canberra: Australian Government Publishing Service.
- Koopman, K.F. 1993. Order Chiroptera; p.137-241 In D.E. Wilson and D.M. Reeder (ed.). *Mammal species of the world: a taxonomic and geographic reference*. Volume II. Washington D.C.: Smithsonian Institution Press.
- Lim, B.K., M.D. Engstrom, R.M. Timm, R.P. Anderson, and L.C. Watson. 1999. First records of 10 bat species in Guyana and comments on diversity of bats in Iwokrama Forest. *Acta Chiropterologica* 1: 179-190.
- Lim, B.K. and M.D. Engstrom. 2005. Mammals of Iwokrama Forest. *Proceedings of the Academy of Natural Sciences of Philadelphia* 154(1): 71-108.
- Linares, J.O. 1998. *Mamíferos de Venezuela*. Caracas: Sociedad Conservacionista Audubon de Venezuela. 691 p.
- Martins, F.D., A. Castilho, J. Campos, F.M. Hatano and S.G. Rolim. 2012. *Fauna da Floresta Nacional de Carajás*. São Paulo: Nitro Imagens. 116 p.
- Mendes, P., T.B.Vieira, M. Oprea, S.R. Lopes, A.D. Ditchfield and M. Zortéa. 2010. O conhecimento sobre morcegos (Chiroptera: Mammalia) do Estado do Espírito Santo, Sudeste do Brasil. *Papéis Avulsos de Zoologia* 50(22): 363-373.
- MMA/IBAMA 2003. *Plano de Manejo Reserva Biológica Guaribas*. Electronic Database accessible at <http://www.icmbio.gov.br/portal/biodiversidade/unidades-de-conservacao/planos-de-manejo.html>. Captured on 12 December 2012.
- Moscoso, P. and D.G. Tirira. 2009. Nuevos registros y comentarios sobre la distribución del murciélago blanco común *Diclidurus albus* (Chiroptera: Emballonuridae) en Ecuador. *Mastozoología Neotropical* 16(1): 233-237.
- Muñoz, A.J. 2001. *Los murciélagos de Colombia: Sistemática, Distribución, Descripción, Historia Natural y Ecología*. Medellín: Editorial Universidad de Antioquia. 391 p.
- Ochoa G., J., J. Sánchez H., F. García, and J. A. Clavijo. 2008. An undescribed behavior of the Isabelle's ghost bat (*Diclidurus isabellus*), with distributional notes in Venezuela. *Memoria de la Fundación La Salle de Ciencias Naturales* 168(70): 141-148.
- Oliveira, J.A. 2004. Diversidade de mamíferos e o estabelecimento de áreas prioritárias para a conservação do bioma Caatinga; p. 263-282 In Silva, J.M.C., M. Tabarelli, M.T. Fonseca and L.V. Lins (ed.). *Biodiversidade da Caatinga: área e ações prioritárias para a conservação*. Brasília: Ministério do Meio Ambiente e Universidade Federal de Pernambuco.
- Peracchi, A.L., L.R. Sansão and A.M. Tanurre. 1984. Quirópteros do Território Federal do Amapá, Brasil (Mammalia: Chiroptera). *Arquivos da Universidade Federal Rural do Rio de Janeiro* 7(2): 89-100.
- Peracchi, A.L., I.P. Lima, N.R. Reis, M.R. Nogueira and H.O. Filho. 2011. Ordem Chiroptera; p. 155-234 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Mamíferos do Brasil*. Volume II. Londrina: Nelio R. dos Reis.
- Reis, N.R. and A.L. Peracchi. 1987. Quirópteros da região de Manaus, Amazonas, Brasil (Mammalia, Chiroptera). *Boletim do Museu Paraense Emílio Goeldi, série Zoológica* 3(2): 161-182.
- Reis, N.R., A.L. Peracchi, W.A. Pedro and I.P. Lima. 2007. *Morcegos do Brasil*. Londrina: Nelio R. dos Reis. 253 p.
- Simmons, N.B. 2005. Order Chiroptera; p. 312-529 In D.E. Wilson and D.M. Reeder (ed.). *Mammal species of the world, a taxonomic and geographic reference*. Volume III. Baltimore: The John Hopkins University Press.
- Sodré, M.M. and W. Uieda. 2006. First record of the Ghost Bat *Diclidurus scutatus* Peters (Mammalia, Chiroptera, Emballonuridae) in São Paulo city, Brazil. *Revista Brasileira de Zoologia* 23(3): 897-898.
- Vaz, S.M. 2005. Mamíferos colecionados pelo serviço de estudos e pesquisas sobre a febre amarela nos municípios de Ilhéus e Buerarema, Estado da Bahia, Brasil. *Arquivos do Museu Nacional* 63(1): 21-28.
- Vieira, C.O.C. 1942. Ensaio monográfico sobre os quirópteros do Brasil. *Arquivos de Zoologia* 3(8): 1-471.

RECEIVED: December 2012

ACCEPTED: July 2013

PUBLISHED ONLINE: August 2013

EDITORIAL RESPONSIBILITY: Paúl M. Velazco