

First record of *Phyllostomus elongatus* (É. Geoffroy, 1810) (Mammalia: Chiroptera) for the state of Espírito Santo, southeastern Brazil

Sergio Barbiero Lage^{1*}, Rodrigo da Silva Cipriano¹, Atilla Colombo Ferreguetti¹ and Rodrigo Lemes Martins²

¹ Faculdades Integradas São Pedro. Rodovia Serafim Derenzi, 3115, São Pedro. CEP 29053-250. Vitória, ES, Brazil.

² Universidade Federal do Rio de Janeiro – Campus Macaé. Av. São José do Barreto, no 764. Bairro São José do Barreto. CEP 27971-550. Caixa Postal 119331. Macaé, RJ, Brazil.

* Corresponding author. E-mail: sergioblage@gmail.com

ABSTRACT: We report the occurrence of *Phyllostomus elongatus* in the municipality of Linhares, state of Espírito Santo. The new record fills a gap in the distribution of the species in southeastern Brazil and increases the state's mammal list.

Phyllostomus Lacépède, 1799 is widely distributed in Central and South America, being represented by four medium to large-sized species (forearms from 58 to 88 mm): *Phyllostomus discolor* (Wagner, 1843), *Phyllostomus elongatus* (É. Geoffroy, 1810), *Phyllostomus hastatus* (Pallas, 1767), and *Phyllostomus latifolius* (Thomas, 1901) (Williams and Genoways 2008; Reis *et al.* 2011).

Phyllostomus elongatus is intermediate, in size, between *P. hastatus* and *P. discolor*. Its measurements include: total length (head, body and tail) 99 to 115 mm; tail 14 to 27 mm; forearm 61 to 71 mm; and weight 30 to 57 g (see Nogueira *et al.* 2007). Mensural data, in addition to wing tip color, fur color, dental morphology, and size of maxillary incisors, can easily differentiate the species (Rodríguez-Posada and Sánchez-Palomino 2009). Other characters may also assist in the identification of *P. elongatus*, including long ears, larger than the head and with rounded edges, well-developed noseleaf with a pointed end, and calcar longer than the foot (Vieira 1942; Husson 1962).

Phyllostomus elongatus is currently known from 11 Brazilian states: Acre, Alagoas, Amazonas, Amapá, Bahia, Mato Grosso, Pará, Pernambuco, Rio de Janeiro, Rondônia, and Roraima (Bernard *et al.* 2011; Tavares *et al.* 2008) (Figure 1). Marinho-Filho (1996) reported its occurrence in the state of Espírito Santo, but no supporting data for this record was provided or is available from more recent compilations (Moreira *et al.* 2008; Mendes *et al.* 2010; Moreira and Mendes 2010).

During a bat inventory in the Povoação Village ($19^{\circ}33'40''$ S and $39^{\circ}48'45''$ W), northern margin of the Doce River, municipality of Linhares (Figure 1), we captured a male *P. elongatus* that represents the first record of this species for the state of Espírito Santo. This specimen was captured in July 2011, in a fragment of "cabruca" located in a tropical lowland rainforest area. The "cabruca" consists of a traditional system where the cacao (*Theobroma cacao* L., Sterculiaceae) is planted

under the native forest canopy. In this system, only 10% of the original forest canopy is maintained (Sambuichi 2002; Rolim and Chiarello 2004). The "cabrucas" keep a stratified structure of forests, ensuring the occurrence of many native animal species (Rolim and Chiarello 2004). For the "cabrucas" of Espírito Santo, we can highlight the maintenance of a non-dense canopy, based on tree species of Moraceae (Rolim *et al.* 2006), a family widely consumed by fruit bats (Pinto and Ortêncio-Filho 2006). The region is classified as Aw, according to Köppen's climate scale, with the dry season from April to September and the rainy season from October to March (Köppen 1948; Schneider and Teixeira 2001). The temperatures range between 24 and 26°C (Pezzopane *et al.* 2004).

Voucher specimen was collected under the SisBio's license 16752-1 and is deposited in the Mammal Collection of the Museu de Biologia Mello Leitão (MBML 3351). Cranial and body measurements were taken, in millimeters, according to Vizotto and Taddei (1973), as follows: forearm length 70.8; tibia length 27.7; thumb length 9.7; ear length 31.5; tragus length 6.6; calcar length 19.9; greatest length of the skull 31.7; condylobasal length 27.8; condylocanine length 26.5; braincase breadth 11.6; postorbital breadth 5.7; mastoid breadth 15.1; lower toothrow length 12.8; cingular-canine external breadth 8.1; breadth across molars 12.1; mandibular length 20.6; upper toothrow length 11.1; zygomatic width 17.5; interorbital width 9.1; palatal width 6.0; palatal length 12.6; and basal length 22.8.

The present record of *P. elongatus* increases the mammal list of the state Espírito Santo, which recently included a new species described from the region of Linhares (Nogueira *et al.* 2012). These new records reinforce the need of additional bat samplings in the Atlantic Forest of the Espírito Santo state, which still has poorly studied areas (Bergallo *et al.* 2003; Moreira *et al.* 2008; Mendes *et al.* 2010).




FIGURE 1. (A) Map with locality records for *Phyllostomus elongatus* in Brazil. The star refers to the new record and the circles refer to existing records, which are also marked by numbers, as follows: (1) Macapá, Amapá (Mok and Lacey 1980); (2) Utinga, Pará (Handley 1967); (3) Recife, Pernambuco (Souza-Lopes 1978); (4) Rio Largo, Alagoas (Vieira 1955); (5) Salvador, Bahia (Vieira 1955); (6) Teresópolis, Rio de Janeiro (Vieira 1955); (7) Rio Branco, Mato Grosso (Cabrera 1958); (8) BR-364, Rondônia (Marques 1989); (9) Parque Nacional da Serra do Divisor, Acre, (Nogueira et al. 1999); (10) Ilha do Maracá, Roraima (Robinson 1998); and (11) Manaus, Amazônia (Reis and Guillaumet 1983). (B) The place of the new record in Espírito Santo (marked with a star).

ACKNOWLEDGMENTS: We thank Ryan Laughlin and Beatriz Rohden Becker for the English review and the Instituto de Pesquisas da Mata Atlântica (IPEMA) for the transfer of funds provided by Kreditanstalt für Wiederaufbau (KfW), by means of Brazilian Biodiversity Fund (FUNBIO), that implements the project called “Funbio n°120: Apoio à criação de unidades de conservação no Estado do Espírito Santo: um processo participativo integrando as comunidades locais – Saberes da Mata”.

LITERATURE CITED

- Bergallo, H.G. and C.E.L. Esbérard, M.A.R. Mello, V. Lins, R. Mangolin, G.G.S. Melo and M. Baptista. 2003. Bat species richness in Atlantic Forest: what is the minimum sampling effort? *Biotropica* 35(2): 278–288.
- Bernard, E., V.C. Tavares and E. Sampaio. 2011. Updated compilation of bat species (Chiroptera) for the Brazilian Amazonia. *Biota Neotropica* 11(1): 35–46.
- Cabrera, A. 1958. Catálogo de los mamíferos de América del Sur. *Revista del Museo Argentino de Ciencias Naturales* 4(1): 1-308.
- Handley, C. O. Jr. 1967. Bats of the canopy of an Amazonian forest. *Atas do simpósio sobre a biota Amazônica* 5: 211–1.
- Husson, A.M. 1962. The bats of Suriname. *Zoologische Verhandelingen. Rijksmuseum Natural History*, Leiden 58: 1–282.
- Köppen, W. 1948. *Climatología: com um estudo de los climas de la tierra*. México: Fondo de Cultura Económica. 477 p.
- Marinho-Filho, J. 1996. Distribution of bat diversity in the southern and southeastern Brazilian Atlantic Forest. *Chiroptera Neotropical* 2(2): 51–54.
- Marques, S.A. 1989. *Levantamento faunístico da área sob influência da BR-364 (Cuiabá-Porto Velho)*. Brasília: SCT/PR CNPq, Programa Polonoroeste. Relatório de Pesquisa n.4. 49 p.
- Mendes, P., T.B. Vieira, M. Oprea, S.R. Lopes, A.D. Ditchfield and M. Zortéa. 2010. O conhecimento sobre morcegos (Chiroptera: Mammalia) do estado do Espírito Santo, sudeste do Brasil. *Papéis Avulsos de Zoologia* 50(22): 363–373.
- Moreira, D.O., B.R. Coutinho and S.L. Mendes. 2008. O status do conhecimento sobre a fauna de mamíferos do Espírito Santo baseado em registro de museus literatura científica. *Biota Neotropica* 8(2): 163173.
- Moreira, D.O. and S.L. Mendes. 2010. Diversidade de mamíferos em ecossistemas costeiros do estado do Espírito Santo; p. 59–74 In L.M. Pessoa, W.C. Tavares and S. Siciliano (org.). *Mamíferos das Restingas e Manguezais do Brasil*. Rio de Janeiro: Sociedade Brasileira de Mastozoologia.
- Mok, W.Y. and L.A. Lacey. 1980. Algumas considerações ecológicas sobre morcegos vampiros na epidemiologia da raiva humana na Bacia Amazônica. *Acta Amazonica* 10: 335–42.
- Nogueira, M.R., A. Pol and A.L. Peracchi. 1999. New records of bats from Brazil with a list of additional species for the chiropteran fauna of the state of Acre, western Amazon basin. *Mammalia* 63: 363–368.
- Nogueira, M.R., A.L. Peracchi and R. Moratelli. 2007. Subfamília Phyllostominae; p. 61–97 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Morcegos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina.
- Nogueira, M.R., I.P. Lima, A.L. Peracchi and N.B. Simmons. 2012. New genus and species of nectar-feeding bat from the Atlantic Forest of Southeastern Brazil (Chiroptera: Phyllostomidae: Glossophaginae). *American Museum Novitates* 3747: 1–30.
- Pezzopane, J.E.M., E.A. Santos, M.M. Eleutério, E.F. Reis and A.R. Santos. 2004. Espacialização da temperatura do ar no estado do Espírito Santo. *Revista Brasileira de Agrometeorologia* 12(1): 151–158.
- Pinto, D. and H. Ortêncio-Filho. 2006. Dieta de quatro espécies de filostomídeos frugívoros (Chiroptera, Mammalia) do Parque Municipal do Cinturão Verde de Cianorte, Paraná, Brasil. *Chiroptera Neotropical* 12(2): 274–279.
- Reis, N.R. and J.L. Guillaumet. 1983. Les chauves-souris frugivores de la région de Manaus et leur rôle dans la dissémination des espèces végétales. *Revue d'Ecologie (Terre Vie)* 38: 147–69.
- Reis, N.R., A.L. Peracchi, W.A. Pedro and I.P. LIMA. 2011. *Mamíferos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina. 441 p.
- Robinson, F. 1998. The bats of the Ilha de Maracá; p.165-188 In J.A. Ratter and W. Milliken (ed.). *Maracá: the biodiversity and environment of an Amazonian Rainforest*. London: Wiley.
- Rodrígues-Posada, M.E. and P. Sánchez-Palomino. 2009. Taxonomía del género *Phyllostomus* (Chiroptera: Phyllostomidae) en Colombia. *Mastozoología Neotropical* 16(1): 153–168.
- Rolim, S.G., N.M. Ivanauskas, R.R. Rodrigues, M.T. Nascimento, J.M.L. Gomes, D.A. Folli and H.Z. Couto. 2006. Composição Florística do estrato arbóreo da Floresta Estacional Semidecidual na Planície Aluvial do rio Doce, Linhares, ES, Brasil. *Acta Botânica Brasilica* 20(3): 549–561.

- Rolim, S.G. and A.G. Chiarello. 2004. Slow death of Atlantic forest trees in cocoa agroforestry southeastern Brazil. *Biodiversity and Conservation* 13: 2679–2694.
- Sambuichi, R.H.R. 2002. Fitossociologia e diversidade de espécies arbóreas em cabruca (Mata Atlântica raleada sobre plantação de cacau) na região sul da Bahia. Brasil. *Acta Botânica Brasilica* 16: 89–101.
- Souza-Lopes, M. J. de. 1978. Cariótipo de duas espécies de morcegos de Pernambuco (Chiroptera-Phyllostomatidae). *Revista Nordestina de Biologia* 1: 113–17.
- Schneider, J.A.P. and R.L. Teixeira. 2001. Relacionamento entre anfíbios anuros e bromélias da Restinga de Regência, Linhares, Espírito Santo, Brasil. *Iheringia, Série Zoologia* 91: 41–48.
- Tavares, V.C., R. Gregorin, and A.L. Peracchi. 2008. Diversidade de morcegos no Brasil: lista atualizada com comentários sobre distribuição e taxonomia; p. 417–426 In S.M. Pacheco, R.V. Marques and C.E.L. Esbérard (org.). *Morcegos no Brasil: biologia, sistemática, ecologia e conservação*. Porto Alegre: Armazém Digital.
- Vieira, C.O.C. 1942. Ensaio monográfico sobre os quirópteros do Brasil. *Arquivos de Zoologia* 3: 219–471.
- Vieira, C. O. C. 1955. Lista remissiva dos mamíferos do Brasil. *Arquivos de Zoologia* 8: 341–474.
- Vizotto, L.D. and V.A. Taddei. 1973. Chave para determinação de quirópteros brasileiros. *Revista da Faculdade de Filosofia, Ciências e Letras de São José do Rio Preto, Boletim de Ciências* 1: 1–72.
- Williams, S.L. and H.H. Genoways. 2008. Subfamily Phyllostominae Gray, 1825; p. 255–300 In Gardner, A.L. (ed.). *Mammals of South America. Volume 1: Marsupials, Xenarthrans, Shrews, and Bats*. Chicago and London: The University of Chicago Press.

RECEIVED: October 2012

ACCEPTED: June 2013

PUBLISHED ONLINE: September 2013

EDITORIAL RESPONSIBILITY: Marcelo R. Nogueira