

# Aquatic macrophytes of Northeastern Brazil: Checklist, richness, distribution and life forms

Edson Gomes de Moura-Júnior<sup>1</sup>, Liliane Ferreira Lima<sup>1</sup>, Simone Santos Lira Silva<sup>2</sup>, Raíssa Maria Sampaio de Paiva<sup>3</sup>, Fernando Alves Ferreira<sup>4</sup>, Carmen Silvia Zickel<sup>5\*</sup> and Arnildo Pott<sup>4</sup>

- 1 Graduate student (PhD) in Plant Biology, Universidade Federal de Minas Gerais, Biology Department. Av. Antônio Carlos, 6627. CEP 31270-901. Belo Horizonte, MG, Brazil.
  - 2 Graduate student (PhD) in Botany, Universidade Federal Rural de Pernambuco, Biology Department. Av. Dom Manoel de Medeiros, s/n°, Dois Irmãos. CEP 52171-900. Recife, PE, Brazil.
  - 3 Graduate student (MSc) in Natural Resource, Universidade Federal de Roraima, Biology Department. Av. Capitão Ene Garcez, 2413, Aeroporto, Boa Vista. CEP 69304-000. Roraima, RR, Brazil.
  - 4 Universidade Federal do Mato Grosso do Sul, Program in Plant Biology, Center for Biological Sciences and Health, Biology Department. Cidade Universitária, s/n - CEP 79070-900. Campo Grande, MS, Brazil.
  - 5 Universidade Federal Rural de Pernambuco, Program in Botany, Biology Department. Av. Dom Manoel de Medeiros, s/n°, Dois Irmãos. CEP 52171-900. Recife, PE, Brazil.
- \* Corresponding Author. E-mail: [zickelbr@yahoo.com](mailto:zickelbr@yahoo.com)

**ABSTRACT:** Aquatic plants have great influence on the structure and dynamics of aquatic ecosystems, thereby contributing considerably to biodiversity. In Brazil, knowledge of the biodiversity of aquatic macroflora is still limited. We present a checklist of aquatic macrophytes occurring in the northeastern region of Brazil through a bibliographic search. We recorded a total of 412 species, 217 genera and 72 families. The most representative families were Cyperaceae (70 species), Poaceae (38), Fabaceae (27) and Asteraceae (20). The States with highest number of species were Pernambuco (370), Bahia (360), Ceará (267) and Paraíba (261). The best-represented life forms were amphibious (193 species) and emergent (100). The aquatic flora of Northeastern Brazil exhibits high species richness; however, there is a scarcity of records of aquatic macrophytes for the States of Alagoas, Rio Grande do Norte, Maranhão, Sergipe and Piauí.

## INTRODUCTION

Wetlands represent one of the vital components in the maintenance of the global balance, considering the high diversity of species found in these ecosystems (Gopal and Junk 2000). Part of this biodiversity can be explained by the presence of aquatic plants, which are known to exert large structural and metabolic influences on the environment. These plants participate in the production of organic matter and nutrient cycling in the water, constitute the base of the food chain as the primary source of energy, protect the margins of rivers, ponds and lakes against erosion, and serve as shelter and protection for aquatic and/or amphibious organisms (Pott and Pott 2000; Murphy *et al.* 2003; Thomaz and Cunha 2010). On the other hand, the excessive growth of aquatic vegetation (mostly ruderal species) can become a problem for water use (navigation and electric power generation), contributing to loss of diversity of submersed plants and phytoplankton (Lembi 2009; Thomaz and Cunha 2010).

Throughout the world, the first studies on aquatic macrophytes were performed in temperate ecosystems and only later began to be performed in the tropics (Thomaz and Bini 2003; Pompêo and Moschini-Carlos 2003). Due to this geographic bias, the existing literature on aquatic plants of tropical regions does not correlate with the biodiversity of the group. Even now, few works address the biodiversity of aquatic macrophytes in African (Raynal-Roques 1980; Obot and Mbagwu 2008) and Asian countries (Subramanyam 1962). In the Americas, studies have been performed in Mexico (Ramos and Novelo 1993; Bonilla-Barbosa and Novelo 1995), in Argentina (Neiff

1982; Lahitte and Hurrell 1996; Miretzky *et al.* 2004), in Paraguay (Mereles *et al.* 1992; Neiff *et al.* 2000), and in Venezuela (Vilarrubia and Cora 1993).

In Brazil, the analysis of biodiversity of aquatic macrophytes has been restricted to the South, Southeast and Central-West regions and some sporadic studies developed in other regions (Thomaz and Bini 2003). Irgang and Gastal-Jr. (1996) published a list with identification keys and photos of approximately 400 species of aquatic macrophytes of the coastal plain of Rio Grande do Sul. Scremin-Dias *et al.* (1999) produced an identification guide of ca. 50 species of aquatic plants occurring in Bonito and Bodoquena. Pott and Pott (2000) presented brief morphological descriptions, ecological aspects, geographic distribution and simplified identification keys of 246 species of aquatic macrophytes found in the Pantanal. Recently, Amaral *et al.* (2008) published a field guide for aquatic and palustrine plants of the State of São Paulo, including the description of ca. 400 species, plus illustrations.

Knowledge of the aquatic macroflora of the Northeast of Brazil is also quite limited. Some floristic studies were carried out by Matias *et al.* (2003) in Ceará, by Neves *et al.* (2006) in coastal lakes in Bahia, and by França *et al.* (2003), also in Bahia. In the State of Pernambuco, Sobral-Leite *et al.* (2010) presented a checklist of vascular macrophytes, Lima *et al.* (2009) made a floristic survey of herbaria, and Moura-Júnior *et al.* (2009) compared the richness of aquatic macrophytes among water supply reservoirs. Pompêo and Moschini-Carlos (2003) observed that in spite of the growing number of studies, research and discussion

of aquatic macrophytes in congresses, symposia and scientific journals, there are few professionals that are currently fully dedicated to study this community in Brazil.

Although they have yet to cover the diversity of aquatic macrophytes, the floristic studies carried out in Brazil are important because they contribute to the quantification and qualification of the flora of aquatic ecosystems, as well as to the knowledge of geographic distribution of species. In such a context, obtaining a global vision of the quantity of records and of the species richness of aquatic macrophytes can serve as subsidy for ecological studies, can help visualize patterns related to biodiversity and can help elucidate the relation of flora with environmental factors.

Therefore, our study aims to produce a checklist of aquatic macrophytes occurring in the Northeast of Brazil through a bibliographic survey based on inventories in this region. Thereafter, we want to answer the following questions: (i) What is the number of species recorded for the Northeast of Brazil? (ii) What is the number of existing records for each State of the region? (iii) What are the life forms? (iv) What is the floristic similarity between the states? and (v) Which species are considered ruderal or opportunistic?

## MATERIALS AND METHODS

### Study site

Distinguished from other tropical regions by certain peculiar aspects, the Northeast region of Brazil occupies an area of 1,554,257 km<sup>2</sup> (18.25% of Brazil) and contains nine states: Alagoas, Bahia, Ceará, Maranhão, Paraíba, Pernambuco, Piauí, Rio Grande do Norte and Sergipe (IBGE 2002) (Figure 1). This region is characterized by strikingly distinct phytogeographic zones (forest, subhumid and semiarid Caatinga, and middle North palmlands) and an annual regime of rain irregularly distributed between March and August, with the peak in May and June, the dry period being between September and February (IBGE 2002; Silva *et al.* 2002).

Covering a large drainage area of the Northeast region of Brazil, there are important Brazilian hydrographic basins, among which we note the following: São Francisco Basin, which has an area of 643,000 km<sup>2</sup> and covers 521 municipalities of seven states (Bahia, Minas Gerais, Pernambuco, Alagoas, Sergipe, Goiás and Distrito Federal); Parnaíba Basin, which occupies 344,112 km<sup>2</sup> nearly covering the whole state of Piauí and part of the states of Maranhão and Ceará; Oriental Northeastern Atlantic Basin, basically situated in the state of Maranhão and in a small oriental portion of the state of Pará, has an area of 254,100 km<sup>2</sup>; and the Eastern Atlantic Basin, with an area of 374,677 km<sup>2</sup>, within two states of the Northeast (Sergipe and Bahia) and two of the Southeast (Minas Gerais and Espírito Santo) (ANA 2012).

### Data Collection


The checklist was created by a bibliographic search of studies on aquatic plants performed in the Northeast of Brazil, published between August 2000 and 2010. To accomplish these objectives, we adopted the conceptualization of aquatic macrophytes proposed by Cook (1996), in which the author includes plants in which

photosynthetically active organs are either permanently or for several months of the year totally or partially submersed in freshwater or floating in aquatic habitats. More recently, Chambers *et al.* (2008) also included charophytes within the definition of macrophytes.

Aquatic macrophytes have been grouped according to their life forms into submerged, free floating, rooted floating, emergent and amphibious (Irgang *et al.* 1984), and epiphyte (Tur 1972). Usually, life form zoning occurs according to water depth: amphibians on the littoral, emergents on the shallow belts, and others in the deeper zones (Pott *et al.* 2011).

We consulted floristic and/or taxonomic work such as book chapters and national and international scientific articles, as well as theses and dissertations made available by graduate programs in Botany and Ecology (Table 1). You can see the locations studied in some states (Figure 1, Table 1), except those related to study of Matias *et al.* (2010), who were not available. Based on those publications, we compiled the floristic data (including ruderal taxa) of life forms of the aquatic macrophytes by state.

To consolidate the checklist, in addition to the referred bibliography, we verified the species records for each



**FIGURE 1.** Hidrographic map of the Northeast region of Brazil, with the inventoried sites (in red). AL = State of Alagoas; BA = State of Bahia; CE = State of Ceará; MA = State of Maranhão; PB = State of Paraíba; PE = State of Pernambuco; PI = State of Piauí; RN = State of Rio Grande do Norte; SE = State of Sergipe.

state of Northeast of Brazil and consulted the biological collections on the data system of “speciesLink” (SPLINK 2011), which has information from the main Brazilian herbaria.

The species list followed the classification of families proposed by APG III (2009) for angiosperms, by Smith *et al.* (2006) for pteridophytes, Buck and Goffinet (2000) for bryophytes, and Wood and Imahori (1964) for macroalgae. Plant names and respective authors were checked by consulting the data bank of the Missouri Botanical Garden (MOBOT 2011). The characterization of species as ruderal was based on “Plantas daninhas do Brasil” (Lorenzi 2008), which considers species ruderal if they are the first to colonize disturbed wetlands.

#### Data analyses

The evaluation of the floristic similarity between the records of different states was accomplished through a

cluster analysis. The date of floristic composition was subjected to an analysis of similarity by the Jaccard index (Magurran 2004) and ordered by the WPGMA method, using the software PRIMER 6.0 (Clarke and Gorley 2006). To identify significantly similar groups ( $p < 0.05$ ), we used the Simprof method of randomization with 1000 replications, using the software PRIMER 6.0 (Clarke and Gorley 2006).

We compiled all records of aquatic macrophytes in a matrix of presence and absence of species. The frequency of occurrence of each species ( $F$ ) was estimated through the equation:

$$F(i) = \frac{n_i}{N}$$

where:

$n_i$  = number of sites where the species  $i$  was found, and  
 $N$  = number of sampled sites

**TABLE 1.** List of the studies used to elaborate the checklist. AL = State of Alagoas; BA = State of Bahia; CE = State of Ceará; MA = State of Maranhão; PB = State of Paraíba; PE; State of Pernambuco; PI = State of Piauí; RN = State of Rio Grande do Norte; SE = State of Sergipe; NO = Northeastern.

AUTHOR(S)	STUDY REGION	TYPE OF ENVIRONMENT
Mathias and Nunes (2001)	CE	Coastal lagoon
França <i>et al.</i> (2003)	BA	Artificial pond
Matias <i>et al.</i> (2003)	CE	Coastal lagoon
Neves <i>et al.</i> (2006)	BA	Coastal lagoon
Pedro <i>et al.</i> (2006)	PB	River
Moura-Júnior <i>et al.</i> (2009)	PE	Artificial pond
Lima <i>et al.</i> (2009)	PE	Shallow lakes, rivers, floodplains, oxbow lakes, temporary ponds, permanent ponds, Coastal lagoons, lakes, permanent and temporary swamps, Artificial ponds
Nascimento (2009)	PE	Artificial pond
Henry-Silva <i>et al.</i> (2010)	RN	Rivers
Matias (2010)	NO	Shallow lakes, rivers, floodplains, oxbow lakes, temporary ponds, permanent ponds, Coastal lagoons, lakes, permanent and temporary swamps, Artificial ponds
Moura-Júnior <i>et al.</i> (2010)	BA	Artificial pond
Silva and Zickel (2010)	PE	Artificial pond
Sobra-Leite <i>et al.</i> (2010)	PE	Shallow lakes, rivers, floodplains, oxbow lakes, temporary ponds, permanent ponds, Coastal lagoons, lakes, permanent and temporary swamps, Artificial ponds

## RESULTS AND DISCUSSION

We compiled 412 species of 217 genera and 72 families (Table 2). According to Agostinho *et al.* (2005), the vascular aquatic flora of wetlands in Brazil is estimated to 600 species, which makes the number of taxa identified in our study somewhat representative. The high richness of aquatic macrophytes for the Northeast can be attributed to seasonal and hydrological influence on the aquatic ecosystems of the region (França *et al.* 2003; Neves *et al.* 2006; Moura-Júnior *et al.* 2009; Campelo *et al.* 2012)

The families with highest richness of aquatic macrophytes in the Northeast of Brazil were Cyperaceae, with 70 species, Poaceae (36 spp.), Fabaceae (27 spp.), Alismataceae (23 spp.) and Asteraceae (20 spp.) (Table 2). The floristic representativeness of Cyperaceae and Poaceae was also recorded in studies on aquatic macrophytes of the other Brazilian regions, including the North (Junk and Piedade 1993), South (Irgang and Gastal-Jr. 1996; Mormul *et al.* 2010) and Southeast (Ferreira *et al.* 2010). The present estimate is *ca.* 700 species of Cyperaceae and 1500 of Poaceae in Brazil (Souza and Lorenzi 2008). According to Goetghebeur (1998), the taxonomic abundance of these families must be related to their efficiency in vegetative

propagation, with underground systems formed by rhizomes or stolons. The richness of Asteraceae and Fabaceae in our study converged with the result reported by Lima *et al.* (2009) in the checklist of aquatic macrophytes of the state of Pernambuco, wherein both of these families as well as the two previous families are the four richest. It is known that species of Asteraceae are particularly common in several vegetation types in Brazil (Souza and Lorenzi 2008), likely due to the morphological adaptations of the fruits, which have a persistent pappus transformed into a dispersal structure, primarily for anemochory and zoochory (Heiden *et al.* 2007). Considering Fabaceae, we believe that the morphological, ecophysiological and reproductive plasticity of the sub-families Faboideae, Caesalpinioideae and Mimosoideae can explain the wide specific richness of the family in ecotonal areas and in wetlands of the Northeast. As was discussed for Cyperaceae, Poaceae, Fabaceae and Asteraceae, the high representation of floristic Alismataceae in northeastern Brazil may also be related to morphological adjustments of their representatives. According to Matias (2010), individuals of *Echinodorus* and *Sagittaria* (the most common representative of Alismataceae in the study area) have

an underground system composed of perennial rhizomes with buds that can withstand long periods of drought and thus can colonize aquatic ecosystems permanently and/or intermittently until ecotone environments (swamps) are created.


The states which presented the highest number of records of species were Pernambuco with 369, Bahia (360 spp.), Ceará (267 spp.) and Paraíba (261 spp.). Next came the states of Alagoas with 208 species, Rio Grande do Norte (192 spp.), Maranhão (187 spp.), Sergipe (181 spp.) and Piauí (158 spp.). The number of species restricted to only one state also follows the pattern of total richness, the most representative being Pernambuco (24 spp.), Bahia (13 spp.), Ceará (3 spp.) and Rio Grande do Norte (2 spp.) (Table 2).

According to Magurran (1988), the number of inventoried species in an area invariably increases with the size of sampled area and/or the sampling effort. Therefore, the scarce record of aquatic macrophytes for the states of Alagoas, Rio Grande do Norte, Maranhão, Sergipe and Piauí compared to the other states of the Northeast might have been caused by the low number of inventoried areas and by a small range of studied areas. Although some states in the Northeast presented a negative bias regarding the record of aquatic plant species, Pernambuco and Bahia showed similar species richness to that observed in the Amazon region, where 388 species were identified (Junk and Piedade 1993), and higher than the richness of 273 species found in the Pantanal (Pott and Pott 2000).

According to the Simprof test, macrophyte communities that had Jaccard (S) scores above 54% were considered significantly similar in floristic structure (Figure 2). Thus, groups of species of aquatic plants in the states of Alagoas, Bahia, Ceará, Paraíba, Pernambuco and Sergipe can be considered similar (S = 55%) (Figure 2). However, within

the group of those six states, there was the formation of three clusters whose similarity indices were the highest recorded in the analysis: the first included the aquatic macrophytes of the states of Bahia and Pernambuco (S = 79.8%), the second was represented by macrophytes of Ceará and Paraíba (S = 69.3%), and third was formed by hydrophytes of Alagoas and Sergipe (S = 58.5%) (Figure 2). Some studies have shown that the composition and distribution of species of aquatic plants in tropical ecosystems are primarily explained by their hydrological (e.g., elevation, flow) and/or limnological characteristics (e.g., turbidity, conductivity, transparency, pH, temperature and concentration of oxygen, nitrogen and dissolved phosphorus) (Thomaz *et al.* 2003; Murphy *et al.* 2003; Pedro *et al.* 2006; Souza *et al.* 2009; Moura Júnior *et al.* 2011). Most of these researchers concede that changes in hydrological characteristics and/or limnological ecosystem changes, in short, the processes of interaction between species (e.g., competition, facilitation), modify the structure of aquatic communities. In this context, the similarity in the species composition of hydrophytes of Alagoas, Bahia, Ceará, Paraíba, Pernambuco and Sergipe can be explained by similarities in the hydrological and limnological conditions of aquatic ecosystems in these states. Similarly, the floristic dissimilarity of macrophytes of Maranhão, Piauí and Rio Grande do Norte in relation to other states of Northeastern Brazil is most likely related to the differences in hydrological and limnological ecosystem states. Moreover, the lack of information about the species of aquatic plants in the states of Maranhão, Piauí and Rio Grande do Norte may also explain the differences for these flora with the rest of the Northeast, which implies the need for floristic and taxonomic studies for these three states.

Regarding the aquatic macrophytes, which are restricted to only one of the life forms of the Northeast,



**FIGURE 2.** Dendrogram of floristic similarity (Jaccard coefficient) of aquatic macrophytes between the states, by Simprof randomization test = 54% (with 2,000 replications;  $p < 0,05$ ). AL = State of Alagoas; BA = State of Bahia; CE = State of Ceará; MA = State of Maranhão; PB = State of Paraíba; PE; State of Pernambuco; PI = State of Piauí; RN = State of Rio Grande do Norte; SE = State of Sergipe.

the most representative groups were amphibious and emergent, with 193 and 100 of the species, respectively. Next, there appeared rooted submersed (22 spp.), free floating (18 spp.), rooted floating (16 spp.), free submersed (eight spp.) and a single species of epiphyte. Relative to the plasticity of life forms, 47 species stood out for being amphibious as well as emergent. As for the results of the life forms per state, we also observed a higher number of species categorized as amphibious and emergent when compared to other analyzed categories (Figure 3). The other categories of life forms contained two species (amphibious/rooted floating) and only one species each (emergent/epiphyte/amphibious, emergent/free floating, emergent/rooted floating, emergent/rooted submersed and free floating/rooted floating). The predominance of species that presented amphibious and/or emergent life forms (for the nine states of Northeastern Brazil) was similar to results obtained in various aquatic environments of tropical regions (Pott and Pott 2000; Ferreira *et al.* 2011). We attribute the representativeness of these species to their morphophysiological adjustments, which allow them to survive in the interface between aquatic and terrestrial environments (Sculthorpe 1967; Matias *et al.* 2003). We further suggest that the high richness of amphibious and/or emergent aquatic plants in tropical ecosystems may be related to the number of species of Cyperaceae and Poaceae (Ribeiro *et al.* 2011).


In our list, 141 species were considered ruderal or opportunistic, corresponding to 34.2% of the number of aquatic macrophytes cited for the Northeast. Such species are grouped into 44 families, of which Poaceae (with 23 species), Asteraceae (14), Cyperaceae (14), Fabaceae (12) and Pontederiaceae (6) had the highest number of species (Table 2). Some studies have reported that the proliferation of aquatic ruderal weeds is primarily related to the dumping of industrial, urban and agricultural nutrient-rich nitrogen and phosphate (anthropogenic eutrophication) in rivers, lakes, ponds and reservoirs (Pott and Pott 2000; Esteves 2011). It is also known that eutrophication is an agent that interferes with interspecific interactions of natural selection, causing a decline in the biodiversity of aquatic communities (Tundisi and Tundisi 2008; Bicudo

*et al.* 2010; Esteves 2011). From this perspective, the growing process of deterioration in the quality of aquatic limnological ecosystems in Northeastern Brazil (Cyril *et al.* 2010; Silva and Zickel 2010) can be considered to be a factor threatening the aquatic biodiversity of the region, especially if academic studies and/or governmental strategies for the conservation of these resources are not developed.

The occurrence of ruderal species per state was the following: Maranhão (43.85%), Sergipe (42.85%), Alagoas (42.78%), Rio Grande do Norte (41.96%), Ceará (41.04%), Piauí (39.87%), Paraíba (39.31%), Bahia (36.16%) and Pernambuco (35.58%). It is worth pointing out that of the 71 species common to all states of the Northeast, 39 species were considered ruderal, while of the 41 taxa considered restricted, *i.e.*, that occurred only in a single state, six were ranked as ruderal (Table 2). As we observed in our study, other researchers also recorded a large percentage of ruderal species, *e.g.*, Kita and Souza (2003) for the floodplain of the upper Paraná, Bove *et al.* (2003) in a temporary aquatic environment of the coastal plain in Rio de Janeiro, and Henry-Silva *et al.* (2010) for the hydrographic basin of Apodi/Mossoró, at 29%, 23% and 47.5%, respectively. The high representativeness of species considered ruderal or opportunistic can be attributed to some features favoring dispersion and survival, such as high adaptation and resistance, as well as longevity and good seed dispersion (Souza and Lorenzi 2008). In addition, the major weedy species of aquatic macrophytes that occur in Brazil spread vegetatively (plus via seed). This factor facilitates the proliferation and exaggerated abundance of some species in many aquatic environments. Many species thus become an ecological problem (Lembi 2009; Thomaz and Cunha 2010).

Moura *et al.* (2009) cited the weedy species *Egeria densa*, *Ceratophyllum demersum*, *Brachiaria mutica*, *B. subquadripara*, *Eichhornia crassipes*, *Pistia stratiotes*, *Polygonum lapathifolium*, *Echinochloa polystachya* and *Salvinia auriculata* as the species that have the most impact in the State of São Paulo because they can hinder multiple uses of water resources, such as the generation of electricity, irrigation, navigation, fishing and recreation. These species, except *Urochloa arrecta* (above cited as *Brachiaria subquadripara*, although not a synonym) and *Polygonum lapathifolium*, have all been recorded in Northeastern Brazil.

We verified that the existing aquatic flora in Northeastern Brazil exhibits high species richness, a fact that is easily observed by the number of recorded taxa. The record of species varied among states and among categories of life forms according to the amount of sampled areas and/or collection effort in each state. Although the Northeast presented a high richness of aquatic plants (34,2% of which were classified as ruderal), we observed a scarcity in the records of aquatic plants for the states of Alagoas, Rio Grande do Norte, Maranhão, Sergipe and Piauí. We detected a need for more data collection and research directed at the aquatic community. We suggest efforts on floristic and taxonomic work be performed in the Northeast of Brazil to gather information to better determine the contribution of aquatic macrophytes present in wetlands of this region.



**FIGURE 3.** Life forms of aquatic macrophytes by state in Northeast Brazil. AL = State of Alagoas; BA = State of Bahia; CE = State of Ceará; MA = State of Maranhão; PB = State of Paraíba; PE, State of Pernambuco; PI = State of Piauí; RN = State of Rio Grande do Norte; SE = State of Sergipe.

**TABLE 2.** List of macrophytes by life forms (LF) and percent occurrence (PO%), recorded in the states of Northeast Brazil. (+) = occurrence; (-) = absence; (\*) = ruderal specie; AL = State of Alagoas; BA = State of Bahia; CE = State of Ceará; MA = State of Maranhão; PB = State of Paraíba; PE; State of Pernambuco; PI = State of Piauí; RN = State of Rio Grande do Norte; SE = State of Sergipe; AM = amphibious; EM = emergent; EP = epiphyte; FF = free floating; FS = free submerged; RF = rooted floating; and RS = rooted submerged.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<b>Acanthaceae</b>											
<i>Dicliptera ciliaris</i> Juss.	AM	-	+	+	-	+	+	-	+	-	0.23
<i>Dyschoriste maranhonis</i> Kuntze	EM	-	+	-	-	-	+	-	-	-	0.09
<i>Hygrophila costata</i> Nees	EM	-	+	-	-	-	+	-	-	+	0.14
<i>Justicia aequilabris</i> (Nees) Lindau	EM	-	+	+	-	+	+	-	+	-	0.23
<i>Justicia laevilinguis</i> (Ness) Lindau	AM	-	+	-	-	+	+	-	-	-	0.14
<i>Nelsonia brunelloides</i> (Lam.) Kuntz	AM	-	+	-	-	+	+	-	-	-	0.14
<i>Ruellia bahiensis</i> (Nees) Morong *	AM	+	+	+	-	+	+	-	+	+	0.32
<i>Ruellia paniculata</i> L.	AM	-	-	-	-	-	+	-	+	-	0.09
<b>Aizoaceae</b>											
<i>Sesuvium portulacastrum</i> (L.) L.	AM	+	+	+	+	+	+	-	+	+	0.37
<b>Alismataceae</b>											
<i>Echinodorus andrieuxii</i> (Hook et Arn.) Small	EM	-	+	+	-	-	+	+	-	-	0.18
<i>Echinodorus floribundus</i> (Seub.) Seub. *	EM	-	+	-	-	-	+	-	-	-	0.09
<i>Echinodorus glandulosus</i> Rataj *	EM	-	+	+	-	-	+	-	-	-	0.14
<i>Echinodorus grandiflorus</i> (Cham. and Schltld.) Micheli *	EM	-	-	+	-	-	+	-	+	-	0.14
<i>Echinodorus lanceolatus</i> Rataj	EM	-	+	+	-	-	-	-	-	-	0.09
<i>Echinodorus longipetalus</i> Micheli	EM	-	+	+	+	+	+	+	+	+	0.37
<i>Echinodorus macrophyllus</i> (Kunth) Micheli	EM	+	+	+	-	+	-	-	-	-	0.18
<i>Echinodorus palaefolius</i> (Nees and Mart.) J.F. Macbr.	EM	-	+	-	-	-	-	-	-	-	0.05
<i>Echinodorus paniculatus</i> Micheli	EM	-	+	-	-	-	+	+	-	+	0.18
<i>Echinodorus pubescens</i> (Mart.) Seub. and Warm.	EM	-	+	+	-	+	+	+	-	+	0.27
<i>Echinodorus ranunculoides</i> (L.) Engelm.	EM	-	+	+	-	-	+	+	-	+	0.23
<i>Echinodorus reticulatus</i> R.R. Haynes and Holm-Niels.	EM	-	+	-	-	-	-	-	-	-	0.05
<i>Echinodorus scaber</i> Rataj	EM	-	+	+	+	-	+	+	-	+	0.27
<i>Echinodorus subalatus</i> (Mart.) Griseb.	EM	+	+	+	+	+	+	+	+	-	0.37
<i>Echinodorus tenellus</i> (Mart. ex Schult. and Schult. f.) Buchenau	AM	-	+	+	+	+	+	+	-	+	0.32
<i>Hydrocleys martii</i> Seub.	RF	+	+	+	-	+	+	+	+	-	0.32
<i>Hydrocleys nymphoides</i> (Willd.) Buchenau	RF	+	+	+	-	+	+	-	+	+	0.32
<i>Hydrocleys parviflora</i> Seub.	RF	-	-	-	-	-	-	-	+	-	0.05
<i>Limnocharis flava</i> (L.) Buchenau *	EM/AM	-	+	+	+	+	+	+	+	-	0.32
<i>Sagittaria guayanensis</i> Kunth *	EM	-	+	+	+	-	+	+	-	+	0.27
<i>Sagittaria lancifolia</i> L.	EM	+	+	-	-	+	-	-	-	+	0.18
<i>Sagittaria planitiana</i> G. Agostini	EM	-	+	+	-	-	-	-	-	-	0.09
<i>Sagittaria rhombifolia</i> Cham.	EM	-	+	-	-	-	+	-	-	-	0.09
<b>Amaranthaceae</b>											
<i>Alternanthera philoxeroides</i> (Mart.) Griseb. *	EM/AM	+	+	-	-	-	+	-	+	-	0.18
<i>Alternanthera tenella</i> Colla *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Blutaparon portulacoides</i> (A. St.-Hil.) Mears	AM	+	+	+	+	+	+	-	+	+	0.37
<i>Gomphrena demissa</i> Mart.	AM	-	+	+	-	+	+	+	+	+	0.32
<i>Salicornia gaudichaudiana</i> Moq.	AM	-	-	-	-	-	-	-	+	-	0.05
<b>Apiaceae</b>											
<i>Centella asiatica</i> (L.) Urb. *	AM	+	+	-	-	+	+	-	-	-	0.18
<i>Lilaeopsis brasiliensis</i> (Glaz.) Affolter	RS	-	-	-	-	-	+	-	-	-	0.05
<b>Apocynaceae</b>											
<i>Ditassa hastata</i> Decne.	AM	+	+	+	+	+	+	+	+	+	0.41
<b>Araceae</b>											
<i>Lemna aequinoctialis</i> Welw.	FF	+	+	+	-	-	+	+	+	-	0.27
<i>Lemna gibba</i> L.	FF	-	-	-	-	-	+	-	-	-	0.05
<i>Lemna valdiviana</i> Phil. *	FF	-	+	-	-	+	+	+	+	-	0.23
<i>Montrichardia linifera</i> (Arruda) Schott	EM	-	+	+	-	+	+	-	+	+	0.27
<i>Philodendron rudgeanum</i> Schott	EM	+	+	-	+	+	+	-	-	-	0.23
<i>Pistia stratiotes</i> L. *	FF	+	+	+	-	+	+	+	+	+	0.37
<i>Spirodela intermedia</i> W. Koch *	FF	-	-	-	-	-	+	-	-	-	0.05
<i>Wolffia brasiliensis</i> Wedd. *	FF	+	+	+	-	+	-	+	+	-	0.27
<i>Wolffiella welwitschii</i> (Hegelm.) Monod	FF	-	+	-	-	-	+	-	-	-	0.09

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<b>Araliaceae</b>											
<i>Hydrocotyle bonariensis</i> Lam. *	RF	+	+	+	-	-	+	-	+	-	0.23
<i>Hydrocotyle leucocephala</i> Cham. and Schtdl. *	RF	+	+	+	-	+	+	-	-	+	0.27
<i>Hydrocotyle ranunculoides</i> L.f.	EM	-	+	-	-	-	+	-	-	-	0.09
<b>Asteraceae (Compositae)</b>											
<i>Acanthospermum hispidum</i> DC. *	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Acmella uliginosa</i> (Sw.) Cass.	AM	+	+	+	+	+	+	+	+	-	0.37
<i>Ageratum conyzoides</i> L. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Blainvillea dichotoma</i> (Murray) Stewart	AM	+	+	+	+	+	-	+	+	+	0.37
<i>Centratherum punctatum</i> Cass. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Conyza bonariensis</i> (L.) Cronquist *	AM	+	+	+	+	+	+	-	+	+	0.37
<i>Delilia biflora</i> (L.) Kuntze*	AM	+	+	+	+	+	+	-	+	+	0.37
<i>Eclipta prostrata</i> (L.) L. *	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Elephantopus mollis</i> Kunth *	AM	+	+	+	+	+	+	+	-	-	0.32
<i>Emilia coccinea</i> (Sims) G. Don *	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Enydra radicans</i> (Willd.) Lack.	AM	-	+	+	-	-	+	-	-	-	0.14
<i>Enydra rivularis</i> Gardner	AM	-	+	-	-	+	+	-	-	-	0.14
<i>Gamochoa americana</i> (Mill.) Wedd. *	AM	-	+	-	-	-	+	-	-	-	0.09
<i>Pluchea sagittalis</i> (Lam.) Cabrera *	AM	+	+	+	+	+	+	-	-	+	0.32
<i>Rolandra fruticosa</i> (L.) Kuntze	AM	+	+	+	+	+	+	+	-	+	0.37
<i>Sonchus oleraceus</i> L. *	EM	+	+	+	-	+	+	-	-	-	0.23
<i>Sphagneticola trilobata</i> (L.) Pruski *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Synedrella nodiflora</i> (L.) Gaertn. *	AM	+	+	+	+	+	+	+	-	+	0.37
<i>Vernonia scorpioides</i> (Lam) Pers. *	AM	+	+	+	+	+	+	-	+	+	0.37
<i>Wedelia alagoensis</i> Baker	AM	+	+	-	-	-	+	-	-	-	0.14
<b>Begoniaceae</b>											
<i>Begonia fischeri</i> Schrank	AM	-	+	-	-	-	+	-	-	-	0.09
<i>Begonia reniformis</i> Dryand.	AM	+	+	+	-	+	+	-	-	+	0.27
<b>Blechnaceae</b>											
<i>Blechnum brasiliense</i> Desv.	EM	+	+	-	-	+	+	-	+	-	0.23
<b>Boraginaceae</b>											
<i>Cordia curassavica</i> (Jacq.) Roem. and Schult.	AM	-	+	-	-	-	-	-	-	-	
<i>Cordia multispicata</i> Cham.	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Cordia superba</i> Cham.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Heliotropium angiospermum</i> Murray	AM	+	+	+	-	+	+	-	+	+	0.32
<i>Heliotropium indicum</i> L. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Heliotropium procumbens</i> Mill. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Tournefortia bicolor</i> Sw.	AM	+	+	-	+	+	+	-	+	-	0.27
<b>Burmanniaceae</b>											
<i>Burmannia capitata</i> (Walter ex J.F. Gmel.) Mart.	AM	+	+	+	-	+	+	+	+	+	0.37
<b>Cabombaceae</b>											
<i>Cabomba aquatica</i> Aubl.	RS	+	+	-	+	-	+	-	-	-	0.18
<i>Cabomba furcata</i> (Schult.) Schult. F.	FF	-	+	-	-	-	+	-	-	-	0.09
<b>Campanulaceae</b>											
<i>Lobelia xalapensis</i> Kunth	AM	-	+	-	-	+	+	-	-	-	0.14
<b>Cannaceae</b>											
<i>Canna glauca</i> L.	EM	-	+	+	+	+	+	-	-	+	0.27
<b>Capparaceae</b>											
<i>Capparis flexuosa</i> (L.) L.	AM	+	+	+	+	+	+	+	+	+	0.41
<b>Caryophyllaceae</b>											
<i>Drymaria cordata</i> (L.) Willd. ex Roem. and Schult. *	AM	+	+	+	-	+	+	-	-	+	0.27
<b>Ceratophyllaceae</b>											
<i>Ceratophyllum demersum</i> L. *	RS	-	-	+	-	-	-	-	+	-	0.09
<i>Ceratophyllum submersum</i> L.	RS	-	+	-	-	-	-	-	+	-	0.09
<b>Characeae</b>											
<i>Chara zeylanica</i> Willd.	RS	+	+	-	-	+	+	-	+	-	0.23
<b>Cleomaceae</b>											
<i>Cleome guianensis</i> Aubl.	AM	+	+	+	+	+	+	-	+	+	0.37
<i>Cleome hassleriana</i> Chodat *	AM	-	+	+	-	-	+	-	-	-	0.14
<i>Cleome spinosa</i> Jacq.*	AM	+	+	+	+	+	+	+	+	+	0.41

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<b>Commelinaceae</b>											
<i>Callisia filiformis</i> (M. Martens and Galeotti) D.R. Hunt	AM	+	+	+	-	+	+	-	+	-	0.27
<i>Commelina erecta</i> L.*	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Commelina schomburgkiana</i> Klotzsch	AM	-	+	-	-	-	-	-	-	-	0.05
<b>Convolvulaceae</b>											
<i>Aniseia cernua</i> Moric.	AM	-	+	-	-	+	-	-	-	-	0.09
<i>Evolvulus filipes</i> Mart.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Ipomoea asarifolia</i> (Desr.) Roem. and Schult.*	EM/AM	+	+	+	+	+	+	+	+	-	0.37
<i>Ipomoea carnea</i> Jacq.	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Ipomoea fistulosa</i> Mart. ex Choisy*	AM	+	+	+	-	-	+	+	-	-	0.23
<i>Ipomoea setosa</i> Ker Gawl.*	AM	-	+	+	-	+	-	-	+	+	0.23
<i>Ipomoea wrightii</i> A. Gray	AM	-	+	-	-	-	+	-	+	-	0.14
<i>Merremia aegyptia</i> (L.) Urb.*	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Merremia umbellata</i> (L.) Hallier f.*	AM	+	+	+	+	+	+	-	+	+	0.37
<b>Cucurbitaceae</b>											
<i>Fevillea trilobata</i> L.	AM	-	+	+	-	+	+	-	-	-	0.18
<b>Cymodoceaceae</b>											
<i>Halodule emarginata</i> Hartog	RS	-	-	-	-	-	+	-	-	-	0.05
<i>Halodule wrightii</i> Asch.	RS	-	+	+	-	-	+	-	+	-	0.18
<b>Cyperaceae</b>											
<i>Becquerelia cymosa</i> Brongn.	AM	+	+	+	+	+	+	-	-	+	0.32
<i>Bulbostylis capillaris</i> (L.) Kunth ex C.B. Clarke*	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Bulbostylis hirtella</i> (Schrad. ex Schult.) Nees ex Urb.	AM	-	+	+	-	+	-	-	-	-	0.14
<i>Bulbostylis junciformis</i> (Kunth) C.B. Clarke	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Cladium jamaicense</i> Crantz	EM	-	+	-	-	+	+	-	-	-	0.14
<i>Cyperus aggregatus</i> (Willd.) Endl.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus amabilis</i> Vahl	AM	+	+	+	+	+	+	-	+	-	0.32
<i>Cyperus articulatus</i> L.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus cayennensis</i> Willd. ex Link	EM/AM	-	+	-	-	-	-	-	-	-	0.05
<i>Cyperus compressus</i> L.	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus cuspidatus</i> Kunth	AM	+	+	+	-	+	+	+	+	+	0.37
<i>Cyperus esculentus</i> L.*	EM/AM	-	+	-	-	-	-	-	+	-	0.09
<i>Cyperus gardneri</i> Nees	EM	-	-	-	+	-	+	-	+	-	0.14
<i>Cyperus giganteus</i> Vahl*	EM	-	+	+	-	-	+	-	-	-	0.14
<i>Cyperus haspan</i> L.	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus hermaphroditus</i> (Jacq.) Standl.	EM/AM	+	+	+	-	+	+	-	+	+	0.32
<i>Cyperus imbricatus</i> Retz.	EM	-	+	+	+	+	+	-	-	-	0.23
<i>Cyperus involucratus</i> Rottb.	EM	-	-	-	-	-	+	-	-	-	0.05
<i>Cyperus iria</i> L.	EM	+	+	+	+	-	+	+	+	+	0.37
<i>Cyperus lanceolatus</i> Poir.*	EM	-	+	-	-	+	-	-	-	-	0.09
<i>Cyperus laxus</i> Lam.	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus ligularis</i> L.	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus luzulae</i> (L.) Rottb. ex Retz.*	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus odoratus</i> L.*	EM	+	+	+	+	+	+	-	+	+	0.37
<i>Cyperus papyrus</i> L.	EM	+	-	-	-	-	+	+	-	+	0.18
<i>Cyperus surinamensis</i> Rottb.	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Cyperus uncinulatus</i> Schrad. ex Nees	EM	+	+	+	-	+	+	+	+	+	0.37
<i>Cyperus virens</i> Michx.	EM/AM	-	+	-	-	-	+	-	+	-	0.14
<i>Dialium guianense</i> (Aubl.) Sandwith	AM	+	+	-	+	+	+	-	-	-	0.23
<i>Diplacrum longifolium</i> (Griseb.) C.B. Clarke	AM	-	+	-	+	+	+	-	-	-	0.18
<i>Eleocharis acutangula</i> (Roxb.) Schult.*	AM	-	+	+	+	+	+	-	+	+	0.32
<i>Eleocharis atropurpurea</i> (Retz.) J. Presl and C. Presl	AM	+	+	+	-	+	+	-	-	-	0.23
<i>Eleocharis barrosii</i> Svenson	AM	-	+	+	-	+	+	-	-	-	0.18
<i>Eleocharis elata</i> Boeckeler	AM	-	+	+	-	-	+	+	-	-	0.18
<i>Eleocharis filiculmis</i> Kunth	AM	-	+	-	+	+	+	+	+	+	0.32
<i>Eleocharis flavescens</i> (Poir.) Urban	EM	-	+	+	-	+	+	-	-	+	0.23
<i>Eleocharis geniculata</i> (L.) Roem. and Schult.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Eleocharis interstincta</i> (Vahl) Roem. and Schult.*	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Eleocharis maculosa</i> (Vahl) Roem. and Schult.	EM	-	+	+	-	+	+	-	-	+	0.23
<i>Eleocharis minima</i> Kunth	EM/AM	+	+	+	+	+	+	+	+	-	0.37



TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<i>Eleocharis mutata</i> (L.) Roem. and Schult.	EM/RS	+	+	+	+	+	+	+	+	+	0.41
<i>Eleocharis nodulosa</i> (Roth) Schult.	EM/AM	-	+	+	-	-	-	-	-	-	0.09
<i>Eleocharis sellowiana</i> Kunth *	AM	-	+	+	+	+	+	-	-	-	0.23
<i>Fimbristylis autumnalis</i> (L.) Roem. and Schult. *	EM	+	+	-	-	-	+	-	-	-	0.14
<i>Fimbristylis cymosa</i> R. Br.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Fimbristylis diphylla</i> (Reitz) Vahl *	EM	-	+	-	-	-	+	-	-	-	0.09
<i>Fuirena umbellata</i> Rottb.	EM/AM	+	+	+	+	+	+	-	+	+	0.37
<i>Kyllinga pumila</i> Michx.	AM	-	+	+	+	+	+	-	-	+	0.27
<i>Kyllinga vaginata</i> Lam.	AM	+	+	+	-	+	+	+	+	+	0.37
<i>Lipocarpha micrantha</i> (Vahl) G.C. Tucker	AM	+	+	+	-	+	+	+	+	+	0.37
<i>Oxycaryum cubense</i> (Poepp. and Kunth) Palla	EM/AM/EP	-	+	+	+	+	+	+	+	-	0.32
<i>Pycneus flavescens</i> (L.) P. Beauv. ex Rchb.	AM	-	+	+	-	-	+	+	-	-	0.18
<i>Pycneus macrostachyos</i> (Lam.) J. Raynal	EM	+	+	+	-	+	+	+	+	+	0.37
<i>Pycneus polystachyos</i> (Rottb.) P. Beauv. *	EM/AM	+	+	+	+	+	+	-	+	+	0.37
<i>Rhynchospora cephalotes</i> (L.) Vahl	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Rhynchospora contracta</i> (Nees) J. Raynal	EM	+	+	+	-	+	+	-	+	+	0.32
<i>Rhynchospora corymbosa</i> (L.) Britton *	EM	+	+	+	+	-	+	-	-	+	0.27
<i>Rhynchospora cyperoides</i> Mart.	AM	-	-	+	-	-	-	-	-	-	0.05
<i>Rhynchospora gigantea</i> Link	EM	+	+	-	-	+	+	-	-	+	0.23
<i>Rhynchospora holoschoenoide</i> (Rich.) Herter	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Rhynchospora pubera</i> (Vahl) Boeckeler	EM	+	+	-	+	+	+	+	-	-	0.27
<i>Rhynchospora riparia</i> (Nees) Boeckeler	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Rhynchospora tenerrima</i> Nees ex Spreng.	EM	-	+	-	+	+	+	+	+	+	0.32
<i>Rhynchospora tenuis</i> Willd. ex Link	EM	-	+	+	+	+	+	+	-	+	0.32
<i>Rhynchospora trispicata</i> (Nees) Schrad. ex Steud.	EM	-	+	-	+	-	+	-	-	-	0.14
<i>Scleria bracteata</i> Cav.	EM	+	+	+	+	+	+	-	+	+	0.37
<i>Scleria hirtella</i> Sw.	AM	-	+	+	-	-	+	+	+	+	0.27
<i>Scleria latifolia</i> Sw.	EM	+	+	+	+	+	+	-	-	+	0.32
<i>Scleria melalaeuca</i> Rchb. ex Schldl. and Cham. *	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Websteria confervoides</i> (Poir.) S.S. Hooper	RS	+	-	-	-	+	+	-	-	-	0.14
<b>Eriocaulaceae</b>											
<i>Eriocaulon aquatile</i> Körn.	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Leiothrix pilulifera</i> (Körn.) Ruhland	AM	+	+	-	-	+	+	-	-	+	0.23
<b>Euphorbiaceae</b>											
<i>Euphorbia hyssopifolia</i> L. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Euphorbia thymifolia</i> L.	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Tonina fluviatilis</i> Aubl.	AM	+	+	+	+	+	+	+	+	-	0.37
<b>Fabaceae</b>											
<i>Aeschynomene evenia</i> C. Wright and Sauvalle	EM	+	+	+	+	+	+	+	-	+	0.37
<i>Aeschynomene filosa</i> Mart.	AM	-	+	+	+	+	+	+	+	-	0.32
<i>Aeschynomene sensitiva</i> Sw.	AM	+	+	+	+	+	+	+	-	+	0.37
<i>Anadenanthera colubrina</i> (Vell.) Brenan	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Centrosema brasilianum</i> (L.) Benth.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Centrosema pubescens</i> Benth.	AM	+	+	+	+	+	+	+	-	-	0.32
<i>Chamaecrista repens</i> (Vogel) H.S. Irwin and Barneby	EM	+	+	+	+	+	+	+	-	+	0.37
<i>Desmodium barbatum</i> (L.) Benth. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Desmodium incanum</i> DC. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Desmodium tortuosum</i> (Sw.) DC. *	AM	-	+	+	+	+	+	+	+	-	0.32
<i>Dioclea grandiflora</i> Mart. ex Benth.	EP	+	+	+	-	+	+	+	+	+	0.37
<i>Indigofera hirsuta</i> L. *	AM	+	+	+	+	-	+	-	-	-	0.23
<i>Lonchocarpus sericeus</i> (Poir.) Kunth ex DC.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Macroptilium lathyroides</i> (L.) Urban *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Mimosa bimucronata</i> (DC.) Kuntze	AM	-	+	-	-	-	+	-	-	-	0.09
<i>Mimosa pudica</i> L. *	AM	+	+	-	-	+	+	-	+	+	0.27
<i>Neptunia plena</i> (L.) Benth. *	AM	+	+	+	+	+	+	+	+	-	0.37
<i>Rhynchosia minima</i> (L.) DC.	AM	-	+	+	+	+	+	+	+	-	0.32
<i>Senna obtusifolia</i> (L.) H.S. Irwin and Barneby *	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Senna uniflora</i> (Mill.) H.S. Irwin and Barneby *	AM	-	+	+	+	+	+	-	-	-	0.23
<i>Stylosanthes angustifolia</i> Vogel	AM	-	+	+	+	+	+	+	+	+	0.37
<i>Stylosanthes gracilis</i> Kunth *	AM	+	+	+	+	+	+	+	+	+	0.41

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<i>Stylosanthes guianensis</i> (Aubl.) Sw. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Stylosanthes scabra</i> Vogel	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Stylosanthes viscosa</i> (L.) Sw. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Vigna luteola</i> (Jacq.) Benth.	AM	-	+	-	-	-	-	-	-	-	0.05
<i>Zornia latifolia</i> Sm.	AM	-	-	+	-	-	-	-	-	-	0.05
<b>Gentianaceae</b>											
<i>Irlbachia alata</i> (Aubl.) Maas	AM	+	+	-	+	+	+	-	-	-	0.23
<i>Irlbachia purpurascens</i> (Aubl.) Maas	AM	+	+	+	-	+	+	-	+	+	0.32
<i>Schultesia guianensis</i> (Aubl.) Malme	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<b>Haloragaceae</b>											
<i>Laurembergia tetrandra</i> (Schott) Kanitz	RS	-	+	-	-	-	+	-	-	-	0.09
<i>Myriophyllum aquaticum</i> (Vell.) Verdc. *	RS	+	+	-	-	-	+	-	-	-	0.14
<b>Hydrocharitaceae</b>											
<i>Apalanthe granatensis</i> (Bonpl.) Planch.	RS	-	+	+	+	+	+	-	+	-	0.27
<i>Egeria densa</i> Planch. *	RS	-	-	+	-	+	+	-	-	+	0.18
<i>Elodea canadensis</i> Michx.	RS	-	-	-	-	-	+	-	-	-	0.05
<i>Halophila decipiens</i> Ostenf.	RS	-	+	-	-	-	+	-	+	-	0.14
<i>Limnobium laevigatum</i> (Humb. and Bonpl. ex Willd.) Heine	FF	-	+	-	-	-	+	-	-	-	0.09
<i>Najas conferta</i> (A. Braun) A. Braun	FS	-	+	+	-	-	+	-	-	-	0.14
<i>Najas guadalupensis</i> (Spreng.) Magnus	FS	-	+	-	-	-	-	-	-	-	0.05
<i>Najas marina</i> L.	RS	-	+	-	-	-	+	-	-	-	0.09
<b>Hydroleaceae</b>											
<i>Hydrolea spinosa</i> L. *	EM	+	+	+	+	+	+	+	+	+	0.41
<b>Juncaceae</b>											
<i>Juncus microcephalus</i> Kunth *	AM	-	+	-	-	-	+	-	-	-	0.09
<b>Lamiaceae</b>											
<i>Hyptis atrorubens</i> Poit. *	AM	+	+	+	+	+	+	+	-	+	0.37
<i>Hyptis suaveolens</i> (L.) Poit.*	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Marsiphanthes chamaedrys</i> (Vahl) Kuntze *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Ocimum campechianum</i> Mill. *	AM	+	+	+	+	+	-	+	-	+	0.32
<i>Vitex megapotamica</i> (Spreng.) Moldenke	AM	-	+	-	-	+	+	+	-	-	0.18
<b>Lentibulariaceae</b>											
<i>Genlisea filiformis</i> A. St.-Hil.	AM	-	+	-	-	+	+	-	-	-	0.14
<i>Utricularia adpressa</i> Salzm. ex A. St.-Hil. and Girard	AM	-	+	+	-	-	-	-	-	-	0.09
<i>Utricularia breviscapa</i> Wright ex Griseb.	FS	-	+	-	-	-	-	-	-	-	0.05
<i>Utricularia foliosa</i> L. *	FS	+	+	+	+	+	+	+	+	+	0.41
<i>Utricularia gibba</i> L.	AM	+	+	+	-	+	+	-	+	-	0.27
<i>Utricularia hydrocarpa</i> Vahl	FS	-	+	+	+	+	+	-	-	-	0.23
<i>Utricularia juncea</i> Vahl	FS	+	+	-	-	+	+	-	-	-	0.18
<i>Utricularia laciniata</i> A. St.-Hil. and Girard	FS	-	-	-	-	-	+	-	-	-	0.05
<i>Utricularia pusilla</i> Vahl	AM	+	+	-	-	-	+	-	-	+	0.18
<i>Utricularia trichophylla</i> Spruce ex. Oliver	FS	-	+	+	-	-	-	-	-	+	0.14
<b>Linderniaceae</b>											
<i>Lindernia crustacea</i> (L.) F. Muell.	AM	+	-	+	+	+	+	-	-	+	0.27
<i>Lindernia microcalyx</i> Pennell and Stehlé	AM	+	+	+	-	-	+	-	-	-	0.18
<i>Lindernia rotundifolia</i> (L.) Alston	AM	-	+	-	-	-	+	-	-	-	0.09
<i>Torenia thouarsii</i> (Cham. and Schltld.) Kuntze	EM/AM	-	+	-	-	-	+	-	+	+	0.18
<b>Loganiaceae</b>											
<i>Spigelia anthelmia</i> L. *	AM	+	+	+	+	+	+	+	+	+	0.41
<b>Lythraceae</b>											
<i>Ammannia latifolia</i> L.	AM	-	+	+	-	+	+	-	+	-	0.23
<i>Cuphea campestris</i> Koehne	EM/AM	-	+	+	-	+	+	+	+	-	0.27
<i>Cuphea carthagenensis</i> (Jacq.) J.F. Macbr. *	EM/AM	-	+	+	-	-	+	-	-	+	0.18
<i>Cuphea circaeoides</i> Sm. ex Sims	EM/AM	+	+	+	-	-	+	-	+	-	0.23
<i>Cuphea racemosa</i> (L.f.) Spreng. *	EM	+	+	-	-	+	+	-	-	+	0.23
<i>Pleurophora anomala</i> (A. St.-Hil.) Koehne	EM/AM	+	+	+	-	+	+	+	+	+	0.37
<i>Rotala mexicana</i> Schltld. and Cham.	EM	-	+	-	-	-	-	-	-	-	0.05
<i>Rotala ramosior</i> (L.) Koehne	AM	-	+	-	-	+	+	-	-	-	0.14
<b>Malvaceae</b>											
<i>Corchorus hirtus</i> L.	AM	+	+	+	+	+	+	-	-	+	0.32

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<i>Pavonia cancellata</i> (L.) Cav. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Sida anomala</i> A. St. Hil.	AM	-	+	+	+	+	-	+	+	-	0.27
<i>Sida galheirensis</i> Ulbr.	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Sida spinosa</i> L. *	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Sidastrum multiflorum</i> (Jacq.) Fryxell	EM	-	+	+	-	+	+	-	-	-	0.18
<i>Triumfetta althaeoides</i> Lam.	EM	+	+	+	+	+	+	-	-	-	0.27
<i>Urena lobata</i> L. *	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Waltheria indica</i> L. *	AM	-	-	+	-	-	-	-	-	-	0.05
<b>Marantaceae</b>											
<i>Thalia geniculata</i> L. *	EM	-	+	+	+	+	+	-	+	-	0.27
<b>Marsileaceae</b>											
<i>Marsilea crotophora</i> D.M. Johnson	RF	-	+	-	-	-	-	-	-	-	0.05
<i>Marsilea deflexa</i> A Braun	AM/RF	-	+	-	-	-	+	-	-	-	0.09
<b>Mayaceae</b>											
<i>Mayaca fluviatilis</i> Aubl.	RS	-	+	-	-	-	+	-	+	-	0.14
<b>Melastomataceae</b>											
<i>Clidemia hirta</i> (L.) D. Don	AM	+	+	+	+	+	+	+	+	+	0.41
<b>Menyanthaceae</b>											
<i>Nymphoides indica</i> (L.) Kuntze *	RF	+	+	+	+	+	+	+	+	+	0.41
<b>Molluginaceae</b>											
<i>Mollugo verticillata</i> L. *	EM	+	+	+	+	+	+	+	+	+	0.41
<b>Nymphaeaceae</b>											
<i>Nymphaea alba</i> L.	RF	-	-	-	-	+	+	+	+	-	0.18
<i>Nymphaea amazonum</i> Mart. and Zucc.	RF	-	+	+	+	+	+	+	+	-	0.32
<i>Nymphaea ampla</i> (Salisb.) DC. *	RF	+	+	+	+	+	+	-	+	-	0.32
<i>Nymphaea capensis</i> Thunb.	RF	-	-	-	-	-	+	-	-	-	0.05
<i>Nymphaea lasiophylla</i> Mart. and Zucc.	RF	+	+	+	-	+	+	-	-	+	0.27
<i>Nymphaea rudgeana</i> G. Mey.	RF	-	+	+	+	+	+	-	-	-	0.23
<b>Ochnaceae</b>											
<i>Sauvagesia erecta</i> L.	EM	+	+	+	+	+	+	+	+	+	0.41
<b>Onagraceae</b>											
<i>Ludwigia erecta</i> (L.) H. Hara	EM/AM	+	+	+	-	+	+	-	+	+	0.32
<i>Ludwigia filiformis</i> (Micheli) Ramamoorthy	AM	-	+	-	-	-	-	-	-	-	0.05
<i>Ludwigia helminthorrhiza</i> (Mart.) H. Hara	RF/FF	-	+	+	+	+	+	-	+	-	0.27
<i>Ludwigia inclinata</i> (L.f.) M. Gómez	AM/RF	-	+	-	+	-	-	-	-	-	0.09
<i>Ludwigia leptocarpa</i> (Nutt.) H. Hara *	AM	-	+	+	+	+	+	+	+	+	0.37
<i>Ludwigia linifolia</i> Poir.	EM	-	-	-	-	-	+	+	-	-	0.09
<i>Ludwigia octovalvis</i> (Jacq.) P.H. Raven *	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Ludwigia peploides</i> (Kuntze) P.H. Raven	AM	-	+	-	-	+	+	-	+	-	0.18
<i>Ludwigia suffruticosa</i> Walter *	EM	-	-	-	-	-	+	-	+	-	0.09
<b>Orchidaceae</b>											
<i>Anacheilium alagoense</i> (Pabst) Pabst, Moutinho ex A.V. Pinto	EM/AM	-	-	-	-	-	+	-	-	-	0.05
<i>Epidendrum tridactylum</i> Lindl.	EM/AM	-	+	-	-	-	+	-	-	-	0.09
<i>Habenaria pratensis</i> (Lindl.) Rchb. f.	EM	+	+	-	+	+	+	+	-	+	0.32
<i>Habenaria repens</i> Nutt.	EM	-	+	-	-	+	+	-	-	-	0.14
<b>Orobanchaceae</b>											
<i>Melasma melampyroides</i> (Rich.) Pennell	EM/AM	-	+	-	+	+	+	-	-	+	0.23
<b>Passifloraceae</b>											
<i>Piriqueta racemosa</i> (Jacq.) Sweet	AM	-	+	-	-	+	-	-	-	-	0.09
<b>Plantaginaceae</b>											
<i>Angelonia gardneri</i> Hook.	EM	+	+	+	-	+	+	-	+	+	0.32
<i>Bacopa aquatica</i> Aubl.	EM/AM	+	+	+	-	-	+	+	-	-	0.23
<i>Bacopa monnieri</i> (L.) Wettst.	AM	-	+	+	+	+	+	-	+	+	0.32
<i>Bacopa stricta</i> (Schrad.) Edwall	AM	-	+	-	-	-	+	-	-	-	0.09
<i>Micranthemum umbrosum</i> (J.F. Gmel.) S.F. Blake *	AM	-	+	+	-	+	+	-	-	-	0.18
<i>Scoparia dulcis</i> L.*	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Stemodia durantifolia</i> (L.) Sw.	EM	-	+	+	+	+	+	-	-	-	0.23
<i>Stemodia maritima</i> L.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Stemodia pratensis</i> (Aubl.) C.C. Cowan	AM	+	+	+	-	+	+	-	+	-	0.27

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<b>Poaceae</b>											
<i>Antheophora hermaphrodita</i> (L.) Kuntze	AM	+	+	+	-	+	+	+	+	+	0.37
<i>Brachiaria brizantha</i> (Hochst. Ex A.Rich.) Stapf*	EM	-	+	-	-	+	+	-	-	-	0.14
<i>Brachiaria decumbens</i> (Stapf) R.D. Webster*	AM	-	+	+	-	-	+	-	+	-	0.18
<i>Brachiaria mutica</i> (Forssk.) Stapf*	EM	+	+	-	-	+	+	-	-	-	0.18
<i>Brachiaria plantaginea</i> (Link) Hitchc.*	EM	-	+	+	+	+	+	+	-	-	0.27
<i>Cenchrus echinatus</i> L.*	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Chloris barbata</i> Sw.*	EM/AM	+	+	+	-	+	+	+	+	+	0.37
<i>Chloris exilis</i> Renvoise	AM	+	+	-	+	-	+	-	-	-	0.18
<i>Cynodon dactylon</i> (L.) Pers.*	EM	+	+	+	+	+	+	+	+	-	0.37
<i>Dactyloctenium aegyptium</i> (L.) Willd.*	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Digitaria ciliaris</i> (Retz.) Koeler*	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Echinochloa colona</i> (L.) Link*	EM/AM	+	+	+	+	+	+	+	+	+	0.41
<i>Echinochloa crus-gavonis</i> (Kunth) Schult.*	AM	-	+	+	-	+	+	-	-	-	0.18
<i>Echinochloa polystachya</i> (Kunth) Hitchc.*	EM	+	+	+	+	+	+	+	+	-	0.37
<i>Eragrostis pilosa</i> (L.) P. Beauv.*	AM	-	+	+	-	+	+	+	+	+	0.32
<i>Eriochloa punctata</i> (L.) Desv. ex Ham.*	AM	+	+	+	+	+	+	-	+	+	0.37
<i>Heleochloa schoenoides</i> (L.) Host	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Homolepis isocalycia</i> (G. Mey.) Chase	EM	+	+	+	-	-	+	-	-	+	0.23
<i>Hymenachne amplexicaulis</i> (Rudge) Nees*	EM/AM	-	+	+	+	+	+	+	+	-	0.32
<i>Leersia hexandra</i> Sw.*	EM	-	+	-	-	-	+	-	-	-	0.09
<i>Leptochloa fascicularis</i> (Lam.) A. Gray	EM	+	+	+	-	+	+	+	+	-	0.32
<i>Luziola brasiliiana</i> Moric.	EM/AM	-	+	+	-	+	+	+	-	-	0.23
<i>Oryza sativa</i> L.*	AM	+	-	-	+	+	+	-	-	-	0.18
<i>Panicum aquaticum</i> Poir.*	EM	-	+	+	+	+	+	-	-	-	0.23
<i>Panicum boliviense</i> Hack.	EM/AM	-	+	-	-	+	-	-	-	-	0.09
<i>Panicum dichotomiflorum</i> Michx.*	EM	-	+	+	+	+	+	-	+	-	0.27
<i>Panicum maximum</i> Jacq.*	EM/AM	+	+	+	+	+	+	-	+	+	0.37
<i>Panicum milleflorum</i> Hitchc. and Chase	AM	-	-	+	+	-	+	-	-	-	0.14
<i>Panicum parvifolium</i> Lam.	EM	+	+	-	+	+	+	+	-	+	0.32
<i>Paratheria prostrata</i> Griseb.	EM	-	-	-	-	-	+	-	-	-	0.05
<i>Paspalidium geminatum</i> (Forssk.) Stapf	EM/AM	-	+	+	+	+	+	+	+	-	0.32
<i>Paspalum millegrana</i> Schrad.	EM	+	+	+	+	+	+	+	+	+	0.41
<i>Paspalum repens</i> P.J. Bergius*	EM	+	+	+	-	+	+	-	-	-	0.23
<i>Paspalum vaginatum</i> Sw.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Setaria parviflora</i> (Poir.) Kerguélen.	EM	+	+	+	-	+	+	-	+	+	0.32
<i>Sporobolus indicus</i> (L.) R.BR.*	EM	-	+	-	+	+	+	-	+	+	0.27
<b>Podostemaceae</b>											
<i>Apinagia richardiana</i> (Tul.) P. Royen	RS	+	-	+	-	-	+	-	-	-	0.14
<i>Mourera fluviatilis</i> Aubl.	RS	+	-	-	+	-	+	-	-	-	0.14
<i>Tristicha trifaria</i> (Bory ex Willd.) Spreng.	RS	-	-	+	-	-	+	-	-	-	0.09
<b>Polygonaceae</b>											
<i>Coccoloba confusa</i> R.A. Howard	EM/AM	+	+	+	-	-	+	-	-	+	0.23
<i>Coccoloba ochreolata</i> Wedd.	EM/AM	+	+	-	-	-	+	+	-	+	0.23
<i>Polygonum acre</i> Lam.	AM	+	+	-	-	-	+	-	-	-	0.14
<i>Polygonum ferrugineum</i> Wedd.	EM/AM	+	+	+	+	+	+	-	-	-	0.27
<i>Polygonum hispidum</i> Kunth	EM/AM	-	+	+	+	+	+	+	+	+	0.37
<i>Polygonum hydroppiperoides</i> Michx.*	EM/RF	-	+	+	-	+	+	+	-	+	0.27
<i>Polygonum punctatum</i> Elliott	AM	+	+	+	-	+	+	+	-	+	0.32
<i>Ruprechtia laxiflora</i> Meisn.	AM	+	+	+	-	+	+	-	-	+	0.27
<i>Triplaris gardneriana</i> Wedd.	AM	+	+	+	+	+	+	+	+	+	0.41
<b>Pontederiaceae</b>											
<i>Eichhornia azurea</i> (Sw.) Kunth*	FF	-	+	-	-	-	+	-	-	-	0.09
<i>Eichhornia crassipes</i> (Mart.) Solms*	EM/FF	+	+	+	+	+	+	-	+	-	0.32
<i>Eichhornia diversifolia</i> (Vahl) Urb.	RF	-	+	+	+	+	+	-	-	-	0.23
<i>Eichhornia heterosperma</i> Alexander	AM	-	+	+	-	+	+	+	+	-	0.27
<i>Eichhornia paniculata</i> (Spreng.) Solms*	EM/AM	+	+	+	-	+	+	-	+	+	0.32
<i>Heteranthera limosa</i> (Sw.) Willd.*	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Heteranthera oblongifolia</i> C. Mart. ex Roem. and Schult.	AM	+	+	+	+	+	+	+	+	+	0.41
<i>Heteranthera reniformis</i> Ruiz and Pav.*	RF	+	+	+	-	-	+	-	-	-	0.18

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<i>Heteranthera seubertiana</i> Solms	AM	-	-	+	-	-	+	-	+	-	0.14
<i>Hydrothrix gardneri</i> Hooker F.	RS	-	+	+	-	+	+	+	+	-	0.27
<i>Pontederia cordata</i> L. *	EM	+	+	+	+	+	+	+	-	-	0.32
<b>Portulacaceae</b>											
<i>Portulaca marginata</i> Kunth *	AM	-	+	-	-	-	-	-	-	-	0.05
<b>Potamogetonaceae</b>											
<i>Potamogeton polygonus</i> Cham. and Schltldl.	RS	-	-	-	-	-	+	-	-	-	0.05
<b>Pteridaceae</b>											
<i>Acrostichum aureum</i> L.	AM	-	+	+	+	+	+	-	+	+	0.32
<i>Acrostichum danaeifolium</i> Langsd. and Fisch.	AM	-	+	-	+	-	+	-	-	-	0.14
<i>Ceratopteris pteridoides</i> (Hook.) Hieron.	EM/AM	-	+	+	+	-	+	+	-	-	0.23
<i>Ceratopteris thalictroides</i> (L.) Brongn.	EM/AM	-	+	-	-	-	+	-	-	-	0.09
<i>Pityrogramma calomelanos</i> (L.) Link	EM/AM	-	+	+	+	-	+	-	-	-	0.18
<i>Pteris vittata</i> L. *	AM	-	-	-	-	-	+	-	-	-	0.05
<b>Ricciaceae</b>											
<i>Ricciocarpos natans</i> (L.) Corda	RF	+	+	-	-	-	-	-	-	-	0.09
<b>Rubiaceae</b>											
<i>Borreria alata</i> (Aubl.) DC.	AM	-	-	-	-	-	+	-	+	-	0.09
<i>Borreria scabiosoides</i> Cham. and Schltldl.	AM	-	+	+	-	+	+	+	-	-	0.23
<i>Borreria verticillata</i> (L.) G. Mey. *	AM	-	+	+	-	-	-	-	-	-	0.09
<i>Coutarea hexandra</i> (Jacq.) K. Schum.	AM	-	-	+	+	-	+	+	-	-	0.18
<i>Machaonia spinosa</i> Cham. and Schltldl.	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Mapouria corymbifera</i> Müll. Arg.	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Pentodon pentandrus</i> (Schumach. and Thonn.) Vatke	AM	-	+	-	-	-	+	-	-	-	0.09
<i>Psychotria deflexa</i> DC.	AM	+	+	+	+	-	+	-	+	-	0.27
<i>Psychotria erecta</i> (Aubl.) Standl. and Steyerf.	AM	-	+	+	-	+	+	-	-	+	0.23
<b>Ruppiaceae</b>											
<i>Ruppia maritima</i> L.	RS	-	-	-	-	+	+	-	+	-	0.14
<b>Salviniaceae</b>											
<i>Azolla caroliniana</i> Willd. *	FF	-	+	-	-	-	+	-	-	-	0.09
<i>Azolla filiculoides</i> Lam.	FF	-	+	-	-	-	+	-	-	-	0.09
<i>Azolla microphylla</i> Kaulf.	FF	-	-	-	-	-	+	-	-	-	0.05
<i>Salvinia auriculata</i> Aubl. *	FF	+	+	+	+	+	+	-	+	+	0.37
<i>Salvinia biloba</i> Raddi	FF	-	-	-	-	-	+	-	-	-	0.05
<i>Salvinia martynii</i> Kopp.	FF	-	+	-	-	-	-	-	-	-	0.05
<i>Salvinia minima</i> Baker	FF	-	+	-	-	-	+	-	-	-	0.09
<i>Salvinia oblongifolia</i> Martius	FF	-	+	-	-	-	+	-	-	-	0.09
<b>Solanaceae</b>											
<i>Brunfelsia uniflora</i> (Pohl) D. Don	AM	+	+	+	-	-	+	+	-	-	0.23
<i>Nicotiana glauca</i> Graham	AM	+	+	+	-	+	+	-	+	-	0.27
<i>Physalis pubescens</i> L. *	AM	-	+	+	-	-	-	-	-	+	0.14
<i>Solanum asperum</i> Rich. *	AM	+	+	+	+	+	+	+	-	-	0.32
<i>Solanum baturitense</i> Huber	AM	+	+	+	+	+	+	-	-	-	0.27
<i>Solanum capsicoides</i> All. *	AM	-	+	+	-	+	+	-	-	+	0.23
<i>Solanum paludosum</i> Moric.	AM	+	+	+	+	+	+	-	-	+	0.32
<i>Solanum paniculatum</i> L. *	AM	-	+	+	+	-	+	-	+	+	0.27
<i>Solanum stipulaceum</i> Roem. and Schult. *	AM	+	+	+	-	+	+	+	-	+	0.32
<b>Sphenocleaceae</b>											
<i>Sphenoclea zeylanica</i> Gaertn. *	AM	-	-	+	+	-	+	-	-	-	0.14
<b>Thelypteridaceae</b>											
<i>Thelypteris interrupta</i> (Willd.) K. Iwats.	EM/AM	+	+	-	+	-	+	-	-	-	0.18
<b>Typhaceae</b>											
<i>Typha domingensis</i> Pers. *	EM	-	+	-	-	-	+	-	-	-	0.09
<b>Urticaceae</b>											
<i>Cecropia adenopus</i> Mart. ex Miq. *	AM	-	+	-	-	-	+	-	-	-	0.09
<b>Verbenaceae</b>											
<i>Lantana camara</i> L. *	AM	+	-	+	+	+	+	-	-	+	0.27
<i>Priva bahiensis</i> A. DC. *	AM	+	+	-	-	+	+	-	-	+	0.23
<i>Priva lappulacea</i> (L.) Pers.	AM	-	-	-	-	-	+	-	-	-	0.05
<i>Stachytarpheta cayennensis</i> (Rich.) Vahl *	AM	-	+	-	+	-	+	-	-	-	0.14

TABLE 2. CONTINUED.

FAMILY/SPECIES	LF	AL	BA	CE	MA	PB	PE	PI	RN	SE	PO %
<i>Stachytarpheta elatior</i> Schrad. ex Schult. *	AM	-	+	-	-	-	+	-	-	-	0.09
<b>Xyridaceae</b>											
<i>Xyris anceps</i> Lam.	EM	+	-	+	-	+	+	-	-	-	0.18
<i>Xyris capensis</i> Thunb.	EM	-	-	-	-	-	+	-	-	-	0.05
<i>Xyris fallax</i> Malme	EM	-	+	-	-	-	+	-	-	-	0.09
<i>Xyris jupicai</i> Rich.	EM	-	+	-	+	+	+	-	-	+	0.23
<i>Xyris laxifolia</i> Mart.	AM	-	+	+	-	-	-	-	-	-	0.09
<b>Zingiberaceae</b>											
<i>Hedychium coronarium</i> J. König *	EM	+	+	-	-	+	+	-	-	-	0.18

## LITERATURE CITED

- Agostinho, A.A., S.M. Thomaz, and L.C. Gomes. 2005. Conservation of the Biodiversity of Brazil's Inland Waters. *Conservation Biology* 19 (3): 646-652.
- Amaral, M.C.E., V. Bittrich, L.O. Anderson and L.Y. Aona. 2008. *Guia de Campo de Plantas Aquáticas e Palustres do Estado de São Paulo*. Ribeirão Preto: Holos. 452 p.
- ANA 2012. *Agência Nacional das Águas*. Electronic Database accessible at <http://www2.ana.gov.br>. Captured on 10 January 2012.
- APG III 2009. Angiosperm Phylogeny Group. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Botanical Journal of the Linnean Society* 161: 105-121.
- Bicudo, C.E.M., J.G. Tundisi, and M.C.B. Scheuenstuhl. 2010. *Águas do Brasil: análises estratégicas*. São Paulo: Instituto de Botânica. 224 p.
- Bonilla-Barbosa, J.R. and R.A. Novelo. 1995. *Manual de identificación de plantas acuáticas del Parque Nacional Lagunas de Zempoala, México*. Ciudad de Mexico: Instituto de Biología, Universidad Nacional Autónoma de México. 186 p.
- Bove, C.P., A.S.B. Gil, C.B. Moreira and R.F.B. Anjos. 2003. Hidrófitas fanerogâmicas de ecossistemas aquáticos temporários da planície costeira do Estado do Rio de Janeiro, Brasil. *Acta Botanica Brasilica* 17(1):119-135.
- Buck, W.R. and B. Goffinet. 2000. Morphology and classification of mosses; p. 72-124. In J.A. Shaw and B. Goffinet (ed.). *Bryophyte Biology*. Cambridge: Cambridge University Press.
- Campelo, M.J.A., J.A. Siqueira-Filho, V.M. Cotarelli, E.B. Souza, W.A. Pimenta and J.V. Pott, 2012. Macrófitas Aquáticas nas Áreas do Projeto da Integração do Rio São Francisco; p. 192-229. In J.A. Siqueira-Filho, E.M.C. Leme. (ed.). *Flora das Caatingas do Rio São Francisco*. Rio de Janeiro: Andrea Jakobsson Estúdio Editorial.
- Chambers, P.A., P. Lacombe, K.J. Murphy and S.M. Thomaz. 2008. Global diversity of aquatic macrophytes in freshwater. *Hydrobiologia* 595: 9-26.
- Clarke, K.R. and R.N. Gorley. 2006. *PRIMER - Plymouth Routines in Multivariate Ecological Research*. Ver. 6: User Manual/Tutorial. PRIMER-E, Plymouth.
- Cook, C.D.K. 1996. *Aquatic plant book*. Amsterdam: SPB Academic Publishing. 228 p.
- Cirilo, J.A., S.M.G.L. Montenegro and J.N.B. Campos. 2010. Questão da água no semiárido brasileiro; p. 81-91. In Bicudo, C.E.M., J.G. Tundisi and M.C.B. Scheuenstuhl (org). *Águas do Brasil: análises estratégicas*. São Paulo: Instituto de Botânica.
- Esteves, F.A. 2011. *Fundamentos de limnologia*. 3ed. Rio de Janeiro: Interciência. 826 p.
- Ferreira, F.A., R.P. Mormul, S.M. Thomaz, A. Pott, and V.J. Pott, 2011. Macrophytes in the upper Paraná river floodplain: checklist and comparison with other large South American wetlands. *Revista de Biologia Tropical* 59 (2): 541-556.
- Ferreira, F.A., R.P. Mormul, G. Pedralli, V.J. Pott and A. Pott, 2010. Estrutura da comunidade de macrófitas aquáticas em três lagoas do Parque Estadual do Rio Doce, Minas Gerais, Brasil. *Hoehnea* 37(1): 43-52.
- França, F., E. Melo, I.B. Oliveira, A.T.C.C. Reis, G.L. Alves and M.F. Costa. 2010. Plantas vasculares das áreas alagadas dos Marimbus Chapada Diamantina, BA, Brasil. *Hoehnea* 37(4): 719-730.
- França, F., E. Melo, A.G. Neto, D. Araújo, M. Bezerra, H.M. Ramos, I. Castro and D. Gomes. 2003. Flora vascular de açudes de uma região do semiárido da Bahia, Brasil. *Acta Botanica Brasilica* 17(4): 549-559.
- Goetghebeur, P. 1998. Cyperaceae; p. 141-190. In K. Kubitzki (ed.). *The families and genera of vascular plants*. Berlin: Springer.
- Gopal, B. and W.J. Junk. 2000. Biodiversity in wetlands: in introduction; p. 1-10. In B. Gopal, W.J. Junk and J.A. Davis (ed.). *Biodiversity in wetlands: assessment, function and conservation*. Leiden: Backhuys Publishers.
- Heiden, G., J.R.V. Iganci, V.L. Bobrowski and L. Macias. 2007. Biogeografia de Baccharis sect. Caulopterae (Asteraceae) no Rio Grande do Sul, Brasil. *Rodriguésia* 58 (4): 787-796.
- Henry-Silva, G.G., R.S.T. Moura and L.L.O. Dantas, 2010. Richness and distribution of aquatic macrophytes in Brazilian semi-arid aquatic ecosystems. *Acta Limnológica Brasiliensis* 22(2): 147-156.
- IBGE 2002. Instituto Brasileiro de Geografia e Estatística. Electronic Database accessible at <http://www.ibge.gov.br>. Captured on 10 January 2012.
- Irgang, B.E. and C.V.S. Gastal-Jr. 1996. *Macrófitas Aquáticas da Planície Costeira do RS*. Porto Alegre: Editora da UFRGS. 290 p.
- Irgang, B.E., G. Pedralli, and J.I. Waechter 1984. Macrófitas aquáticas da Estação Ecológica do Taim, Rio Grande do Sul, Brasil. *Roessléria* 6: 395-404.
- Junk, W.J. and M. Piedade 1993. Herbaceous plants of the Amazon floodplain near Manaus: Species diversity and adaptation to the fluid pulse. *Amazoniana* 12(4): 467-484.
- Lahitte, H.B. and J.A. Hurrell. 1996. *Plantas hidrófilas de la Isla Martín García*. Buenos Aires: Ministerio de La Producción Provincia de Buenos Aires. 200 p.
- Lembi, C.A. 2009. *Aquatic plant Management: Identifying and Managing Aquatic Vegetation*. Indiana: Purdue extension. 19 p.
- Lima, L.F., P.B. Lima, R.C. Soares-Júnior, R.M.M. Pimentel and C.S. Zickel. 2009. Diversidade de macrófitas aquáticas no estado de Pernambuco: levantamento em herbário. *Revista de Geografia*. 26(3): 307-319.
- Lorenzi, H. 2008. *Plantas daninhas do Brasil: terrestres, aquáticas, parasitas e tóxicas*. Nova Odessa: Instituto Plantarum. 261 p.
- Kita, K.K. and M.C. Souza 2003. Levantamento florístico e fitofisionomia da lagoa Figueira e seu entorno, planície alagável do alto rio Paraná, Porto Rico, Estado do Paraná, Brasil. *Acta Scientiarum. Biological Sciences* 25(1):145-155.
- Magurran, A.E. 2004. *Measuring biological diversity*. Malden: Blackwell Publishing. 256 p.
- Margurran, A.E. 1988. *Ecological diversity and its measurement*. Princeton: Princeton University. 179 p.
- Matias, L.G. 2010. A synopsis of Alismataceae from the semi-arid region of Northeastern Brazil. *Revista Caatinga* 23(4): 46-53.
- Matias, L.Q., E.R. Amado and E.P. Nunes. 2003. Macrófitas aquáticas da lagoa de Jijoca de Jericoacoara, Ceará, Brasil. *Acta Botanica Brasilica* 4(17): 623-631.
- Matias, L.Q. and E.P. Nunes. 2001. Levantamento florístico da área de proteção ambiental de Jericoacoara, Ceará. *Acta Botânica Brasilica* 15(1): 35-43.
- Mereles, F., R. Degen, N.L. Kochalca. 1992. Humedales en el Paraguai: Breve reseña de su vegetación. *Amazoniana* 12(2): 305-316.
- Miretzky, P., A. Saralegui and C.A. Fernandez. 2004. Aquatic macrophytes potential for the simultaneous removal of heavy metals (Buenos Aires, Argentina). *Chemosphere* 57: 997-1005.
- MOBOT 2011. *Missouri Botanical Garden*. Electronic Database accessible at <http://www.mobot.org>. Captured on 30 October 2011.
- Mormul, R.P., F.A. Ferreira, P. Carvalho, T.S. Michelan, M.J. Silveira and S.M. Thomaz 2010. Aquatic macrophytes in the large, sub-tropical Itaipu Reservoir. *Revista de Biologia Tropical* 58: 1437-1452.
- Moura, M.A.M., D.A.S. Franco and M.B. Matallo. 2009. Manejo integrado de macrófitas aquáticas. *Biológico* 71(1): 77-82.
- Moura-Júnior, E.G., M.C. Abreu, W. Severi and G.A.S.T. Lira. 2011. Are floristic composition, richness and life forms of aquatic macrophytes affected by the dam-river gradient of the Sobradinho Reservoir? *Rodriguésia* 62: 731-742.
- Moura-Júnior, E.G., M.C. Abreu, W. Severi and G.A.S.T. Lira. 2010. Macroflora aquática do Reservatório Sobradinho – BA, trecho sub-médio do Rio São Francisco; p. 189-212. In Moura, A.M., E.L. Araújo, M.C. Bittencourt-Oliveira, R.M.M. Pimentel and U.P. Albuquerque (eds.). *Reservatórios do Nordeste do Brasil: biodiversidade, ecologia e*

- manejo*. Recife: Nuppea.
- Moura-Júnior, E.G., S.S.L. Silva, L.F. Lima, P.B. Lima, E.B. Almeida-Jr., L.M. Pessoa, F.S. Santos-Filho, D.P.W. Medeiros, R.M.M. Pimentel and C.S. Zickel. 2009. Diversidade de plantas aquáticas vasculares em açudes do Parque Estadual de Dois Irmãos (PEDI), Recife-PE. *Revista de Geografia* 26: 178-293.
- Murphy, K.J., G. Dickinson, S.M. Thomaz, L.M. Bini, K. Dick, K. Greaves, M.P. Kennedy, S. Livingstone, H. McFerran, J.M. Milne, J. Oldroyd, and R.A. Wingfield. 2003. Aquatic plant communities and predictors of diversity in a sub-tropical river floodplain: the upper Rio Paraná, Brazil. *Aquatic Botany* 77: 257-276.
- Nascimento, P.R.F. 2009. *Levantamento florístico e produtividade de macrófitas aquáticas ocorrentes em ambientes limnéticos do estado de Pernambuco – Brasil*. Tese (Doutorado em Botânica) – Universidade Federal Rural de Pernambuco, Recife - PE.
- Neiff, J.J. 1982. Esquema sucessional de La vegetación en las islas flotantes del Chaco Argentino. *Boletim da Sociedade Argentina de Botânica* 21(1-4): 325-341.
- Neiff, J.J., P.A.S.G. Neiff, C.A.E. Patiño and B.I. Chiozzi. 2000. Prediction of colonization by macrophytes in the Yaciretá reservoir of the Paraná river (Argentina and Paraguay). *Revista Brasileira de Biologia* 60(4): 615-626.
- Neves, E.L., K.R.B. Leite, F. França and E. Melo. 2006. Plantas aquáticas vasculares em uma lagoa de planície costeira no município de Candeias, Bahia, Brasil. *Sitientibus, Série Ciências Biológicas* 6(1): 24-29.
- Obot, E.A. and I.G. Mbagwu. 2008. Successional patterns of aquatic macrophytes in Jebba Lake, Nigeria. *African Journal of Ecology* 26(4): 295-299.
- Pedro, F., L. Maltchik and I. Bianchini Jr, 2006. Hydrologic cycle and dynamics of aquatic macrophytes in two intermittent rivers of the semi-arid region of Brazil. *Brazilian Journal of Biology* 2: 575-585.
- Pompêo, M.L.M. and V. Moschini-Carlos, 2003. *Macrófitas aquáticas e perifiton, aspectos ecológicos e metodológicos*. São Carlos: RIMA. 130 p.
- Pott, V.J. Pott, A. Lima, L.C.P. Moreira, S.N. and A.K.M. Oliveira. 2011. Aquatic macrophyte diversity of the Pantanal wetland and upper basin. *Brazilian Journal of Biology* 1: 255-263.
- Pott, V.J. and A. Pott. 2000. *Plantas aquáticas do Pantanal*. Brasília: Embrapa. 414 p.
- Ramos, V.L. and R.A. Novelo, 1993. Vegetación y flora acuática de la laguna de Yuriria, Guanajuato, México. *Acta Botánica* 25: 61-79.
- Raynal-Roques, A. 1980. Les plantes aquatiques; p. 63-152. In J.R. Durand and C. Lévêque (ed.). *Flore et faune Aquatiques de L'Afrique Sahelosoudanienne*. Paris: Orstom.
- Ribeiro, J.P.N., L.K. Takao, R.S. Matsumoto, C. Urbanetz and M.I.S. Lima. 2011. Plantae, aquatic, amphibian and marginal species, Massaguaçu River Estuary, Caraguatatuba, São Paulo, Brazil. *Check List* 7(2): 133-138.
- Scremin-Dias, E., V.J. Pott, R.C. Hora and P.R. Souza. 1999. *Nos jardins submersos da Bodoquena: guia para identificação de plantas aquáticas de Bonito e região*. Campo Grande: Editora da UFMS. 160 p.
- Sculthorpe, C.D. 1967. *The biology of aquatic vascular plants*. London: Edward Arnold Ltd. 610p.
- Silva, S.S.L. and C.S. Zickel 2010. Macrófitas aquáticas: conceitos e metodologia para os reservatórios nordestinos; p 71-186. In Moura, A.M., E.L. Araújo, M.C. Bittencourt-Oliveira, R.M.M. Pimentel and U.P. Albuquerque, (eds.). *Reservatórios do Nordeste do Brasil: biodiversidade, ecologia e manejo*. Recife: Nuppea.
- Silva, V.P.R., M.J.F. Guedes, W.F.A. Lima and J.H.B.C. Campos. 2002. Modelo de previsão de rendimento de culturas de sequeiro, no semi-árido do Nordeste do Brasil. *Revista Brasileira de Engenharia Agrícola e Ambiental* 6: 83-87.
- Smith, A.R., K.M. Pryer, E. Schuettpelz, P. Korall, H. Schneider and P.G. Wolf. 2006. A classification for extant ferns. *Taxon* 55(3): 705-731.
- Sobral-Leite, M., M.J.A. Campelo, J.A. Siqueira-Filho and S.I. Silva. 2010. Checklist das macrófitas vasculares de Pernambuco: riqueza de espécies, formas biológicas e considerações sobre distribuição; p. 253-280. In U.P. Albuquerque, A.N. Moura and E.L. Araujo (ed.). *Biodiversidade, potencial econômico e processos ecofisiológicos em ecossistemas nordestinos*. Volume II. Recife: Nuppea.
- Sousa, W.T.Z., S.M. Thomaz, K.J. Murphy, M.J. Silveira and R.P. Mormul. 2009. Environmental predictors of the occurrence of exotic *Hydrilla verticillata* (L.f.) Royle and native *Egeria najas* Planch. in a sub-tropical river floodplain: the Upper River Paraná, Brazil. *Hydrobiologia* 632: 65-78.
- Souza, V.C. and H. Lorenzi. 2008. *Botânica Sistemática*. 2 ed. Nova Odessa: Instituto Plantarum. 544 p.
- SPLINK 2011. *System that integrates real-time primary data from scientific collections of Brazil*. Electronic Database accessible at <http://www.splink.cria.org.br> Captured on 30 march 2011.
- Subramanyam, K. 1962. *Aquatic Angiosperms. A systematic account of common Indian aquatic Angiosperms*. New Delhi: C.S.I.R. 190 p.
- Thomaz, S.M. and E.R. Cunha. 2010. The role of macrophytes in habitat structuring in aquatic ecosystems: methods of measurement, causes and consequences on animal assemblages composition and biodiversity. *Acta Limnologica Brasiliensia* 22(2): 218-236.
- Thomaz, S.M. and L.M. Bini. 2003. Análise crítica dos estudos sobre macrófitas aquáticas desenvolvidos no Brasil; p. 19-38. In S.M. Thomaz and L.M. Bini, (ed.). *Ecologia e manejo de macrófitas aquáticas*. Maringá: EDUEM.
- Thomaz, S.M., M.C. Souza and L.M. Bini. 2003. Species richness and beta diversity of aquatic macrophytes in a large subtropical reservoir (Itaipu Reservoir, Brazil): the influence of limnology and morphometry. *Hydrobiologia* 505:119-128.
- Tundisi, J.G. and Tundisi, T.M. 2008. *Limnologia*. São Paulo: Oficina de Texto. 632 p.
- Tur, N.M. 1972. Um caso de epifitismo acuático. *Boletim de la Sociedad Argentina de Botânica* 10(4): 323-327.
- Vilarrubia, T.V. and M. Cora. 1993. Estudio sobre la distribución y ecología de macrofitos acuáticos en el embalse de Guri. *Interiencia* 18: 77-82.
- Wood, R.D. and K. Imahori. 1964. *A revision of the Characeae. 2: iconograph of the Characeae*. Weinheim: J. Cramer. 797 p.

RECEIVED: May 2012

ACCEPTED: March 2013

PUBLISHED ONLINE: April 2013

EDITORIAL RESPONSIBILITY: Frederico Augusto Guimarães Guilherme

# Aquatic macrophytes of Northeastern Brazil: Checklist, richness, distribution and life forms

Edson Gomes de Moura-Júnior<sup>1</sup>, Liliane Ferreira Lima<sup>1</sup>, Simone Santos Lira Silva<sup>2</sup>, Raíssa Maria Sampaio de Paiva<sup>3</sup>, Fernando Alves Ferreira<sup>4</sup>, Carmen Silvia Zickel<sup>5\*</sup> and Arnildo Pott<sup>4</sup>

- 1 Graduate student (PhD) in Plant Biology, Universidade Federal de Minas Gerais, Biology Department. Av. Antônio Carlos, 6627. CEP 31270-901. Belo Horizonte, MG, Brazil.
  - 2 Graduate student (PhD) in Botany, Universidade Federal Rural de Pernambuco, Biology Department. Av. Dom Manoel de Medeiros, s/n°, Dois Irmãos. CEP 52171-900. Recife, PE, Brazil.
  - 3 Graduate student (MSc) in Natural Resource, Universidade Federal de Roraima, Biology Department. Av. Capitão Ene Garcez, 2413, Aeroporto, Boa Vista. CEP 69304-000. Roraima, RR, Brazil.
  - 4 Universidade Federal do Mato Grosso do Sul, Program in Plant Biology, Center for Biological Sciences and Health, Biology Department. Cidade Universitária, s/n - CEP 79070-900. Campo Grande, MS, Brazil.
  - 5 Universidade Federal Rural de Pernambuco, Program in Botany, Biology Department. Av. Dom Manoel de Medeiros, s/n°, Dois Irmãos. CEP 52171-900. Recife, PE, Brazil.
- \* Corresponding Author. E-mail: [zickelbr@yahoo.com](mailto:zickelbr@yahoo.com)

## ERRATUM

**AUTHORS AFFILIATION:** Liliane Ferreira Lima<sup>1</sup>, Simone Santos Lira Silva<sup>2</sup> **should be** Liliane Ferreira Lima<sup>2</sup>, Simone Santos Lira Silva<sup>5</sup>

We regret these errors.

May 2013

**ABSTRACT:** Species richness in Pernambuco **should be** 369, not 370.

**PAGE 310** within **TABLE 2:**

### Salviniaceae

<i>Salvinia biloba</i> Raddi	FF	-	-	-	-	-	+	-	-	-	0.05
<i>Salvinia martynii</i> Kopp.	FF	-	+	-	-	-	-	-	-	-	0.05

**Should be**

### Salviniaceae

<i>Salvinia biloba</i> Raddi	FF	-	+	-	-	-	+	-	-	-	0.09
<i>Salvinia martynii</i> Kopp.	FF	-	-	-	+	-	-	-	-	-	0.05