

# *Tropidophorus robinsoni* Smith, 1919 (Squamata: Scincidae): New distribution record and map

Likhit Waiprom<sup>1,2</sup>, Yodchaiy Chuaynkern<sup>3</sup>, Prateep Duengkae<sup>2,4\*</sup>, Chantip Chuaynkern<sup>3</sup> and Kriangsak Sriboarod<sup>5</sup>

- 1 Thailand Natural History Museum, National Science Museum, Technopolis, Khlong 5, Khlong Luang, Pathum Thani, 12120 Thailand.
  - 2 Kasetsart University, Faculty of Forestry, Department of Forest Biology, Chatuchak, Bangkok, 10900 Thailand.
  - 3 Khon Kaen University, Faculty of Science, Department of Biology, Mueang, Khon Kaen, 40002 Thailand.
  - 4 Kasetsart University, Center for Advanced Studies in Tropical Natural Resources, Chatuchak, Bangkok, 10900 Thailand.
  - 5 Khlong Saeng Wildlife Research Station, P.O. Box 9, Makham, Surat Thani, 22150 Thailand.
- \* Corresponding author. E-mail: [prateep.du@ku.ac.th](mailto:prateep.du@ku.ac.th)

**ABSTRACT:** We present a new locality for *Tropidophorus robinsoni* Smith, 1919 based on a specimen collected from Khlong Saeng Wildlife Sanctuary, Surat Thani Province, southern Thailand, a new provincial record. The geographical distribution of the species is reviewed and an updated distribution map is presented.

The lygosomine genus *Tropidophorus* Duméril and Bibron, 1839 (type species: *T. cocincinensis*) currently contains 28 species (Nguyen *et al.* 2010b). Its morphology is characterized by exposure of the tympanum and presence of a single scale at the corner of the eyelid (*e.g.*, Greer 1970; Hikida *et al.* 2002; Greer and Biswas 2004; Honda *et al.* 2006). The members of *Tropidophorus* are known to occur in two regions: (1) Indochina and adjacent parts of continental Eurasia exclusive of Peninsular Malaysia, and (2) the South-east Asian islands of Borneo, Sulawesi, and the Philippines (Honda *et al.* 2006).

Within Indochina, 15 species are recognized (Nguyen *et al.* 2010a). Among them, the Thai *Tropidophorus* contain nine species (Chuaynkern *et al.* 2005): *T. bermorei* Blyth, 1853, *T. cocincinensis* Duméril and Bibron, 1839, *T. hangnam* Chuaynkern, Nabhitabhata, Inthara, Kamsook and Somsri, 2005, *T. laotus* Smith, 1923, *T. laticutatus* Hikida, Orlov, Nabhitabhata and Ota, 2002, *T. matsuii* Hikida, Orlov, Nabhitabhata and Ota, 2002, *T. microlepis* Günther, 1861, *Tropidophorus robinsoni* Smith, 1919, and *T. thai* Smith, 1919. The distribution of the *Tropidophorus* in Thailand is summarized in several works (see *e.g.*, Smith 1919, 1923; Taylor 1963; Nabhitabhata *et al.* 2004; Nabhitabhata and Chan-ard 2005; Das 2010). These works show that reports on localities of *Tropidophorus* species in southern Thailand are rare. The present work represents an additional record of *Tropidophorus* in southern Thailand which fills a gap in this skink's known geographical distribution.

The specimen was caught by hand on 28<sup>th</sup> April 2010 by Likhit Waiprom from a small stream whose bottom is covered with small igneous rocks. The site coordinates are 8° 59' 32.748"N, 98° 48' 18.1542"E. The stream flows in dry evergreen forest in Khlong Saeng Wildlife Sanctuary, Surat Thani Province, southern Thailand. The skink was preserved in 10% buffered formalin, and later transferred into 70% ethanol. It was catalogued as THNHM 17962 in the collections of the Thailand Natural History Museum (THNHM), Pathum Thani, Thailand. Measurements were made with a digital caliper to the nearest 0.1 mm.

Photographs were taken using a digital camera and modified by using Photoshop CS2 programs. The head drawing was made by L. Waiprom using ArcView program.

The specimen is an adult female (Figures 1 and 2) with a snout-vent length of 61.5 mm and a total length of 99 mm (tail regenerated). We assigned this specimen to *Tropidophorus robinsoni* based on its morphological characteristics that match the original description by Smith (1919) and other subsequent taxonomic works (see Taylor 1963; Chuaynkern *et al.* 2005; Nguyen *et al.* 2010a). This skink can be distinguished by the following characters: head scales keeled and rugose; dorsal scales strongly keeled; frontal not divided; frontonasal single; flank scales keeled and pointing straight backwards.


*Tropidophorus robinsoni* was previously known only from the type locality "Tasan, W. of Chumphon, P. Siam" (Smith, 1919), Khao Lak in Phang-nga Province, Thailand (Pauwels *et al.* 2000) and from Tanintharyi Division in Myanmar (Wogan *et al.* 2008). The new locality thus represents the first provincial record for Surat Thani Province (southern Thailand), the fourth locality for the species, and it partly fills approximately 200 km hiatus




**FIGURE 1.** Dorsal (A) and ventral (B) views of an adult female *Tropidophorus robinsoni* (THNHM 17962) from Khlong Saeng Wildlife Sanctuary, Surat Thani province, southern Thailand.

between Tasan and Khao Lak.

Current data seem to indicate that *Tropidophorus robinsoni* is restricted to peninsular Thailand and peninsular Myanmar (Figure 3), on both sides of the Isthmus of Kra. Further studies are recommended for better defining the distribution range of this rare species.



**FIGURE 2.** Dorsal (A), lateral (B), and ventral (C) head views of an adult female *Tropidophorus robinsoni* (THNHM 17962) from Khlong Saeng Wildlife Sanctuary, Surat Thani Province, southern Thailand. Scale bar indicates 10 mm.



**FIGURE 3.** Updated distribution map of *Tropidophorus robinsoni*: (★), type locality "Tasan, Chumphon Province" (=Tasan, Chumphon Province, Thailand); (■), Khao Lak, Phang-nga Province (Thailand); (●), Khlong Saeng Wildlife Sanctuary (Surat Thani Province, Thailand); and (▲), Tanintharyi Division (Myanmar).

**ACKNOWLEDGMENTS:** We thank Chorphaka Vijittrakoolchai, Sutasinee Saosong, Jaruwan Somphoa, Watchara Chitchamnong and Auaiphon Kangwal for assistance in the field. Atchara Teerawattananon is thanked for her kindness to edit our manuscript. The following institutions are thanked for supporting this work: the National Science Museum (Thailand Natural History Museum, Pathum Thani), the Khlong Saeng Wildlife Research Station (Surat Thani), Kasetsart University (Bangkok) and Khon Kaen University (Khon Kaen).

**LITERATURE CITED**

Chuaynkern, Y., J. Nabhitabhata, C. Inthara, M. Kamsook and K. Somsri. 2005. A new species of the Water Skink *Tropidophorus* (Reptilia: Squamata: Scincidae) from northeastern Thailand. *The Thailand Natural History Museum Journal* 1:165-175.

Das, I. 2010. *A Field Guide to the Reptiles of Thailand and South-East Asia*. Bangkok: Asia Books Co. Ltd. 376 p.

Greer, A.E. 1970. A subfamilial classification of scincid lizards. *Bulletin of the Museum of Comparative Zoology* 139: 151-184.

Greer, A.E. and S. Biswas. 2004. A generic diagnosis for the Southeast Asian scincid lizard *Tropidophorus* Duméril and Bibron, 1839 with some additional comments on its morphology and distribution. *Journal of Herpetology* 38: 426-430.

Hikida, T., N.L. Orlov, J. Nabhitabhata and H. Ota. 2002. Three new depressed-bodied water skinks of the genus *Tropidophorus* (Lacertilia: Scincidae) from Thailand and Vietnam. *Current Herpetology* 21: 9-23.

Honda, M., H. Ota, R.W. Murphy and T. Hikida. 2006. Phylogeny and biogeography of water skinks of the genus *Tropidophorus* (Reptilia: Scincidae): a molecular approach. *Zoologica Scripta* 35: 85-95.

Nabhitabhata J., T. Chan-ard and Y. Chuaynkern. 2004. *Checklist of Amphibians and Reptiles in Thailand*. Bangkok: Office of Environmental Policy and Planning. 152 p.

Nabhitabhata, J. and T. Chan-ard. 2005. *Thailand Red Data: Mammals, Reptiles and Amphibians*. Bangkok: Office of Natural Resources and Environmental Policy and Planning. 158 p.

Nguyen, Q.T., V.S. Nguyen, N. Orlov, N.T. Hoang, W. Böhme and T. Ziegler. 2010a. A review of the genus *Tropidophorus* (Squamata, Scincidae) from Vietnam with new species records and additional data on natural history. *Journal of Zoological Systematics and Evolutionary Research* 86: 5-19.

Nguyen, Q.T., T.T. Nguyen, A. Schmitz, N.L. Orlov and T. Ziegler. 2010b. A new species of the genus *Tropidophorus* Duméril & Bibron, 1839 (Squamata: Sauria: Scincidae) from Vietnam. *Zootaxa* 2439: 53-68.

Pauwels, O.S.G., O.-A. Laohawat, P. David, R. Bour, P. Dangsee, C. Puangjit and C. Chimsunchart. 2000. Herpetological investigations in Phang-Nga Province, southern Peninsular Thailand, with a list of reptile species and notes on their biology. *Dumerilia* 4: 123-154.

Smith, M.A. 1919. The lizards of the genus *Tropidophorus* in Siam, with description of two new species. *Journal of Natural History Society of Siam* 3: 223-228.

Smith, M.A. 1923. A review of the lizards of the genus *Tropidophorus* on the Asiatic mainland. *Proceedings of the Zoological Society of London* 1923: 775-781.

Taylor, E.H. 1963. The lizards of Thailand. *The University of Kansas Science Bulletin* 44: 687-1077.

Wogan, G.O.U., J.V. Vindum, J.A. Wilkinson, M.S. Koo, J.B. Slowinski, H. Win, T. Thin, S.W. Kyi, S.L. Oo, K.S. Lwin and A.K. Shein. 2008. New country records and range extensions for Myanmar amphibians and reptiles. *Hamadryad* 33: 83-96.

RECEIVED: November 2011

ACCEPTED: February 2013

PUBLISHED ONLINE: May 2013

EDITORIAL RESPONSIBILITY: Olivier S. G. Pauwels