

Checklist of Lizards of the Juruti, state of Pará, Brazil

Ana Lúcia Costa Prudente*, Fernanda Magalhães, Alessandro Menks and João Fabrício de Melo Sarmento

1 Museu Paraense Emílio Goeldi, Departamento de Zoologia, Av. Perimetral, 1901. CEP 66077-530. Belém, Pará (PA), Brazil.

* Corresponding author. E-mail: prudente@museu-goeldi.br

ABSTRACT: We present the first lizard species list for the municipality of Juruti, state of Pará, Brazil. The list was drawn up as a result of data obtained from specimens deposited in the Herpetological Collection of the Museu Paraense Emílio Goeldi and from inventories conducted in 2008-2011. Sampling methods included pitfall traps with drift fences and time constrained searches. We considered the data collected by other researchers, incidental encounters and records of dead individuals on the road. We recorded 33 species, 26 genera and ten families. *Norops tandai* was the most abundant species. Compared with the other regions of Amazonia, the region of Juruti presented a large number of lizards. However, further studies with an increase in the sampling effort, could prove this area to be richer in lizards than that observed so far.

INTRODUCTION

The Amazon rainforest is one of the world's biomes that exhibits high biological diversity and complexity of environments (Duellman 1978). Of the more than six million km² estimated to be the current total area of the Amazon, about 60% are in Brazilian territory (Capobianco *et al.* 2001), giving it the status of the largest and most diverse rainforest in the world (Silva *et al.* 2005).

The call for information on Amazonian fauna is undisputed among researchers; however, lack of knowledge is mainly related to low sampling (Vivo 1996; Silva Jr. 1998) and insufficient collection. For amphibians and reptiles, this scenario is no different. According to Ávila-Pires *et al.* (2010), due to the large extension of the Amazon region, the herpetofauna has not yet been identified completely, particularly for some groups of reptiles.

The richness of the lizard fauna of the Brazilian Amazon, although it is not fully known, is one of the highest amongst the Brazilian biomes (Rodrigues 2005). Of the 248 species of lizards present in Brazil (Bérnils and Costa 2012), 94 species are recorded in the Brazilian Amazon, which corresponds to 72% of all known species for the whole Amazon region (Ávila-Pires *et al.* 2007). This number grows each year due to studies and faunal inventories, which provide material for further and ongoing descriptions of Amazonian species (Ávila-Pires *et al.* 2007).

Among the main studies that list the composition of the Amazonian lizard fauna, the following can be highlighted: in the state of Amazonas, the studies of Martins (1991) in the region of Balbina (20 species), Duellman (1990) and Vitt *et al.* (2008) in the Manaus region (34 species), Ávila-Pires *et al.* (2009) in the Rio Ituxi (26 species), Ilha and Dixo (2010) in the municipality of Rio Preto da Eva (20 species), and Pantoja and Fraga (2012) in the Extractive Reserve of Rio Gregorio (15 species); in the state of Pará the work of Cunha *et al.* (1985) and Nascimento *et al.* (1987) in the Carajás area (25 species), Ávila-Pires and Hoogmoed (1997) and Bernardi *et al.* (2002) in the

Caxiuanã National Forest (28 species), Gardner *et al.* (2007) in the municipality of Monte Dourado (30 species), Ávila-Pires *et al.* (2009) in the Curuá-Una (22 species), Mendes-Pinto and Souza (2011) in the National Forest of Trairão (23 species), Silva *et al.* (2011) in the region of Barcarena (17 species) and Bernardo *et al.* (2012) in Tapirapé Biological Reserve (14 species); in the state of Rondônia, Macedo *et al.* (2008) in the municipality of Espigão do Oeste (29 species) and Ávila-Pires *et al.* (2009) in Guajará-Mirim State Park (23 species); and in the state of Acre, the studies of Ávila-Pires *et al.* (2009) in Porto Walter (29 species) and Bernarde *et al.* (2011) in the Extractive Reserve of Riozinho da Liberdade (29 species).

Located south of the Amazon River, state of Para, Brazil, the municipality of Juruti presents preserved tropical rainforest, with some areas destined for the exploration of bauxite. Herein, we present a list of lizard species for this region, based on fieldwork and collection data.

MATERIALS AND METHODS

Study area

The activities were developed in the Area under the Influence of the Project Juruti/Alcoa, in the west of the state of Pará, Brazil (02°09'09"S; 56°05'42"W) (Figure 1). The vegetation of the area is classified as dense ombrophilous forest, with llatsoil type Dystrophic Oxisol (MCT 2008). The climate is classified as AMw following Köppen's system (Kottek *et al.* 2006), which is characterized by a dry season of short duration, with annual rainfall between 2000 and 2250 mm/year. The mean annual temperature is 26°C and humidity is above 80% (MCT 2008).

Data Collection

To compose the list of lizard species in the region, we used data from expeditions conducted from 2008 to 2011 and from the material deposited in the herpetological collection of the Museu Paraense Emílio Goeldi (MPEG).

Eight expeditions were conducted, totaling 123 days, in the following years: 2008 (17 February to 3 March - 17 to 26 March, 1 to 21 December, in the rainy season), 2009 (19

to 28 January - 13 to 29 May, in the rainy season), 2010 (15 to 30 July - 15 to 30 September, in the dry season) and 2011 (15 to 30 January, in the rainy season).

Collections were conducted in six areas with different phytophysionomies and degrees of environmental disturbance (Figure 2), and located in the "Area under the Influence of the Juruti Project", municipality of Juruti, state of Pará, Brazil (Figure 1).

For descriptions of the areas were considered as characteristic of the vegetation circumference at breast height (CBH) and litter depth. For inferences of CBH trees were defined plots of 100m² (10 x 10), randomly chosen and set next to each trap, totaling six plots at each collection point (600 m²). All the individuals with circumference at breast height (CBH) ≥ 15cm were sampled. The depth of litter was measured within the same plots where the CBH were measured (five measurements per plot) using a millimeter ruler.

The six study areas are described below:

A) *Barroso* (02°28'18"S, 56°00'44"W): An area formed by dense ombrophilous to submontane forest (MCT 2008), which has streams (igarapés) with vegetation consisting of palm trees (*Socratea* spp., *Mauritia* sp., *Oenocarpus* sp. and *Astrocaryum* sp.), large trees reaching 35m in height (Santos *et al.* 2011), and CBH above 400cm (n= 92; mean of 57.5cm), with a leaf litter layer up to 14cm deep (n= 30; mean of 11.4cm) (Figure 2A). This area is barely impacted, with absence of artificial clearings, subsistence hunting activities or paved roads. These features make this area one of the most preserved among the areas studied. The collections in this area were performed only in the years 2010 and 2011.

B) *Capiranga* (02°28'57"S, 56°09'41"W): An area formed by dense ombrophilous to submontane forest (MCT 2008), with primary forest consisting of large trees

with height reaching 40m (Santos *et al.* 2011), and CBH above 390cm (n= 93; mean of 50.8cm), with a leaf litter layer up to 13cm deep (n= 30; mean of 10.8cm) (Figure 2B). Paved roads and ramifications are present in this area with moderate flow of vehicles. Although it is a preserved area, logging and population growth of the communities in its surroundings cause this area to be under great potential risk of degradation. The collections in this area were performed only in the 2008 to 2011.

C) *Adutora* (02°27'51"S, 56°10'59"W): An area formed by dense ombrophilous to submontane forest (MCT 2008), with secondary forest vegetation and the presence of igarapés and large artificial clearings formed by the extraction of timber. There are many palms and vines, maximum depth of leaf litter is 6.2cm (n= 30; mean of 5.8cm), and trees have a height of 35m (Santos *et al.* 2011), CBH above 92cm (n= 84; mean of 25cm) (Figure 2C). Timber extraction and a large flow of vehicles make this area the most impacted with large anthropic disturbance. The collections in this area were performed only in the years 2008 to 2011.

D) *Pacoval* (02°32'40"S; 56°10'02"W): An area formed by dense ombrophilous to submontane forest (MCT 2008). Presence of secondary forest (Figure 2E), with large numbers of trumpet trees (*Cecropia* spp.), leaf litter with maximum depth of 8cm (n= 30; mean of 59cm), trees with height of 35m (Santos *et al.* 2011) CBH above 102cm (n= 79; mean of 29cm). This area is crossed by a road that gives access to various private properties, where logging and subsistence hunting occur. The collections in this area were performed only in the years 2008 to 2010.

E) *Galiléia* (02°32'51"S; 56°13'32"W): An area formed by dense ombrophilous to submontane forest (MCT 2008), with secondary forest located on flat terrain. Trumpet trees (*Cecropia* spp.) are present, maximum depth of leaf litter is 8cm (n= 30; mean of 7.0cm) and trees have a CBH of 185cm (n= 82; mean of 27cm) (Figure 2D). This area shows great human disturbance caused by the large movement of vehicles and by logging. The collections in this area were performed only in the years 2008 to 2011.

F) *Mutum* (02°36'34"S; 56°11'46"W): An area formed by dense ombrophilous to submontane forest (MCT 2008). It features well-preserved primary forest, with the presence of streams (igarapés) and vegetation consisting of palm trees (*Socratea* sp.1, *Socratea* sp.2, *Mauritia* sp., *Oenocarpus* sp. and *Astrocaryum* sp.), large trees above 30m in height (Santos *et al.* 2011), and CBH of 432 (n = 95; mean of 62cm) and 14cm of leaf litter (n= 30; mean of 12cm). This is the most preserved area of primary forest studied (Figure 2F). It is likely that the good conservation of the area is related to the difficulty of access, which is made by an abandoned secondary road. The collections in this area were performed only in the 2008 to 2011.

The habitats were classified using the following categories: Primary Forest (PF), with a predominance of trees of medium to large size and without significant human disturbance; Secondary forest (SF), with some degree of disturbance and impact primarily caused by logging, with few large trees; Igarapé (IG), represented by riparian vegetation present along watercourses in lowland areas; and Disturbed area (DA), deforested areas for building roads, railroads or other structures present in




FIGURE 1. Location of the study area (in red) with their collections sites: (A) Barroso; (B) Capiranga; (C) Adutora; (D) Pacoval; (E) Galiléia; (F) Mutum.

the Area under the Influence of Juruti Project (Figure 2).

We used two collection methods: Pitfall Traps (PFT) (Cechin and Martins 2000; Foster 2012) and Time Constrained Search (TCS) (Martins and Oliveira 1999). Specimens recorded through Incidental Encounters (IE), run over on roads (Road) and collected by third parties (CTP) were also considered.

The PFT were installed in the five areas (A, B, C, E and F) being arranged in a “Y” shape, 100m to 300m from one another; six trap stations were installed in each area. Each trap station consisted of four 35L buckets connected together by a drift fence, placed 10m from one another. Drift fences were buried approximately 15cm deep and were kept upright with wooden stakes that passed through the center of the buckets. Traps were checked daily during the sampling periods. The sampling effort was 7080 buckets/day (24 buckets x 5 areas x 59 days).

The time constrained search (TCS) was performed in the six areas (A-F), for four hours per expedition in each area (two during the day and two at night), with a staff of four people. The sampling effort per area was 128 hours-man (16 hours-man/day x eight expeditions).

The combined use of these methods aided in the compilation of a representative sample for the region, considering that they are complementary methods and access different faunas (Ávila-Pires *et al.* 2007; Maschio *et*

al. 2009; Foster 2012).

The collecting and transportation of the material were made under proper authorization by IBAMA (process N° 063/07; 073/08; 020/09; 011/10; 25/10). All material collected in this study is deposited in the Herpetological collection of MPEG (see Appendix). The nomenclature used herein followed the list of species for Brazilian reptiles by Bérnils and Costa (2012), considering the recent changes proposed by Gamble *et al.* (2011) for the genus *Coleodactylus*, Hedges and Conn (2012) for the genus *Mabuya* and Nicholson *et al.* (2012) for the genus *Anolis*.

RESULTS AND DISCUSSION

A total of 33 species (n= 622) of lizards were collected in the municipality of Juruti, state of Pará, belonging to 26 genera and ten families (Table 1, Figures 3 and 4). Gymnophthalmidae, with eight species (25% of the total species) was the family with highest species richness, followed by Teiidae, with six species (18% of the total species), Sphaerodactylidae with five species (15% of the total species), Dactyloidae and Tropiduridae with four species (12% of the total species) and Mabuyidae with two species (6% of the total species). In the families Gekkonidae, Iguanidae, Phyllodactylidae and Polychrotidae only one species was recorded (Table 1).



FIGURE 2. Environments sampled in the study area in the municipality of Juruti, State of Pará: (A) Igarapé, (B) Primary forest, (C) Secondary forest, (D) Disturbed area.

TABLE 1. Lizards recorded in municipality of Juruti, west of Pará, Brazil. Collection sites: B= Barroso; C= Capiroanga; A= Adutora; G= Galiléia; P= Pacoval; M= Mutum; R= Road, S= Surroundings. Methods Collection: PFT= Pitfall traps; TCS= Time Constrained Search; FR= Fauna Rescue; IE= Incidental Encounters. Habitats: PF= Primary Forest; SF= Secondary Forest; IG= Igarape; DA= Disturbed Area.

SPECIES	LOCALITY	METHODS COLLECTION	HABITAT	THIS STUDY	DATA COLLECTION (MPEG)
DACTYLOIDAE					
<i>Norops fuscoauratus</i> (D'Orbigny, 1837)	A, B, C, G, M, P, S, R	PFT, TCS, IE, FR	PF, SF, DA, IG	28	42
<i>Norops tandai</i> (Ávila-Pires, 1995)	A, B, C, G, M, P, R, S	PFT, TCS, IE	PF, SF, DA, IG	59	26
<i>Norops ortonii</i> (Cope, 1869)	M, R, S	TCS, IE, FR	PF, DA	2	7
<i>Dactyloa punctata</i> (Daudin, 1802)	A, C, G, P, M, B, R, S	PFT, TCS, IE, FR	PF, SF, DA, IG	7	13
GÉKKONIDAE					
<i>Hemidactylus mabouia</i> (Moreau de Jonnés, 1818)	C	IE	DA		1
GYMNOPHTHALMIDAE					
<i>Alopoglossus angulatus</i> (Linnaeus, 1758)	A, M	TCS, IE	PF, IG	4	1
<i>Arthrosaura reticulata</i> (O'Shaughnessy, 1881)	B, C, M, R	IE, PFT, FR	PF, SF, DA, IG	1	4
<i>Bachia flavescens</i> (Bonnaterre, 1789)	C, M, R, S	PFT, TCS, FR	PF, SF, DA, IG	2	3
<i>Cercosaura</i> sp.	A, B, C, G, P, M, R, S	PFT, IE, TCS, FR	PF, SF, DA	10	31
<i>Iphisa elegans</i> Gray, 1851	A, B, G, M	PFT, IE, TCS	PF, SF	8	8
<i>Leposoma osvaldoi</i> Ávila-Pires, 2005	A, B, C, M, P, S	PFT, TCS	PF, SF, DA, IG	8	15
<i>Leposoma percarinatum</i> (Müller, 1923)	C, M, S	TCS, PFT	PF, SF, IG		7
<i>Ptychoglossus brevifrontalis</i> Boulenger, 1912	M	PFT	PF	1	
IGUANIDAE					
<i>Iguana iguana</i> (Linnaeus, 1758)	C, R	TCS, IE	DA		2
PHYLLODACTYLIDAE					
<i>Thecadactylus rapicauda</i> (Houttuyn, 1782)	A, G, M, S	PFT, TCS, IE, FR	PF, SF, DA, IG	7	3
POLYCHROTIDAE					
<i>Polychrus marmoratus</i> (Linnaeus, 1758)	R	FR	DA		2
MABUYIDAE					
<i>Varzea bistriata</i> (Spix, 1825)	S	TCS	DA		1
<i>Copeoglossum nigropunctatum</i> (Spix, 1825)	A, C, G, P, M, B, S, R	PFT, TCS, IE	PF, SF, DA, IG	18	20
SPHAERODACTYLIDAE					
<i>Chatogekko amazonicus</i> (Andersson, 1918)	A, B, C, G, M, P, S, R	PFT, TCS, IE, FR	PF, SF, DA, IG	37	29
<i>Gonatodes hasemani</i> Griffin, 1917	M, S	PFT	PF	1	2
<i>Gonatodes humeralis</i> (Guichenot, 1855)	A, B, C, G, M, P, S, R	PFT, TCS, IE	PF, SF, DA, IG	39	37
<i>Lepidoblepharis heyerorum</i> Vanzolini, 1978	B, C, M	TCS, PFT	PF, SF		6
<i>Pseudogonatodes guianensis</i> Parker, 1935	B, C, G, M, S	IE, PFT, FR	PF, SF, DA	8	6
TEIIDAE					
<i>Ameiva ameiva</i> (Linnaeus, 1758)	A, B, C, G, S, M	TCS, PFT, IE	PF, SF, DA	13	8
<i>Crocodilurus amazonicus</i> Spix, 1825	S	IE	IG	1	
<i>Kentropyx altamazonica</i> Cope, 1876	S	TCS	PF, DA		2
<i>Kentropyx calcarata</i> Spix, 1825	A, C, P, M, B, S	PFT, TCS, IE, FR	PF, SF, DA	19	26
<i>Tupinambis longilineus</i> Ávila-Pires, 1995	S, R	FR	DA		2
<i>Tupinambis teguixin</i> (Linnaeus, 1758)	M, R	TCS, FR	PF, DA		2
TROPIDURIDAE					
<i>Plica plica</i> (Linnaeus, 1758)	A, B, G, M	PFT, TCS	PF, SF, IG	5	9
<i>Plica umbra ochrocollaris</i> (Spix, 1825)	C, G, M, S	PFT, IE, FR	PF, SF, DA	8	7
<i>Uracentron azureum</i> (Linnaeus, 1758)	C, S	PFT, FR	SF, DA	1	1
<i>Uranoscodon superciliosus</i> (Linnaeus, 1758)	A, B	TCS	IG	11	1
TOTAL				298	324

Norops tandai (n= 85), *Gonatodes humeralis* (n= 76), *N. fuscoauratus* (n= 70) and *Chatogekko amazonicus* (n= 66) were the most abundant species, being recorded in the six areas and the four types of habitat studied (Table 1).

Norops tandai is usually found in *terra firme* primary forest and, occasionally, in marginally disturbed areas. It is present south of the Amazon River (Ávila-Pires 1995, D'Angiolella et al. 2011). *Gonatodes humeralis* is considered one of the most common species in the Amazon and may occupy *terra firme* forests, primary or secondary, floodplain or wetland environments, plus gallery forests and savannas. It is a diurnal and arboreal

species, which uses tree trunks both inside forests and on the edge of forests (Ávila-Pires 1995; Vitt et al. 1997). *Norops fuscoauratus* is an arboreal and diurnal species that can be found in both primary and secondary forests. It is present in the north of South America, east of the Andes, in the following countries: Brazil, French Guiana, Suriname, Guyana, Venezuela, Colombia, Ecuador, Peru and Bolivia (Duellman 1978, 1990; Vitt 1991; Ávila-Pires 1995; Vitt et al. 2008). *Chatogekko amazonicus* is a diurnal species that inhabits the leaf litter of primary and secondary *terra firme* forests and can occasionally be found in lowland areas. Its distribution comprises eastern and central

Amazonia, including Brazil, French Guiana, Suriname, southern Guyana and southern Venezuela (Ávila-Pires 1995). The species *Chatogekko amazonicus*, *Gonatodes humeralis*, *Ameiva ameiva* e *Kentropyx calcarata* are widely distributed, occurring in different areas of the Brazilian Amazon (Cunha et al. 1985; Duellman 1990; Martins 1991; Ávila-Pires and Hoogmoed 1997; Bernardi et al. 2002; Macedo et al. 2008; Vitt et al. 2008; Ilha and Dixo 2010; Ávila-Pires et al. 2010; Silva et al. 2011; Bernardo et al. 2012) (Table 2). However, according Geurgas and Rodrigues (2010), there are molecular evidences that *C. amazonicus* may represent several distinct species.



FIGURE 3. Some species of lizards recorded in the municipality of Juruti, State of Pará: (A) *Norops tandai*; (B) *Gonatodes humeralis*; (C) *Norops fuscoauratus*; (D) *Chatogekko amazonicus*; (E) *Uracentron azureum*; (F) *Ameiva ameiva*; (G) *Uranoscodon superciliosus*; (H) *Pseudogonatodes guianensis*.

About 30% of lizard species recorded in the Juruti region, occurred in the four habitat types (PF, SF, DA and IG). Most species were recorded both in primary forest (75%) and in disturbed areas (73%). In secondary forest environments and streams (igarapés), this percentage was lower (58% and 45%, respectively) (Table 1).

Among the methods used, the TCS recorded the highest number of specimens ($n = 23$, with three unique species), followed by PFT ($n = 22$, two exclusive species) and IE ($n = 18$, two exclusive species). The number of species collected in the dry season (20 spp.) was similar to that recorded in the rainy season (22 spp.), although the sampling effort has been small in the dry season.




FIGURE 4. Some species of lizards recorded in the municipality of Juruti, State of Pará: (A) *Dactyloa punctata*; (B) *Arthrosaura reticulata*; (C) *Cercosaura* sp.; (D) *Thecadactylus rapicauda*; (E) *Norops ortonii*; (F) *Plica umbra* (Foto: A. Dourado); (G) *Kentropyx calcarata* (Foto: A. Dourado); (H) *Copeoglossum nigropunctatum* (Foto: A. Dourado).

The number of species recorded in the Juruti region (33 species) corresponds to 35% of all known lizards in the Brazilian Amazon, and 25% of lizards known throughout the Amazon region (Ávila-Pires et al. 2007). This number is higher than the number of species recorded in other studies conducted in different herpetofaunistic regions of the Brazilian Amazon (Cunha et al. 1985; Martins 1991; Ávila-Pires and Hoogmoed 1997; Bernardi et al. 2002; Gardner et al. 2007; Macedo et al. 2008; Ávila-Pires et al. 2009; Ilha and Dixo 2010; Silva et al. 2011; Bernardo et al. 2012; Pantoja and Fraga 2012) (Table 2).

If we consider the Amazon region as a whole, the number of species recorded in Juruti exceeds locations as the Cusco Amazónico (25 species) in Peru (Duellman 2005) and Santa Cecilia (30 species) in Ecuador (Duellman 1978, 1987). However, the diversity is lower than the observed in studies conducted in the Manaus region (34 species) in Brazil (Duellman 1990; Vitt et al. 2008) (Table 2) and in the Iquitos region (36 species) in Peru (Dixon and Soini 1986).

The results presented here contribute to the knowledge on the richness of Amazonian lizards and can serve as a basis for future studies of population, distribution, ecology and environmental impact (deforestation and bauxite mining) on communities of Amazonian lizards.

TABLE 2. Lizards recorded in Juruti, west of Pará, Brazil and other studies conducted in different regions of the Brazilian Amazon. **1** = Porto Walter (Ávila-Pires et al. 2009); **2** = Riozinho da Liberdade (Bernarde et al. 2011); **3** = Balbina (Martins 1991); **4** = Manaus (Duellman 1990; Vitt et al. 2008); **5** = Rio Ituxi (Ávila-Pires et al. 2009); **6** = Rio Preto da Eva (Ilha and Dixo 2010); **7** = Rio Gregório (Pantoja and Fraga 2012); **8** = Carajás (Cunha et al. 1985; Nascimento et al. 1987); **9** = Caxiuanã (Ávila-Pires and Hoogmoed 1997; Bernarde et al. 2002); **10** = Curuá-Una (Ávila-Pires et al. 2009); **11** = Barcarena (Silva et al. 2011); **12** = Trairão (Mendes-Pinto and Souza, 2011); **13** = Tapirapé (Bernardo et al. 2012); **14** = Juruti (This study); **15** = Espigão do Oeste (Macedo et al. 2008); **16** = Guajará-Mirim (Ávila-Pires et al. 2009)

SPECIES	ACRE		AMAZONAS				PARÁ					RONDÔNIA				
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
DACTYLOIDAE																
<i>Dactyloa philopunctatus</i> (Rodrigues, 1988)			3	+		+										
<i>Dactyloa punctata</i> (Daudin, 1802)	+	+			+			+	6	+	1	+		20	16	+
<i>Dactyloa transversalis</i> (Duméril, 1851)	+	+			+										5	+
<i>Norops auratus</i> (Daudin, 1802)												+				
<i>Norops brasiliensis</i> (Vanzolini and Williams, 1970)								+								
<i>Norops chrysolepis</i> (Duméril and Bibron, 1837)			17	+		+	1									
<i>Norops fuscoauratus</i> (D'Orbigny, 1837)	+	+	8	+	+	+	1	+	29	+	16	+		70	26	+
<i>Norops tandai</i> (Ávila-Pires, 2005)	+	+			+		1							85	7	
<i>Norops ortonii</i> (Cope, 1869)	+	+		+	+	+		+	3	+	1			9	+	
<i>Norops trachyderma</i> (Cope, 1876)	+	+					1			+		+				
GEKKONIDAE																
<i>Hemidactylus mabouia</i> (Moreau de Jonnés, 1818)		+		+		+			9		1			1	+	
GYMNOPHTHALMIDAE																
<i>Alopoglossus angulatus</i> (Linnaeus, 1758)	+		3	+	+		6		6		3	+		5	+	+
<i>Alopoglossus atriventris</i> Duellman, 1973	+				+											
<i>Alopoglossus buckleyi</i> (O'Shaughnessy, 1881)		+														
<i>Amapasaurus tetradactylus</i> Cunha, 1970																
<i>Arthrosaura kockii</i> (Lidth de Jeude, 1904)									44				+			
<i>Arthrosaura reticulata</i> (O'Shaughnessy, 1881)			12	+	+	+	2		16	+		+	+	5		+
<i>Bachia gr. dorbignyi</i> (Duméril and Bibron, 1839)	+															+
<i>Bachia flavescens</i> (Bonnaterre, 1789)			1	+			1		1	+		+		5		
<i>Bachia peruana</i> (Bonnaterre, 1789)		+					1									
<i>Bachia panoplia</i> Thomas, 1965				+												
<i>Cercosaura argulus</i> Peters, 1863	+	+										+			8	+
<i>Cercosaura eigenmanni</i> (Griffin, 1917)					+										8	+
<i>Cercosaura ocellata</i> Wagler, 1830	+	+			+			+	34	+		+	+	19		
<i>Cercosaura oshaughnessyi</i> (Boulenger, 1885)	+															
<i>Cercosaura</i> sp.														41		
<i>Colobosaura modesta</i> (Reinhardt and Luetken, 1862)								+								
<i>Gymnophthalmus cf. underwoodi</i> Grant, 1958																
<i>Iphisa elegans</i> Gray, 1851	+	+		+		+			1	+		+		16	10	+
<i>Leposoma</i> sp.				+									+			
<i>Leposoma guianense</i> Ruibal, 1952			5	+												
<i>Leposoma osvaldoi</i> Ávila-Pires, 2005												+		23	4	+
<i>Leposoma percarinatum</i> (Müller, 1923)			3	+		+			8	+	3	+		7		+
<i>Neusticurus</i> sp.																
<i>Neusticurus bicarinatus</i> (Linnaeus, 1758)			2	+		+		+								
<i>Neusticurus ecleopus</i> (Cope, 1876)	+							+		+		+				
<i>Neusticurus juruazensis</i> Ávila-Pires and Vitt, 1998	+															
<i>Neusticurus rudis</i> Boulenger, 1900																
<i>Ptychoglossus brevifrontalis</i> Boulenger, 1912	+	+		+	+							+		1		
<i>Tretioscincus agilis</i> (Ruthven, 1916)			3	+		+		+	18							
HOPLOCERCIDAE																
<i>Enyalioides laticeps</i> (Guichenot, 1855)	+	+			+										2	
<i>Enyalioides palpebralis</i>	+	+														
IGUANIDAE																
<i>Iguana iguana</i> (Linnaeus, 1758)		+		+	+		2	+	2		+		+	2	+	
LEIOSAURIDAE																
<i>Enyalius leechii</i> (Boulenger, 1885)															3	
MABUYIDAE																
<i>Varzea bistriata</i> (Spix, 1825)			6	+	+			+	1					1		
<i>Copeoglossum nigropunctatum</i> (Spix, 1825)	+	+		+	+	+			20	+	11	+	+	38	17	+
PHYLLODACTYLIDAE																
<i>Thecadactylus rapicauda</i> (Houttuyn, 1782)	+		2	+	+			+	12	+	2		+	10	+	+

TABLE 2. CONTINUED.

SPECIES	ACRE		AMAZONAS					PARÁ					RONDÔNIA			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<i>Thecadactylus solimoensis</i> Bergmann and Russell, 2007		+														
POLYCHROTIDAE																
<i>Polychrus marmoratus</i> (Linnaeus, 1758)				+				+		+	4			2		
LEIOSAURIDAE																
<i>Enyalius leechii</i> (Boulenger, 1885)								+		+						
SPHAERODACTYLIDAE																
<i>Chatogekko amazonicus</i> (Andersson, 1918)			11	+	+	+		+	150	+	5	+	+	66	9	+
<i>Gonatodes annularis</i> Boulenger, 1887																
<i>Gonatodes eladioi</i> Nascimento, Ávila-Pires and Cunha, 1987								+					+			
<i>Gonatodes hasemani</i> Griffin, 1917	+	+			+				3					3	29	+
<i>Gonatodes humeralis</i> (Guichenot, 1855)	+	+	18	+			+	4	+	61	+	22	+	+	76	12
<i>Lepidoblepharis heyerorum</i> Vanzolini, 1978										1				6		
<i>Pseudogonatodes</i>								1								
<i>Pseudogonatodes guianensis</i> Parker, 1935	+	+		+		+									14	
<i>Ameiva ameiva</i> (Linnaeus, 1758)	+	+	4	+	+	+	4	+	36	+	3	+	+	21	58	+
<i>Cnemidophorus</i> sp.				+		+										
<i>Cnemidophorus cryptus</i> Cole and Dessauer, 1993									27							
<i>Cnemidophorus lemniscatus</i> (Linnaeus, 1758)								+		+	8					
<i>Crocodilurus amazonicus</i> Spix, 1825				+					4					1		
<i>Dracaena guianensis</i> Daudin, 1802		+														
<i>Kentropyx altamazonica</i> (Cope, 1876)		+			+		4							2		+
<i>Kentropyx calcarata</i> Spix, 1825			18	+		+		+	50	+	15	+	+	45	4	+
<i>Kentropyx pelviceps</i> Cope, 1868	+	+			+										33	
<i>Kentropyx striata</i> (Daudin, 1802)																
<i>Tupinambis longilineus</i> Ávila-Pires, 1995						+								2		
<i>Tupinambis teguixin</i> (Linnaeus, 1758)	+	+	4	+	+	+		+	2			+		2	2	+
TROPIDURIDAE																
<i>Plica plica</i> (Linnaeus, 1758)	+	+		+	+	+		+	12	+		+		14	4	+
<i>Plica umbra</i> (Linnaeus, 1758)	+	+	1	+	+	+	1	+	14	+	7	+	+	15	2	+
<i>Stenocercus fimbriatus</i> Ávila-Pires, 1995		+														
<i>Stenocercus roseiventris</i> Duméril and Bibron, 1837	+														6	
<i>Stenocercus sinesaccus</i> Torres-Carvajal, 2005															+	
<i>Tropidurus</i> gr. <i>torquatus-hispidus</i>								+								
<i>Tropidurus hispidus</i> (Spix, 1825)					+											
<i>Uracentron azureum</i> (Linnaeus, 1758)			1	+					1					2		
<i>Uranoscodon superciliosus</i> (Linnaeus, 1758)			8	+	+			+	12	+	4	+	+	12	+	+
TOTAL	29	29	20	34	26	20	15	25	28	22	17	23	14	33	29	23

ACKNOWLEDGMENTS: We thank the Museu Paraense Emílio Goeldi and Alcoa for their support in completing this project, developed in the region of Juruti, as part of the project "Monitoramento de Fauna". We also thank Rafael Salomão for the overall coordination of the network; Angelo Dourado and João Costa for technical support during field expedition sand in the laboratory; and Marinus Hoogmoed and Teresa Cristina Ávila-Pires (Museu Paraense Emílio Goeldi) for their help in species identification. We thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Alcoa for providing technical assistance grants (ACM, FS and JFMS), a productivity scholarship (ALCP, process 308950/2011-9) and financial support (ALCP, PROTAX project 562171/2010-0; Universal project 478347/2011-1).

LITERATURE CITED

Ávila-Pires, T.C.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). *Zoologische Verhandlungen* 299:1-706.
 Ávila-Pires, T.C.S. and M.S. Hoogmoed. 1997. The Herpetofauna; p. 389-402 In P.L.B. Lisboa (ed). *Caxiuanã*. Belém: Museu Paraense Emílio Goeldi.
 Ávila-Pires, T.C.S., M.S. Hoogmoed and L.J. Vitt. 2007. Herpetofauna da Amazônia; p. 13-43 In L.B. Nascimento and M.E. Oliveira (ed.). *Herpetologia no Brasil II*. Belo Horizonte: Sociedade Brasileira de Herpetologia.
 Ávila-Pires, T.C.S., L.J. Vitt. S.S. Sartorius and P.A. Zani. 2009. Squamata (Reptilia) from four sites in southern Amazonia, with a biogeographic analysis of Amazonian lizards. *Boletim do Museu Paraense Emílio*

Goeldi. Ciências Naturais 4(2): 99-118.

Ávila-Pires, T.C.S., M.S. Hoogmoed and W.A. Rocha. 2010. Notes on the Vertebrates of northern Pará, Brazil: A forgotten part of the Guianan Region, I. Herpetofauna. *Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais* 5(1): 13-112.
 Bernarde, P.S., R.A. Machado and L.C.B. Turci, 2011. Herpetofauna da área do Igarapé Esperança na Reserva Extrativista Riozinho da Liberdade, Acre – Brasil. *Biota Neotropica* 11(3): 117-144.
 Bernardi, J.A.R., N. Rufino, R.G.N. Costa and R.A.T. Rocha. 2002. Répteis; p. 533-540 In P.L.B. Lisboa (ed). *Caxiuanã. Populações Tradicionais, Meio Físico & Diversidade Biológica*. Belém: Museu Paraense Emílio Goeldi.
 Bernardo P.H., R.A. Guerra-Fuentes, W. Matiazzi and H. Zaher. 2012. Checklist of Amphibians and Reptiles of Reserva Biológica do Tapirapé, Pará, Brazil. *Check List* 8(5): 839-846.
 Bérnils, R.S. and H.C. Costa (org.). 2012. *Brazilian reptiles – List of species. Version 2012.1*. Accessible at <http://www.sbherpetologia.org.br/>. Sociedade Brasileira de Herpetologia. Captured on October 10th, 2012.
 Capobianco, J.P.R., A. Veríssimo, A. Moreira, D. Sawyer, S. Ikeda and L.P. Pinto. 2001. *Biodiversidade na Amazônia Brasileira: Avaliação de Ações Prioritárias para a Conservação, Uso Sustentável e Repartição de Benefícios*. São Paulo: Estação Liberdade. 540 p.
 Cechin, S.Z. and M. Martins. 2000. Eficiência de armadilhas de queda (pitfall traps) em amostragem de anfíbios e répteis no Brasil. *Revista Brasileira de Zoologia* 17: 729-740.

- Cunha, O.R., F.P. Nascimento and T.C.S. Ávila-Pires. 1985. Os Répteis da área de Carajás, Pará, Brasil (Testudines e Squamata). *Publicações avulsas do Museu Paraense Emílio Goeldi* n. 40.
- D'Angiolella, A.B., T. Gamble, T.C.S. Ávila-Pires, G.R. Colli, B.P. Noonan and L.J. Vitt. 2011. *Anolis chrysolepis* Duméril & Bibron, 1837 (Squamata: Polychrotidae) revisited: Molecular phylogeny and taxonomy of the *Anolis chrysolepis* species group. *Bulletin of the Museum of Comparative Zoology* 160: 35-63.
- Dixon, J.R. and P. Soini. 1986. *The reptiles of the upper Amazon basin, Iquitos region, Peru*. Milwaukee: Milwaukee Public Museum. 154 p.
- Duellman, W.E. 1978. *The biology of an equatorial herpetofauna in Amazonian Ecuador*. Lawrence: University of Kansas. Miscellaneous Publication n. 65. 352 p.
- Duellman, W.E. 1987. Lizards in an Amazonian rain forest community: resource utilization and abundance. *National Geographic Society Research* 3: 489-500.
- Duellman, W.E. 1990. Herpetofaunas in Neotropical Rainforests: Comparative Composition, History, and Resource Use; p. 455-505 In A.H. Gentry (Ed.). *Four neotropical rainforests*. New Haven: Yale University Press.
- Duellman, W.E. 2005. *Cusco Amazónico. The lives of amphibians and reptiles in an Amazonian rainforest*. Ithaca and London: Comstock Publishing Associates. 433 p.
- Foster, M.S. 2012. Standard Techniques for Inventory and Monitoring; p. 205-272 In R.W. McDiarmid, M.S. Foster, C.W. Gibbons and N. Chernoff (ed.). *Reptile Biodiversity. Standard methods for inventory and monitoring*. Berkeley: University of California Press. 412p.
- Gamble, T., J.D. Daza, G.R. Colli, L.J. Vitt and A.M. Bauer 2011. A new genus of miniaturized and pug-nosed gecko from South America (Sphaerodactylidae: Gekkota). *Zoological Journal of the Linnean Society* 163(4): 1244-1266.
- Gardner, T.A., M.A. Ribeiro-Junior, J. Barlow, T.C.S. Ávila-Pires, M.S. Hoogmoed and C.A. Peres. 2007. The value of primary, secondary, and plantation forest for a neotropical herpetofauna. *Conservation Biology* 21(3): 775-787.
- Geurgas S.R. and M.T. Rodrigues. 2010. The hidden diversity of *Coleodactylus amazonicus* (Sphaerodactylinae, Gekkota) revealed by molecular data. *Molecular Phylogenetics and Evolution* 54: 583-593.
- Hedges, S.B., and C.E. Conn. 2012. A new skink fauna from Caribbean islands (Squamata, Mabuyinae, Mabuyinae). *Zootaxa* 3288: 1-244.
- Ilha, P. and M. Dixo. 2010. Anurans and Lizards, Rio Preto da Eva, Amazonas, Brasil. *Check List* 6(1): 17-21.
- Kottek, M., J. Grieser, C. Beck, B. Rudolf and F. Rubel. 2006. World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* 15(3): 259-263.
- Macedo, L.C., P.S. Bernarde and A.S. Abe. 2008. Lagartos (Squamata: Lacertilia) em áreas de floresta e de pastagem em Espigão do Oeste, Rondônia, sudoeste da Amazônia, Brasil. *Biota Neotropica* 8(1): 133-139.
- Martins, M. 1991. The Lizards of Balbina, Central Amazonia, Brazil: A Qualitative Analysis of Resource Utilization. *Studies on Neotropical Fauna and Environment* 26(3): 179-190.
- Martins, M. and M.E. Oliveira. 1999. Natural history of snakes in forests of the Manaus region, Central Amazonia, Brazil. *Herpetological Natural History* 6(2): 78-150.
- Maschio, G. F., M.C. Santos-Costa and A.L.C. Prudente. 2009. Comunidade de Serpentes da região de Caxiuanã com avaliação da eficiência dos métodos de captura; p. 589-603 In P.L.B. Lisboa (ed.). *Caxiuanã: Desafios para a conservação de uma Floresta Nacional na Amazônia*. Belém: Museu Paraense Emílio Goeldi.
- MCT - Ministério da Ciência e Tecnologia. 2008. Conservação e Monitoramento da Vegetação e das Unidades de Paisagem nas Áreas de Mineração da Alcoa, Município de Juruti, Estado do Pará. 1: 14-15.
- Mendes-Pinto, T.J. and S.M. Souza. 2011. Preliminary assessment of amphibians and reptiles from Floresta Nacional do Trairão, with a new snake record for the Pará state, Brazilian Amazon. *Salamandra* 47(4): 199-206.
- Nascimento, F.P., T.C.S. Ávila-Pires and O.R. Cunha. 1987. Os Répteis da área de Carajás, Pará, Brasil (Squamata). II. *Boletim do Museu Paraense Emílio Goeldi. Série Zoologia* 3(1): 33-65.
- Nicholson, K.E., B.I. Crother, C. Guyer and J.M. Savage. 2012. It is time for a new classification of anoles (Squamata: Dactyloidea). *Zootaxa* 3477: 1-108.
- Pantoja, D.L. and R. Fraga. 2012. Herpetofauna of the Reserva Extrativista do Rio Gregório, Juruá Basin, southwest Amazonia, Brazil. *Check List* 8(3): 360-364.
- Rodrigues, M.T. 2005. The Conservation of Brazilian Reptiles: Challenges for a Megadiverse Country. *Conservation Biology* 19(3): 659-64.
- Santos, M.P.D., A. Aleixo, F.M. D'horta and C.E.B. Portes. 2011. Avifauna of the Juruti Region, Pará, Brazil. *Revista Brasileira de Ornitologia* 19(2): 134-153.
- Silva, J.M.C., A.B. Rylands and G.A.B. Fonseca. 2005. O destino das áreas de endemismo da Amazônia. *Megadiversidade* 1(1): 124-131.
- Silva, F.M., A.C. Menks, A.L.C. Prudente, J.C.L. Costa, A.E.M. Travassos and U. Gallati. 2011. Squamate Reptiles from municipality of Barcarena and surroundings, state of Pará, north of Brazil. *Check List* 7(3): 220-226.
- Silva Jr, J.S. 1998. Problemas de amostragem no desenvolvimento da sistemática e biogeografia de primatas neotropicais. *Neotropical Primates* 6(1): 21-22.
- Vitt, L.J. 1991. Ecology and life history of the scansorial arboreal lizard *Plica plica* (Iguanidae) in Amazonian Brazil. *Canadian Journal Zoology* 69: 504-511.
- Vitt, L.J., P.A. Zani and A.M. Barros. 1997. Ecological Variation among Populations of the Gekkonid Lizard *Gonatodes humeralis* in Amazon Basin. *Copeia* 1: 32-43.
- Vitt, L. J., W.E. Magnusson, T.C.S. Ávila-Pires and A.P. Lima. 2008. *Guia de Lagartos da Reserva Adolfo Ducke*. Amazônia Central. Manaus Attema Design Editorial. 176p.
- Vivo, M. 1996. How many species of mammals there are in Brazil? p. 313-321 In C.E.M. Bicudo and N.A. Menezes (ed.). *Biodiversity in Brazil. A First Approach*. Campos do Jordão: Proceedings of the Workshop for the assessment of biodiversity in plants and animals.

RECEIVED: October 2012

ACCEPTED: December 2012

PUBLISHED ONLINE: February 2013

EDITORIAL RESPONSIBILITY: Pedro M. S. Nunes

Appendix. Voucher specimens collected in the municipality of Juruti, state of Pará, Brazil.

Alopoglossus angulatus (MPEG 21853, 25041, 28238, 28239, 28502); *Ameiva ameiva* (MPEG 20790, 20813, 20814, 20825, 20860, 20861, 21849, 25355, 25018, 25019, 25020, 25021, 26470, 27060, 28264, 28487, 28488, 28510, 28504, 28505, 28564); *Norops fuscoauratus* (MPEG 20791, 20793, 20802, 20803, 20804, 20809, 20822, 20840, 20857, 20873, 25401, 25402, 25403, 25404, 25405, 25406, 25407, 25408, 25409, 25410, 25411, 25412, 25413, 25414, 25415, 25416, 25417, 25418, 25419, 25420, 25421, 25422, 25423, 25424, 25425, 25426, 25427, 25428, 25429, 25431, 25432, 25039, 25040, 25119, 25120, 26471, 26472, 26473, 26474, 26475, 26476, 26477, 26478, 27812, 27813, 27814, 27815, 27816, 27065, 27066, 27067, 27068, 27069, 27070, 27071, 28246, 28247, 28248, 28568); *Norops tandai* (MPEG 20785, 20794, 20810, 20811, 20819, 20847, 20854, 20856, 21858, 21854, 21856, 21852, 21851, 25338, 25339, 25341, 25340, 25342, 25343, 25344, 25345, 25346, 25347, 25348, 25349, 25350, 25027, 25028, 25029, 25030, 25102, 25103, 25104, 25105, 25106, 25107, 25108, 25109, 25110, 26479, 6480, 26481, 26482, 26483, 26484, 26485, 26486, 26487, 26488, 26489, 26490, 26491, 26492, 26493, 27818, 27819, 27820, 27821, 27822, 27823, 27824, 27073, 27074, 27075, 27076, 27077, 27078, 27079, 27080, 27081, 27082, 27083, 27084, 27085, 27086, 27087, 28249, 28250, 28251, 28252, 28253, 28254, 28255, 28505, 28578); *Norops ortonii* (MPEG 20866, 20879, 25375, 25376, 25377, 25378, 25379, 27088, 28256); *Dactyloa punctata* (MPEG 20778, 20846, 21863, 25365, 25366, 25367, 25368, 25369, 25370, 25371, 25372, 25373, 25374, 25038, 25131, 27817, 27810, 27089, 28489, 28567); *Arthrosaura reticulata* (MPEG 25351, 25354, 25352, 25353, 28501); *Bachia flavescens* (MPEG 21871, 25385, 25386, 25024, 25121); *Cercosaura* sp. (MPEG 20779, 20781, 20783, 20784, 20823, 20834, 20850, 21870, 21855, 25318, 25319, 25320, 25321, 25322, 25323, 25203, 25334, 25324, 25325, 25326, 25327, 25328, 25329, 25331, 25332, 25333, 25335, 25336, 25337, 25330, 25025, 25026, 26494, 26495, 26496, 26497, 26498, 26499, 28240, 28241); *Chatogekko amazonicus* (MPEG 20786, 20792, 20796, 20824, 20835, 20839, 20848, 21874, 21877, 21866, 21879, 21878, 21873, 21876, 25272, 25273, 25274, 25275, 25276, 25277, 25278, 25279, 25280, 25281, 25282, 25283, 25284, 25285, 25286, 25287, 25031, 25032, 25033, 25034, 25035, 25036, 25037, 25122, 25123, 27811, 26500, 26501, 26502, 26503, 26504, 26505, 26506, 26507, 26508, 26509, 27825, 27826, 27827, 27828, 27829, 27092, 27093, 27094, 27095, 27096, 27097, 27098, 28257, 28258, 28259, 28571, 28572); *Crocodylus amazonicus* (MPEG 26510); *Gonatodes hasemani* (MPEG 20787, 25357, 25133); *Gonatodes humeralis* (MPEG 20797, 20806, 20807, 20808, 20878, 21865, 21875, 21868, 21872, 21861, 21869, 21867, 21857, 21860, 25220, 25221, 25222, 25223, 25225, 25226, 25224, 25227, 25228, 25229, 25230, 25231, 25232, 25233, 25234, 25235, 25236, 25237, 25238, 25239, 25240, 25241, 25242, 25042, 25043, 25044, 25045, 25046, 25047, 25048, 25049, 25050, 25051, 25052, 25053, 25111, 25112, 25113, 25114, 25115, 26511, 26512, 26513, 26514, 26515, 26516, 26517, 26518, 26519, 26520, 27830, 27831, 27832, 27833, 27834, 27835, 27100, 27101, 28260, 28261, 28262, 28506); *Hemidactylus mabouia* (MPEG 25292); *Iguana iguana* (MPEG 20869, 25077); *Iphisa elegans* (MPEG 20821, 20832, 25256, 25257, 25258, 25259, 25260, 25261,

26521, 28242, 28243, 28244, 28491, 28490, 28507, 28565); *Kentropyx altamazonica* (MPEG 20868, 20870); *Kentropyx calcarata* (MPEG 20782, 20799, 20800, 20801, 20805, 20818, 20820, 20831, 20838, 20841, 20842, 20843, 20852, 20862, 20864, 21862, 21864, 25309, 25310, 25311, 25312, 25313, 25314, 25317, 25011, 25012, 25014, 25015, 25016, 25013, 25017, 25124, 25125, 25126, 25127, 25128, 25129, 26522, 26523, 26524, 27836, 27837, 28265); *Lepidoblepharis heyerorum* (MPEG 21859, 25243, 25244, 25245, 25246, 25247); *Leposoma osvaldoi* (MPEG 20780, 20788, 20789, 20798, 20827, 20830, 20855, 25248, 25249, 25250, 25251, 25252, 25253, 25254, 25255, 26525, 26526, 26527, 28245, 28508, 28492, 28576, 28577); *Leposoma percarinatum* (MPEG 20872, 20876, 25267, 25268, 25269, 25270, 25271); *Varzea bistriata* (MPEG 20874); *Copeoglossum nigropunctatum* (MPEG 20795, 20815, 20817, 20844, 20851, 20867, 20877, 20871, 21847, 21848, 25293, 25294, 25295, 25296, 25297, 25298, 25299, 25300, 25301,

25302, 25022, 25132, 26528, 26529, 26530, 26531, 26532, 26533, 26534, 26535, 26536, 27102, 28493, 28494, 28509, 28509, 28569, 28570); *Plica plica* (MPEG 20812, 20816, 20829, 20836, 20849, 20853, 20858, 20859, 25308, 26537, 26538, 28266, 28495, 28563); *Plica umbra ochrocollaris* (MPEG 20828, 20833, 20845, 20863, 20880, 20881, 25381, 25023, 25134, 26539, 27839, 27103, 28267, 28497, 28496); *Polychrus marmoratus* (MPEG 25387, 25388); *Pseudogonatodes guianensis* (MPEG 20865, 25262, 25263, 25264, 25266, 25265, 25054, 25055, 26540, 27104, 28263, 28511, 28573, 28574, *Ptychoglossus brevifrontalis* (MPEG 26541); *Thecadactylus rapicauda* (MPEG 20875, 25358, 25359, 25130, 26542, 27105, 28499, 28512, 28498, 28566); *Tupinambis longilineus* (MPEG 25382, 25383); *Tupinambis teguixin* (MPEG 21880, 25384); *Uracentron azureum* (MPEG 25396, 28500); *Uranoscodon superciliosus* (MPEG 25356, 25116, 25117, 26543, 26544, 26545, 27840, 27841, 27842, 27106, 27107, 28268).