

Uroderma magnirostrum Davis, 1968 (Chiroptera: Phyllostomidae): First record from the state of Sergipe, northeastern Brazil

Patrício A. da Rocha^{1*}, José Anderson Feijó¹, Juan Ruiz-Esparza² and Stephen F. Ferrari³

- 1 Programa de Pós-Graduação em Zoologia, Universidade Federal da Paraíba, Campus I, Departamento de Sistemática e Ecologia. CEP 58059-900. João Pessoa, PB, Brazil,
 - 2 Programa de Pós-Graduação em Desenvolvimento e Meio Ambiente – PRODEMA, Universidade Federal de Sergipe. CEP 49100-000. São Cristóvão, SE, Brazil.
 - 3 Universidade Federal de Sergipe. Departamento de Biologia, CEP 49100-000. São Cristóvão, SE, Brazil.
- * Corresponding author. E-mail: parocha2@yahoo.com.br

ABSTRACT: This study provides the first record of *Uroderma magnirostrum* Davis, 1968 from the state of Sergipe in the Brazilian northeast, based on the capture of two specimens, one male and one female. The morphometric data and morphological characters were consistent with those recorded for the species at other Brazilian sites. This record extends the distributional range of the species within South America approximately 220 km eastwards.

The genus *Uroderma* Peters, 1866 includes two species of medium-sized bats, *Uroderma bilobatum* Peters, 1866 and *Uroderma magnirostrum* Davis, 1968, both of which occur in Brazil (Simmons 2005; Gardner 2007; Peracchi *et al.* 2011). *Uroderma magnirostrum* has been recorded in Bolivia, Colombia, Ecuador, Guyana, Peru, and Venezuela (Gardner 2007). In Brazil, the species has been recorded in the Amazon basin (Martins *et al.* 2006; Bernard and Fenton 2002; Bernard *et al.* 2011), Pantanal (Alho *et al.* 2011), Cerrado (Pine *et al.* 1970; Marinho-Filho 1996), Atlantic Forest (Peracchi and Albuquerque 1993; Nogueira *et al.* 2003; Tavares *et al.* 2007), and Caatinga (Mares *et al.* 1981; Willig and Mares 1989).

Despite its broad geographic distribution and the variety of habitats occupied by the species, *U. magnirostrum* appears to be relatively rare locally (Willig 1983; Nogueira *et al.* 2003). In the Brazilian northeast, *U. magnirostrum* has been recorded at only three localities (Mares *et al.* 1981) (Figure 1). This is the first record of *U. magnirostrum* for the Brazilian state of Sergipe, and represents an approximate 220 km extension of its distribution to the east.

Two specimens of *U. magnirostrum* were captured during a survey of the bat fauna of Serra da Guia, in Poço Redondo, Sergipe (09°58' S, 37°52' W), between May and July, 2009 (Figure 1). The specimens were captured in mist-nets (100 x 2.5m) in an area of arboreal caatinga ("caatinga alta" of Mares *et al.* 1981: p 89), located at the foothills of the Serra da Guia. The study area is covered by this vegetation – dominated by arboreal cacti and trees of the Leguminosae and Euphorbiaceae families – up to an altitude of 650 m a.s.l. The specimens collected are an adult male (UFPB 6107) and an adult female (UFPB 6106), both were fixed in 10% formalin, preserved in 70% ethanol, with subsequent extraction of the skull, and deposited in the Zoological Collections of the Federal University of Paraíba in João Pessoa.

The genus *Uroderma* can be distinguished from all other stenodermatines by the parallel and markedly bifid




FIGURE 1. Records of *Uroderma* in the Brazilian Northeast (bold outline). Biomes: dark gray = Atlantic Forest; medium gray = Cerrado; light gray = Caatinga; white = Amazon. Black Circle = *U. magnirostrum* (1) Km 18, federal highway BR 316, 18 km south of Teresina, Piauí; (2) Exu, Pernambuco; (3) Araripe-Apodí National Forest, 8 km south-southwest of Crato, Ceará (Mares *et al.* 1981); Black Triangle = *U. magnirostrum* (4) Serra da Guia, Poço Redondo, Sergipe (this study); Black Square = *U. bilobatum* (5) São Cristóvão, Sergipe (Rocha *et al.* 2010), (6) Ilhéus, Bahia; (7) Una, Bahia; (8) Itapebi, Bahia (Faria *et al.* 2006).

upper internal incisors, and the contact between the four lower incisors (Gardner 2007). *Uroderma magnirostrum* can be differentiated from *U. bilobatum* by its heavy, deep

rostrum, and by the gradual, nearly straight slope of the dorsal profile of the skull, from the crown to the tip of the snout (Figure 2), and the marked lateral expansion of the mesethmoid bone, which forms a shield-like structure as seen in the frontal view (Figure 3). The facial stripes of *U. magnirostrum* are also poorly developed or absent, and the internal surface of the ear is unicolor, with no white or

yellowish edging (Davis 1968). All the external and cranial characters of the two *U. magnirostrum* specimens collected in the present study are in accordance with characters described by previous authors (Davis, 1968; Nogueira et al. 2003). Additionally, cranial measurements taken from the specimens from Sergipe fall within the range recorded for specimens collected in southeastern Brazil (Table 1).



FIGURE 2. Lateral view of the skull of (A) *Uroderma magnirostrum* (UFPB 6107) and (B) *Uroderma bilobatum* (UFPB 5281). Scale bar = 10 mm.



FIGURE 3. Frontal view of the skull of (A) *Uroderma magnirostrum* (specimen UFPB 6107) and (B) *Uroderma bilobatum* (specimen UFPB 5281). The mesethmoid bone is indicated by the white arrow.

TABLE 1. Cranial measurements of the *Uroderma magnirostrum* specimens collected during the present study, at Serra da Guia, Sergipe, and at other localities in southeastern Brazil. ¹ Data from Nogueira et al. (2003) for specimens collected from Jaíba and Timóteo (Minas Gerais), Linhares (Espírito Santo), and Rio de Janeiro.

Parameter	Present study		Southeastern Brazil ¹	
	Male (UFPB 6107)	Female (UFPB 6106)	Male (n=7)	Female (n=5)
Greatest length of skull	22.84	23.30	23.54 ± 0.54	22.99 ± 0.22
Condylbasal length	21.15	21.59	21.04 ± 0.56	20.62 ± 0.22
Mastoid breadth	10.75	10.92	10.94 ± 0.20	10.92 ± 0.22
Zygomatic breadth	-	12.52	12.72 ± 0.62	12.67 ± 0.31
Breadth of braincase	9.23	9.66	9.54 ± 0.34	9.41 ± 0.25
Postorbital constriction	5.64	5.89	5.82 ± 0.20	5.69 ± 0.15
Palatal length	10.96	11.87	-	-
Palatal breadth	4.68	4.62	-	-
Breadth across upper canines	5.20	5.31	5.71 ± 0.20	5.58 ± 0.12
Breadth across upper molars	8.66	9.01	8.94 ± 0.35	9.11 ± 0.14
Length of maxillary toothrow	7.26	8.30	8.89 ± 0.23	8.87 ± 0.22
Length of mandible	15.04	15.54	15.81 ± 0.44	15.40 ± 0.24
Height of ramus at the coronoid process	4.74	5.21	-	-

While *U. magnirostrum* and *U. bilobatum* are sympatric throughout much of tropical South America (Davis 1968) their distribution in the Brazilian northeast does not appear to follow this pattern (Figure 1), as in this region *U. magnirostrum* occurs exclusively in the semi-arid Caatinga, and *U. bilobatum* in the coastal Atlantic Forest. This scenario may reflect the distinct mosaic of dry and wet climate and vegetation associated of the Brazilian northeast in comparison with the rest of the geographic occurrence of the genus, and may further indicate that *U. magnirostrum* can tolerate arid conditions, *i.e.* the Caatinga better than *U. bilobatum*. This condition is also suggested by the non-overlapping distribution of *U. bilobatum* and *U. magnirostrum* within Sergipe in two closely located sites representative of the Atlantic Forest (Rocha et al. 2010) and Caatinga biomes (this study) respectively.

ACKNOWLEDGMENTS: We are grateful to Jadson Luis da Silva Brito and Carolina Nunes Liberal for technical assistance, to CAPES (PAR, JAF) and FAPITEC (JMRA) for graduate fellowships and CNPq for a research grant to SFF (process no. 302747/2008-7).

LITERATURE CITED

- Alho, C.J.R., E. Fischer, L.F. Oliveira-Pissini and C.F. Passos. 2011. Bat-species richness in the Pantanal floodplain and its surrounding uplands. *Brazilian Journal of Biology* 71(1): 311-320.
- Bernard, E. and M.B. Fenton. 2002. Species diversity of bats (Chiroptera: Mammalia) in forest fragments, primary forests and savannas in Central Amazonia, Brazil. *Canadian Journal of Zoology* 80: 1124-1140.
- Bernard, E., V.C. Tavares and E. Sampaio. 2011. Updated compilation of bat species (Chiroptera) for the Brazilian Amazonia. *Biota Neotropica* 11(1): *in press*.
- Davis, W. B. 1968. Review of the genus *Uroderma* (Chiroptera). *Journal of Mammalogy* 49: 676-698.
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from the Atlantic rainforest of southern Bahia, Brazil. *Biota Neotropica* 6(2): 1-13
- Gardner, A.L. 2007. Subfamily Stenodermatina; p. 300-375 In A.L. Gardner (ed.). *Mammals of South America: marsupials, xenarthrans, shrews, and bats*. Volume 1. Chicago and London: University of Chicago Press.
- Mares, M. A., M. R. Willig, K.E. Streilein and T.E. Lacher. 1981. The mammals of northeastern Brazil: a preliminary assessment. *Annals of the Carnegie Museum* 50: 81-137.
- Marinho-Filho, J.S. 1996. The Brazilian Cerrado fauna and its conservation. *Chiroptera Neotropica* 2: 37-38.
- Martins, A.C.M., E. Bernard and R. Gregorin. 2006. Inventários biológicos rápidos de morcegos (Mammalia, Chiroptera) em três unidades de conservação do Amapá, Brasil. *Revista Brasileira de Zoologia* 23:1175-1184.
- Nogueira M.R., V.C. Tavares and A.L. Peracchi. 2003. New records of *Uroderma magnirostrum* Davis (Mammalia, Chiroptera) from southeastern Brazil, with comments on its natural history. *Revista Brasileira de Zoologia* 20: 691-697.
- Peracchi, A.L. and S. T. Albuquerque. 1993. Quirópteros do município de Linhares, Estado do Espírito Santo, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Biologia*. 53: 575-581.
- Peracchi, A.L., I.P. Lima, N.R. Reis, M.R. Nogueira and H. Ortêncio Filho. 2011. Ordem Chiroptera; p. 155-234 In N.R. Reis, A.L. Peracchi, W.A. Pedro, and I.P. Lima (ed.). *Mamíferos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina. 440 p.
- Pine, R.H., I.R. Bishop and R.L. Jackson. 1970. Preliminary list of mammals of the Xavantina/Cachimbo expedition (Central Brazil). *Transactions of the Royal Society of Tropical Medicine and Hygiene* 64: 668-670.
- Rocha, P.A., J.S. Mikalauskas, S.F. Gouveia, V.V.B. Silveira, and A.L. Peracchi. 2010. Morcegos (Mammalia, Chiroptera) capturados no Campus da Universidade Federal de Sergipe, com oito novos registros para o estado. *Biota Neotropica* 10(3): 183-188.
- Tavares, V.C., F.A. Perini and J.A. Lombardi. 2007. The bat communities (Chiroptera) of the Parque Estadual do Rio Doce, a large remnant of Atlantic Forest in southeastern Brazil. *Lundiana* 8: 35-47.
- Willig, M.R. 1983. Composition, microgeographic variation, and sexual dimorphism in Caatingas and Cerrado bat communities from northeast Brazil. *Bulletin of the Carnegie Museum of Natural History* 23:1-131.
- Willig, M. R. and M. A. MARES. 1989. Mammals of the Caatinga: an updated list and summary of recent research. *Revista Brasileira de Biologia* 49: 361-367.

RECEIVED: August 2011

LAST REVISED: October 2011

ACCEPTED: October 2011

PUBLISHED ONLINE: December 2011

EDITORIAL RESPONSIBILITY: Valeria C. Tavares