

# Gastropoda, Caenogastropoda, Eulimidae, *Annulobalcis aurisflamma* Simone and Martins, 1995: First record to northeastern Brazil

Vinicius Queiroz<sup>1\*</sup>, Licia Sales<sup>1</sup>, Cláudio L. S. Sampaio<sup>2</sup>, Elizabeth G. Neves<sup>1</sup> and Rodrigo Johnsson<sup>1</sup>

<sup>1</sup> Universidade Federal da Bahia, Instituto de Biologia, Departamento de Zoologia. Avenida Adhemar de Barros s/n<sup>o</sup>, Campus Ondina. CEP 40170-290. Salvador, BA, Brazil.

<sup>2</sup> Universidade Federal de Alagoas, Campus Arapiraca. Unidade de Ensino Penedo. Avenida Beira Rio, Centro Histórico. CEP 57200-000. Penedo, AL, Brazil.

\* Corresponding author. E-mail: [vinicius UFBA@yahoo.com.br](mailto:vinicius UFBA@yahoo.com.br)

**ABSTRACT:** The current article provides the first record of *Annulobalcis aurisflamma* Simone and Martins, 1995 outside São Paulo state, Brazil. Herein we extend its geographical distribution to northeastern Brazil.

The biodiversity of marine invertebrates in Brazil is undoubtedly underestimated and the amount of new species being described is continuously increasing with their distributional records being constantly updated. Even in a well established group as Mollusca in which the shell facilitates its preservation, the most relevant book to the knowledge of Brazilian mollusks (Rios 1994; 2009) quickly becomes out of date due to the publication of many new descriptions and records of distribution (e.g. Absalão 2009; Simone 2009; Wiggers and Veitenheimer-Mendes 2009; Breves-Ramos *et al.* 2010; Caetano *et al.* 2010; Dacosta *et al.* 2010; Lima *et al.* 2010; Oliveira and Absalão 2010; Pimpão *et al.* 2010).

Among the gastropods, the family Eulimidae Philippi, 1853 is known as a symbiont of echinoderms (Barel and Kramers 1977; Warén 1983). Currently all Echinodermata classes have been recorded in association with these snails (Bacci 1948; Waren and Sibuet 1981; Cantera and Neira 1987; Waren and Crossland 1991; Lyskin and Britayev 2005; Crandall *et al.* 2008). A large spectrum of symbiosis has been registered among these two taxa, ranging from complex levels of endoparasitism (Altnöder *et al.* 2007) to relatively simple ectoparasites (Dgebuadze and Kantor 2009). Within the eulimids ectoparasites two different behaviors can be observed. The development of a specialized proboscis to feed on dermal tissues and body fluids leading to a permanent attachment to the host, and the mobile species that can move freely over the host while feeding on it (Väitilinen 2004) as in *Annulobalcis* Habe, 1965.


According to Warén (1983) *Annulobalcis* is characterized by relatively large species, showing a wide shell aperture, outer lip strongly projected and highly colored mantle. The genus is composed of three species recorded from the Pacific Ocean, *A. shimazui* Habe, 1965; *A. yamamotoi* Habe, 1974 e *A. marshalli* Warén, 1981, and two Atlantic species, *A. procera* Simone, 2002 and *A. aurisflamma* Simone and Martins, 1995 (Warén 1981;

Warén 1983; Simone and Martins 1995; Simone 2002).

The aim of this paper is to report for the first time the occurrence of *Annulobalcis aurisflamma* out of São Paulo State coast, expanding its distribution up to northeastern Brazil.

The sample was gathered on January 6<sup>th</sup>, 2010 in Barra beach (Figure 1) Salvador-BA, Brazil (13°00'37" S, 38°31'49" W) at a depth of three meters in large natural reef pools. Altogether 32 crinoids of the species *Tropiometra carinata carinata* (Lamarck, 1816) were observed and only five of them were parasitized. Eleven specimens of *A. aurisflamma* were collected from these five specimens of *Tropiometra carinata carinata*. Two specimens of *A. aurisflamma* were deposited in the Mollusca Collection of the Museu de Zoologia da Universidade de São Paulo (MZUSP 98545) and two in the Malacological Collection of the Instituto de Biologia da Universidade Federal do Janeiro (IBUFRJ 19082 and IBUFRJ 19083).

*Annulobalcis aurisflamma* species (Figure 2 A-C) shows all characteristics provided in the original description (Simone and Martins 1995). The host species, *Tropiometra carinata carinata* (Figure 2 G-D), is identified due to its small size. It bears ten arms emerging from five radial



**FIGURE 1.** Geographical distribution of *Annulobalcis aurisflamma*. Star = type locality (Ubatuba) reported by Simone and Martins (1995); circle = second record (São Sebastião channel) reported by L.F. Netto (*unpublished dissertation*); triangle = present record. Locality abbreviations: BA – Bahia State; SP – São Paulo State; BTS – Todos os Santos Bay.

stalks and ending abruptly (Figure 2D). Pinnules arranged in two rows and continuously exposed (Figure 2E). Anus laterally displaced and mouth in the center of calyx which bears an average of 21 cirrus (Figures 2D and 2F). Color variable, ranging from reddish brown with reddish and yellowish bands up to completely yellowish (Clark 1947).

Field observations registered the snails only on the aboral region of the arms, close to the calyx (Figure 3). Further observations during sample procedures confirmed statement made by Simone and Martins (1995) that *A. aurisflamma* is able to easily release from its host probably due to the slightest indication of stress.

It can be inferred that *Annulobalcis* is specifically a crinoid associate except for *A. procera* for which the host is unknown (Warén 1981; Morton and Mladenov 1989; Simone and Martins 1995; Simone 2002). According to Warén (1982; 2008) specificity in generic level can be noted for Eulimidae that usually have its hosts well documented. These associations are so well established that paleontologists use them as tool for identification of poorly preserved fossil echinoderms (Lozouet and Dockery 2001).

Based on the present results, the geographical distribution of *Annulobalcis aurisflamma*, originally




**FIGURE 3.** Photography in situ of *Annulobalcis aurisflamma* parasiting on *Tripometra carinata carinata*.

restricted to São Paulo coast in Ubatuba (Simone and Martins 1995) and São Sebastião channel (L.F. Netto unpublished dissertation), extends to northeastern Brazil, in Salvador, Bahia State. This record implies that the species previously confined to the zoogeographical Paulista Province is now observed as far as the Tropical Province (Palácio 1980; Floeter and Soares-Gomes 1999).

#### LITERATURE CITED

- Absalão, R.S. 2009. New small deep-sea species of Gastropoda from the Campos Basin off Brazil. *American Malacological Bulletin* 27(1-2): 133-140.
- Altnöder, A., J.M. Bohn, I.M. Rückert and E. Schwabe. 2007. The presumed shelled juvenile of the parasitic gastropod *Entocolax schiemenzii* Voigt, 1901 and its holothurian host *Chiridota pisanii* Ludwig, 1886. *Spixiana* 30(2): 187-199.
- Bacci, G. 1948. *Melanella comatulicola* (Graff), un Gasteropodo parassita della *Antedon mediterranea* (Lam.). *Italian Journal of Zoology* 15(1): 89-98.
- Barel, G.D.N and P.G.N. Kramers. 1977. A survey of the echinoderm associates of the north-east Atlantic area. *Zoologische Verhandlungen* 156: 3-159.
- Breves-Ramos, A., A.D. Pimenta, M.T.M. Szechy and A.O.R. Junqueira. 2010. Mollusca, Bivalvia, Mytilidae, *Myoforceps aristatus* (Dillwyn, 1817): Distribution and new record localities at Ilha Grande Bay, Brazil. *Check List* 6(3): 408-409.
- Caetano, C.H.S., V. Scarabino and R.S. Absalão. 2010. Brazilian species of *Gadila* (Mollusca: Scaphopoda: Gadilidae): rediscovery of *Gadila elongate* comb. nov. and shell morphometrics. *Check List* 27(2): 305-308.
- Cantera, J.R.K and R.O. Neira. 1987. Primer registro del genero *Echineulima* Lutzen y Nielsen (Gastropoda: Eulimidae), Moluscos parasitos de erizos de mar en la isla de Gorgona (Pacífico colombiano). *Anais do Instituto de investigacion Marinhas - Punta de Betín* 17: 87-93.
- Clark, A.H. 1947. A monograph of the existing crinoids: Superfamily Mariametrida (concluded the family Colobometridae) and Superfamily Tropiometrida (except the families Thalassometridae and Charitometridae). *Bulletin of United States National Museum* 82(4b): 1-534.
- Crandall, E.D., M.E. Jones, M.M. Muñoz, B. Akinronbi, M.V. Erdmann and P.H. Barber. 2008. Comparative phylogeography of two seastars and their ectosymbionts within the Coral Triangle. *Molecular Ecology* 17: 5276-5290.
- Dacosta, S., V. Padula and M.A. Schrodler. 2010. New Species of *Hypselodoris* and a Redescription of *Hypselodoris picta lajensis* (Nudibranchia: Chromodorididae) from Brazil. *Veliger* 51(1): 15-25.
- Floeter, S.R and A. Soares-Gomes. 1999. Biogeographic and species richness patterns of Gastropoda on the southwestern Atlantic. *Revista Brasileira de Biologia* 59(4): 567-575.
- Lima, S.F.B., J.C.N. Barros and J.A. Francisco. 2010. A new deep-sea species of Mitromorpha (Gastropoda: Conoidea: Conidae) off Brazil. *Journal of the Marine Biological Association of the United Kingdom* 90(3): 599-603.
- Lyskin, S.A and T.A. Britayev. 2005. Symbionts of holothurians from South Vietnam: Intra- and interspecific interactions. *Doklady Biological Sciences* 401(1): 116-119.



**FIGURE 2.** *Annulobalcis aurisflamma* (A, B and C) and its host *Tropiometra carinata carinata* (D, E, F and G): A - dorsal view. B and C - ventral view. D - Oral view of the calyx showing mouth and anus. E - Arm showing some pinnules. F - Aboral view of the calyx showing the cirrus. G - Cirrus in detail. Scale lines: A, B, C, F and G = 1 mm. D = 2 mm. E = 3mm.

- Lozouet, P and D.T. Dockery. 2001. First fossil record of the genus *Pyramidelloides* (Mollusca, Gastropoda, Eulimidae) discovered in the Eocene deposits of Alabama. *C. R. C. Acad. Sci. Paris, Sciences de la Terre et des planètes / Earth and Planetary Sciences* 333: 351-356.
- Oliveira, C.D.C and R.S. Absalão. 2010. Review of the Septibranchia (Mollusca: Pelecypoda) from the deep sea of Campos Basin, Brazil: Family Lyonsiellidae, with description of a new species. *Scientia Marina* 74(2): 305-316.
- Morton, B and P.V. Mladenov. 1989. The associates of *Tropiometra afra macrodiscus* (Echinodermata: Crinoidea) in Hong Kong; p. 431-438. In B. Morton (ed.). *The Marine Flora and Fauna of Hong Kong and Southern China*. Hong Kong: Proceedings of Fourth International Marine Biological Workshop.
- Palacio, F.J. 1980. Revisión zoogeográfica marina del sur del Brasil. *Boletim do Instituto Oceanográfico* 31(1): 69-92.
- Pimpao, D.M., I.L. Veitenheimer-Mendes and F. Scarabino. 2010. Mollusca, Bivalvia, Cuspidariidae, *Plectodon braziliensis* (E. A. Smith, 1915) n. comb.: Record of the genus for the South Atlantic, off Brazil. *Check List* 6(4): 648-651.
- Rios, E.C. 1994. *Seashells of Brazil*. Rio Grande: Fundação Universitária de Rio Grande. 368 p.
- Rios, E.C. 2009. *Compendium of Brazilian Sea shells*. Rio Grande: Evangraf. 668 p.
- Santos, S.B., L.E.M. Lacerda and I.C. Miyahira. 2009. *Uncancylus concentricus* (Mollusca, Gastropoda, Ancyliidae): New occurrence in state of Rio de Janeiro, Brazil. *Check List* 5(3): 513-517.
- Simone, L.R.L. 2009. New Gastropods from the São Pedro e São Paulo Archipelago, Brazil (Vetigastropoda and Caenogastropoda). *Strombus* 16(1-2): 11-18.
- Simone, L.R.L and C.M. Martins. 1995. *Annulobalcis aurisflama*, a new species of Eulimidae (Gastropoda, Prosobranchia) parasitic on a crinoid from Brazil. *Journal of Conchology* 35: 223-235.
- Vaitilingon, D., I. Eackhaunt, D. Fourgon and M. Jangoux. 2004. Population dynamics, infestation and host selection of *Vexilla vexillum*, an ectoparasitic muricid of echinoids, in Madagascar. *Diseases of Aquatic Organisms* 61: 241-255.
- Warén, A. 1981. Eulimid gastropods parasitic on echinoderms in the New Zealand Region. *New Zealand Journal of Zoology* 8(3): 313-324.
- Warén, A. 1983. A generic revision of the family Eulimidae (Gastropoda, Prosobranchia). *Journal of Molluscan Studies, Supplement* 13: 1-96
- Warén, A. 2008. Eulimidae; p. 714-721 In G.T. Poppe (ed.). *Philippine Marine Mollusks*, vol 1, Gastropoda Part 1. Hackenheim: Germany.
- Warén, A and M.R. Crossland. 1991. Revision of *Hypermastus* Pilsbry, 1899 and *Turveria* Berry, 1956 (Gastropoda: Prosobranchia: Eulimidae), Two Genera Parasitic on Sand Dollars. *Records of the Australian Museum* 43(1): 85-112.
- Warén, A and M. Sibuet. 1981. *Ophieulima* (Mollusca, Prosobranchia), a new genus of ophiuroid parasites. *Sarsia* 66: 103-107.
- Wiggers, F and I. Veitenheimer-Mendes. 2009. First record and new combination of a rissoid gastropod from off south Brazil. *Biotemas* 22(4): 251-254.

RECEIVED: April 2011

LAST REVISED: August 2011

ACCEPTED: August 2011

PUBLISHED ONLINE: October 2011

EDITORIAL RESPONSIBILITY: Luis E. Arruda Bezerra