

Mollusca, Gastropoda, Heterobranchia, Ancyliidae, *Gundlachia radiata* (Guilding, 1828): First record of occurrence for the northwestern region of Argentina

Ximena Maria Constanza Ovando^{1*}, Luiz Eduardo Macedo de Lacerda² and Sonia Barbosa dos Santos²

¹ Facultad de Ciencias Naturales e Instituto Miguel Lillo. Miguel Lillo 205. CP 4000 Tucumán, Argentina.

² Universidade do Estado do Rio de Janeiro, Instituto de Biologia Roberto Alcântara Gomes, Laboratório de Malacologia. Rua São Francisco Xavier 524. PHL 525/2, CEP 20550-900, Rio de Janeiro, RJ, Brasil.

* Corresponding author. E-mail: xco1303@hotmail.com

ABSTRACT: In the present paper we report for the first time the presence of *Gundlachia radiata* (Guilding, 1828), in northwestern region (Jujuy province), Argentina. Adult and juveniles specimens of this freshwater limpet were collected in two temporary water bodies. This record represents the first report of this species in Argentina but also is the southernmost point of occurrence of *G. radiata* in South America. As a result, the distributional range of this species is increased and the species richness of Ancyliidae in Argentina is incremented to a total of seven species classified in four genera.

The Ancyliidae *sensu lato* are freshwater pulmonate snails, characterized by a pateliform shell. Ancyliidae are cosmopolitan, and according to Santos (2003) there are seven genera in South America: *Anisancylus* Pilsbry, 1924; *Gundlachia* Pfeiffer, 1849; *Hebetancylus* Pilsbry, 1913; *Uncancylus* Pilsbry, 1913; *Burnupia* Walker, 1912; *Ferrissia* Walker, 1913 and *Laevapex* Walker, 1903. Distributional data as well as the systematic position of Argentinean Ancyliidae are scarcely known with only few species records for this country.

In this paper, we report the first record of occurrence of *Gundlachia radiata* (Guilding, 1828) (Figure 1) in northwestern Argentina. This report is based on the discovery of specimens collected in two small temporary water bodies extending the geographic distribution of this species to the subtropical region (Figure 2).

Nowadays, the genus *Gundlachia* comprises six nominal species distributed in South America: *G. bakeri* Pilsbry, 1913; *G. lutzi* Walker, 1925; *G. radiata* (Guilding, 1828); *G. saulcyana* (Bourguignat, 1853), *G. ticaga* (Marcus & Marcus, 1962) and *G. leucaspis* (Ancey 1901) (Lanzer 1996; Santos 2003).

Although in Argentina previous records from the literature mentioned the presence of the genus *Gundlachia*, they are in fact related to *Uncancylus concentricus* (d'Orbigny 1835) and *Hebetancylus moricandi* (Pilsbry 1924) in northeast Argentina and Patagonian regions. *Uncancylus concentricus*, under the synonyms *Gundlachia nordenskioldi* Pilsbry, 1924, *G. concentrica* and *G. concentrica bonariense* Strobel, 1874, were described from Corrientes Province (Hylton-Scott 1953). *Gundlachia moricandi* is a synonym of *H. moricandi* and together with *U. concentricus* has a wide distribution from northern Argentina (Misiones, Corrientes, Entre Ríos, Formosa, Chaco, Salta and Santiago del Estero), to Santa Fe, Buenos Aires and Northern Patagonia (Castellanos 1982; Castellanos and Landoni 1995; Rumi *et al.* 2008).




FIGURE 1. *Gundlachia radiata*. Drawings of shell (Mol UERJ 7841). (A) Dorsal view, (B) apical microsculpture with irregular punctuations and (C) left lateral view Scale = 1 mm.

Considering the statement above, the first record of the genus *Gundlachia* for Argentina is from Lanzer (1996) who cited *G. ticaga* to Misiones and Corrientes Province. Gútierrez Gregoric *et al.* (2006) confirmed the record of *G. ticaga* for Misiones province. Recently, Rumi *et al.* (2008) also mentioned *G. ticaga* with a “continuous restricted distribution” in Argentina but without giving the exact distributional area where it was found.




FIGURE 2. *Gundlachia radiata*. (A) Distribution map in the Americas, indicating the previous known locality (blue circles) and the new records (red circles).

Concerning *G. radiata* its current distribution ranges from southern United States to north and northern South America, with references to Texas (Basch 1963), Georgia and Florida (Basch 1963; Hubendick 1967); Central America: Antigua (*apud* Lanzer 1996), Bahamas (Hubendick 1967), Barbados (Schomburgk 1848), Costa Rica (Hubendick 1967), Cuba (Aguayo 1938, Hubendick, 1967), Dominican Republic (Malek 1985), Grenada (Harrison 1983), Guadeloupe (Hubendick 1967, Pointier 1976), Guatemala (Walker 1917, Goodrich and van der Schalie 1937, Hubendick 1967), Honduras (Pfeiffer 1858), Jamaica (Walker 1921, Hubendick 1967), Martinique (Aguayo 1966), México (Hubendick 1967), Nicaragua (Tate 1870, Hubendick 1967, Lanzer 1996, Thompson 2008), Panama (Zetek 1918), Puerto Rico (Harry and Hubendick 1963, Hubendick 1967), Saint Lucia (Harrison 1983), Saint Vincent (Harrison 1983) and Trinidad (Smith 1896);

South America: Colombia (Hubendick 1967), Venezuela (Hubendick 1967) and in Brazil, it is found at north and northern regions (Irmeler 1975; Lanzer 1996; Santos 2003) and recently the species was found at the central-western Brazilian region (Thiengo *et al.* 2005), extending its southernmost geographic range (Figure 2).

Lack of studies concerning to occurrence, systematic and distribution of Ancyliidae, particularly in Argentinean northwestern region (NOA), has led to field work in different NOA localities.

Field work was carried out on May 2009 in small and temporary water bodies in Jujuy province (Argentina). Field works were conducted under appropriate legal authorization. These water bodies are originated by precipitation. Specimens of *G. radiata* were found with both shells forms, with and without septum.

One of the water bodies was located on the edge of the

Route 34 on the road to Calilegua (23°33'43" S, 64°23'33" W) (Figure 3A). The second one was located in Zapla (24°13'10.2" S, 65°06'3.9" W) within an eucalyptus forest (Figure 3B). The specimens were found on sand banks surrounded by vegetation. The material was obtained through the examination of riparian and floating vegetation, with the help of fine mesh sieves. Most specimens were found attached to aquatic vegetation, branches and leaves of eucalyptus and grasses obscured by aging.



FIGURE 3. Temporary ponds inhabited by *Gundlachia radiata* in the province of Jujuy, Argentina. (A) Edge of the Route 34 on the road to Calilegua, (B) Zapla. Photos: Eduardo Dominguez.

The material was analyzed in the Malacological Laboratory of the Miguel Lillo Foundation (UNT-Tucumán). The identification of the specimens was conducted following Santos (2003) using morphological and morphometric characters (shell shape and size, teleoconch and protoconch microsculpture, mantle color pattern and pattern of muscular scars). Out of 260 specimens, 117 presented septate shells on different stages of development (Figure 4). The shell shape was elliptical with periostracum pale color. Left side of shell is more projected than right one, apex irregularly punctuate (Figure 1) similar to Hubendick's (1967) description. The average measurements obtained (mm) from the shells and their ranges of values (maximum-minimum, standard deviation) were, 7.56 (10.3-6.5, 1.54) length, 3.79 (5.9-3.0, 1.0) width and 2.08 (2.6-1.2, 0.51) height. The adhesive area is well marked and elongated and the anterior right muscle scar is more or less elliptical with a slight curvature

in the inner portion (Figure 5A). The mantle pigmentation is scarce, concentrated in the anterior side (Figure 5A). Almost 50% of the collected specimens of *G. radiata* were septated ancyliiform shells in shape (Figure 4 and 5B), reflecting the adverse habitat conditions (temporary ponds). The physical-chemical parameters (pH 5 and low conductivity 36.5 $\mu\text{S}/\text{cm}$) registered in each temporary ponds support, even more, the idea that the development of the septum is in response to unfavorable environmental conditions. The presence of the septum would let keeping the water in the body, thereby, increasing the possibilities of survival of the species (Aguayo 1946; Hylton-Scott 1953; Richardot 1974; Santos 2003).

This record is the southernmost of *G. radiata* to South America and the first to Argentinean northwestern region, extending the geographic distribution of the species and increasing the list of freshwater snails to Argentina from six (Rumi et al. 2008) to seven (Figure 2).

The specimens were deposited at the Malacological Collection of the Miguel Lillo Institute (IML 15291, IML 15292 A, IML 15298 A, IML 15299), Tucumán province, Argentina. Specimens used to soft parts illustration were inserted into the Malacological Collection of the Universidade do Estado do Rio de Janeiro (Mol. UERJ 7841).




FIGURE 4. *Gundlachia radiata*. Drawings of shells (IML 15299) with septum on different stages of development.




FIGURE 5. *Gundlachia radiata*. Drawings of animal. (A) mantle color pattern and muscular scars (Mol. UERJ 7841); (B) ancyliiform form in left lateral view showing mantle modification (IML 15298 A). Scale = 1 mm.

ACKNOWLEDGMENTS: We thank the Funding Agency of the Scientific and Technological Promotion of Argentina (FONCyT-AGENCIA), Project BID 1728/ OC-AR PICT 528, for supporting field works; C. Molineri and E. Dominguez for assistance in the field; Coordenação de Pessoal de Ensino Superior (Capes, Brazil) for L.E.M. Lacerda Master scholarship; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil) for S.B. Santos research grant (476682/2004-5 and Protax 562291/2010-5). We specially thank to Gabriela Cuezco and Eugenia Salas for the personal comments.

LITERATURE CITED

- Aguayo, C.G. 1938. Los moluscos fluviatilis cubanos. *Memorias de la Sociedad Cubana de Historia Natural* 12(3): 203-242.
- Aguayo, C.G. 1946. *Gundlachia* y *Hebetancylus* en Cuba. *Revista de la Sociedad Malacologica "Carlos de La Torre"* 4: 55-62.
- Aguayo, C.G. 1966. Una lista de los moluscos terrestres y fluviatilis de Puerto Rico. *Stahli* 5: 89-105.
- Basch, P.F. 1963. A review of the recent freshwater limpet snails of North America. *Bulletin of the Museum of Comparative Zoology at Harvard College* 129: 401-461.
- Castellanos, Z.A. 1982. Estado actual de Ancyliidae Neotropicales. *Neotropica* 28: 101-102.
- Castellanos, Z.A. and N. Landoni. 1995. Mollusca Pelecypoda y Gastropoda; p. 759-801. In E.C. Lopretto and G.T. Tell (ed.). *Ecosistemas de Aguas Continentales. Metodologías de para su estudio*. Tomo II. La Plata: Ediciones Sur.
- Goodrick, C. and H. Van Der Schalie. 1937. Mollusca of Péten and North Alta Vera Paz, Guatemala. *Miscellaneous Publications, Museum of Zoology, University of Michigan* 34: 1-50.
- Gutiérrez G., D.E., V. Nuñez, A. Rumi and M.A. Roche. 2006. Freshwater gastropods from del Plata Basin, Argentina. Checklist and new locality records. *Comunicaciones de la Sociedad Malacológica del Uruguay* 9(89): 51-60.
- Harrison, A.D. 1983. Identity of *Ferrissia irrorata* and *Gundlachia radiata*, Guilding's species of Ancyliidae from St. Vincent. *Wl. Archiv für Molluskenkunde* 113: 7-15.
- Harry, H.W. and B. Hubendick. 1964. The freshwater pulmonate Mollusca of Puerto Rico. *Göteborgs Kungl Vetenskap- Och Vitterhets-Samhallets Handlinger* 9: 60-72.
- Hubendick, B. 1967. Studies on Ancyliidae. The Australian, Pacific and Neotropical formgroups. *Göteborgs Kungl Vetenskap-Och Vitterhets-Samhallets Handlinger Zoologie* 1: 1-52.
- Hylton-Scott, M.I. 1953. Notas sobre morfología de *Gundlachia* Pfr. (Ancyliidae). *Physis* 20: 467-473.
- Irmiler, U. 1975. Ecological studies of aquatic soil invertebrates in three inundation forests of Central-Amazonia. *Amazoniana* 5: 337-409.
- Lanzer, R.M. 1996. Ancyliidae (Gastropoda: Basommatophora) na América do Sul: sistemática e distribuição. *Revista Brasileira de Zoologia* 13(1): 175-210.
- Malek, E.A. 1985. Freshwater and terrestrial snails of Saint Lucia, West Indies. *Nautilus* 100(4): 143-147.
- Pfeiffer, L. 1858. Beschreibung einer neuen *Gundlachia*. *Malakozoologische Blätter* 5: 196-198.
- Pointier, J.P. 1976. Répartition locale et biogeography des mollusques dulçaquicoles de la Guadeloupe (Antilles françaises). *Malacological Review* 9: 85-103.
- Richardot, M. 1974. La forme septifere de *Ferrissia wautieri*: forme de resistance a la secheresse. *Haliotis* 4: 135-139.
- Rumi, A., Gutiérrez G., D.E., V. Nuñez and G. Darrigran. 2008. Malacología Latinoamérica. Moluscos de agua dulce de Argentina. *Revista de Biología Tropical* 56(1): 77-111.
- Santos, S.B. 2003. Estado atual do conhecimento dos ancilídeos na América do Sul. (Mollusca, Gastropoda, Pulmonata, Basommatophora). *Revista de Biología Tropical* 50 (Suppl. 3): 191-224.
- Schomburgk, R.H. 1848. A list of molluscos animal or shells found at Barbados and among the West Indians islands generally; p. 659-664. In R.H. Schomburgk (ed.). *The history of Barbados*. London: Longman, Brown, Green and Longmans, XXI + 722 p.
- Smith, E.A. 1896. A list of land and freshwater Mollusca of Trinidad. *Journal of Conchology* 8: 231-245.
- Tate, R. 1870. On the land and fresh-water Mollusca of Nicaragua. *American Journal of Conchology* 5: 151-162.
- Thiengo, S.C., S.B. Santos and M.A. Fernandez. 2005. Malacofauna límnic da área de influência do lago da usina hidrelétrica de Serra da Mesa, Goiás, Brasil. I. Estudo qualitativo. *Revista Brasileira de Zoologia* 22: 867-874.
- Thompson, F.G. 2008. *An annotated checklist and bibliography of the land and freshwater snails of Mexico and South America*. Gainesville: University of Florida, 903 p.
- Zetek, J. 1918. *Los moluscos de la Republica del Panama*. Panama: Imprenta Nacional. 80 p.
- Walker, B. 1917. A new *Gundlachia* from Guatemala. *Nautilus* 31(2): 51-53.
- Walker, B. 1921. *Ancylus obscurus* Haldeman and species referred to it. *Nautilus* 34(3): 73-76.

RECEIVED: May 2010

LAST REVISED: March 2011

ACCEPTED: April 2011

PUBLISHED ONLINE: May 2011

EDITORIAL RESPONSIBILITY: Inga Ludmila Veitenheimer Mendes