

Reptilia, Squamata, Gymnophthalmidae, *Acratosaura mentalis* (Amaral 1933): Distribution extension and geographic distribution map

Crizanto Brito De-Carvalho^{1*}, Francis Luiz Santos Caldas¹, Daniel Oliveira Santana³, Marcus Vinicius Noronha², Evellyn Borges de Freitas¹, Renato Gomes Faria^{1,2} and Rafael Alves dos Santos¹

1 Universidade Federal de Sergipe, Núcleo de Pós-Graduação em Ecologia e Conservação. CEP 49000-000. São Cristóvão, SE, Brazil.

2 Universidade Federal de Sergipe, Departamento de Biologia, Laboratório de Herpetologia. CEP 49000-000, São Cristóvão, SE, Brazil.

3 Universidade Federal de Sergipe, Núcleo de Pós-Graduação em Ecologia e Conservação. CEP 49000-000, São Cristóvão, SE, Brazil.

* Corresponding author. E-mail: crizantobrito@gmail.com

ABSTRACT: This article provides new records to *Acratosaura mentalis* of two new localities from the state of Sergipe, northeast Brazil. These records extend the geographical distribution of specie for municipalities from Poço Redondo and Itabaiana. Possibly, these points are the first records of *A. mentalis* for the state of Sergipe.

The Gymnophthalmid lizards are small and medium-sized, distributed from southern Mexico to Argentina, the Caribbean, on some continental shelf islands of Central and South America east of the Andes (Pellegrino *et al.* 2001; Vitt and Caldwell 2009). Currently in Brazil there are 82 species in 32 genera (Bérnils 2010), some of which are restricted to the northeastern region (*e.g.* Rodrigues *et al.* 2007; Rodrigues and Santos 2008; Rodrigues *et al.* 2009). The *Acratosaura* genus was created to accommodate *Colobosaura mentalis* Amaral, 1933 (Rodrigues *et al.* 2007), and now also includes *Acratosaura spinosa* Rodrigues, Cassimiro, Freitas and Silva 2009.

Acratosaura mentalis (Figure 1) was described by Amaral in 1933 from municipality of Vila Nova, now Senhor do Bonfim, state of Bahia (BA), Brazil. The second record of its geographical distribution was made by Rodrigues (1986) in the municipality of Cabaceiras, state of Paraíba (PB), while Queiroz and Lema (1996) recorded the species in Brejo da Madre de Deus, state of Pernambuco (PE). Since then, records have been collected in new surveys at other localities (Figure 2).

Herein, we expand the geographic distribution of *Acratosaura mentalis* by presenting two new localities from the state of Sergipe: Serra da Guia (09°58'52" S, 37°52'05" W; 727 above sea level) in the municipality of Poço Redondo, and the Parque Nacional Serra de Itabaiana (10°44'56" S, 37°20'31" W; 240 m above sea level) in municipality of Areia Branca (Figure 2). The latter site presents areas of semideciduous lowland forest interspersed with more open habitat (Vicente *et al.* 2005), and represents the transition zone between the Atlantic Forest and the semi-arid Caatinga, in which species from both biomes coexist (Carvalho and Vilar 2005).

The Serra da Guia site is known locally as *brejos de altitude*, and is characterized by humid forests on hilltops and plateaus over 500 m above sea level (Prado 2005).



FIGURE 1. *Acratosaura mentalis* (C720), Parque Nacional Serra de Itabaiana, state of Sergipe, Brazil. Photo by C.B. De-Carvalho.

These areas appear to be relicts of the Atlantic Forest, characterized by high biodiversity, isolated within the more arid Caatinga vegetation (Barbosa *et al.* 2004).

Our records extend the geographical distribution of *Acratosaura mentalis* ca. 270 km for Serra da Guia and 320 km for Serra de Itabaiana in relation to the type locality at Senhor do Bonfim, state of Bahia. The nearest point is located in Xingó Power Plant (about 50 km), where we do not know if the specimen was collected in the state of Alagoas or Sergipe. Possibly, these points presented here are the first records of *A. mentalis* for the state of Sergipe.

Identification of the specimens was based in Amaral (1933), Queiroz and Lema (1996) and Rodrigues *et al.* (2009). Voucher specimens were deposited at the Herpetological Collection of the Universidade Federal de Sergipe, Brazil (C296 and C720). Collecting permit (# 10504-1) was granted by Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis - IBAMA.




FIGURE 2. Geographic distribution of *Acratosaura mentalis*: 1. Santana do Pirapama (Rodrigues *et al.* 2009); 2. Diamantina (Rodrigues *et al.* 2009); 3. Turmalina (Rodrigues *et al.* 2009); 4. Cristália (Rodrigues *et al.* 2009); 5. Grão Mogol (Rodrigues *et al.* 2009); 6. Parque Nacional do Peruaçu (Rodrigues *et al.* 2009); 7. Maracás (Rodrigues *et al.* 2009); 8. Mucugê (MZUSP 74223); 9. Santo Inácio (MZUSP 79743); 10. Senhor do Bonfim (Type-Locality, MZUSP 40084-5) 11. Sento Sé (Santos *et al.* 2008); 12. Parque Nacional Serra de Itabaiana (New Record); 13. Serra da Guia (New Record); 14. UHE Xingó (MZUSP 80182-3); 15. Delmiro Gouveia (Rodrigues *et al.* 2009); 16. Garanhuns (MZUSP 65675); 17. Parque Nacional do Catimbau (Rodrigues and Santos 2008); 18. Brejo da Madre de Deus (Queiroz and Lema 1996); 19. São José dos Cordeiros and Sumé (Delfim and Freire 2007); 20. Cabaceiras (Rodrigues 1986); 21. Cacimba de Dentro (Arzabe *et al.* 2005), 22. Parque Estadual Pedra da Boca (Arzabe *et al.* 2005) and 23. Tenente Laurentino Cruz (Gogliath *et al.* 2010). States Abbreviations: AL = Alagoas, BA = Bahia, MG = Minas Gerais, PE = Pernambuco, PB = Paraíba, RN = Rio Grande do Norte and Sergipe = SE.

ACKNOWLEDGMENTS: We thank Hussam Zaher and Carolina Castro-Mello for geographic data about specimens housed in the herpetological collection of Museu de Zoologia da USP, Marcelo Brito for comments on the map, Steve Ferrari, Bruno Duarte and Luiz Gustavo Gomes Borges for their revision of the text and two anonymous referees for critically reviewing the manuscript. The Secretaria de Meio Ambiente e Recursos Hídricos - SEMARH/SE and Universidade Federal de Sergipe for logistic and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for fellowships.

LITERATURE CITED

Amaral, A. 1933. Estudos sobre lacertílios neotrópicos. *Memórias do Instituto Butantan* 7: 1-74.

Arzabe, C., G. Skuk, G.G. Santana, F.R. Delfim, Y.C.C. Lima and S.H.F. Abrantes. 2005. Herpetofauna da área do Curimataú, Paraíba; p. 264-280. In F.S. Araújo, M.J.N. Rodal and M.R.V. Barbosa (ed.). *Análise das variações da biodiversidade do Bioma Caatinga: suporte a estratégias regionais de conservação*. Brasília: Ministério do Meio Ambiente.

Barbosa, M.R.V., M.F. Agra, E.V.S.B. Sampaio, J.P. Cunha and L.A. Andrade. 2004. Diversidade florística da Mata de Pau Ferro, Areia, Paraíba; p. 111-122 In K.C. Porto, J.J.P. Cabral and M. Tabarelli. (ed.). *Brejos de altitude em Pernambuco e Paraíba: História natural, ecologia e conservação*. Brasília: Ministério do Meio Ambiente.

Bérnils, R.S. 2010. *Brazilian reptiles - List of species*. Electronic Database accessible at <http://www.sbherpetologia.org.br>. Brazilian Society of Herpetology, Brazil. Captured on 06 June 2010.

Carvalho, C.M. and J.C. Vilar. 2005. Levantamento da biota do Parque Nacional Serra de Itabaiana. *Biologia Geral e Experimental* 30(1): 9-14.

Delfim, F.R. and E.M.X. Freire. 2007. Os lagartos Gimnophthalmídeos (Squamata: Gymnophthalmidae) do Cariri Paraibano e do Seridó do Rio Grande do Norte, Nordeste do Brasil: Considerações acerca da distribuição geográfica e ecologia. *Oecologia Brasiliensis* 11(3): 365-382.

Gogliath, M., L.B. Ribeiro and E.M.X. Freire. 2010. *Acratosaura mentalis* (NCN) Geographic Distribution. *Herpetological Review* 41(1): 105-106.

Pellegrino, K.C.M., M.T. Rodrigues, Y. Yonenaga-Yassuda and J.W. Sites Jr. 2001. A molecular perspective on the evolution of microteiid lizards (Squamata, Gymnophthalmidae), and a new classification for the family. *Biological Journal of the Linnean Society* 74(3): 315-338.

Rodrigues, M.T. 1986. Uma nova espécie do gênero *Phyllopezus* de Cabaceiras: Paraíba: Brasil; com comentários sobre a fauna de lagartos da área (Sauria Gekkonidae). *Papéis Avulsos de Zoologia* 36(20): 237-250.

Rodrigues, M.T. 2005. Herpetofauna da Caatinga; p. 181-236 In I.R. Leal, M. Tabarelli and J.M.C. Silva (ed.). *Ecologia e conservação da Caatinga*. Recife: Editora Universitária, Universidade Federal de Pernambuco.

- Rodrigues, M.T., K.C.M. Pellegrino, M. Dixo, V.K. Verdade, D. Pavan, A.J.S. Argôlo and J.W. Sites Jr. 2007. A new genus of microteiid lizard from the Atlantic Forests of state of Bahia, Brazil, with a new generic name for *Colobosaura mentalis*, and a discussion of relationships among the Heterodactylini (Squamata, Gymnophthalmidae). *American Museum Novitates* 3565: 1-27.
- Rodrigues, M.T. and E.M. Santos. 2008. A new genus and species of eyelidless and limb reduced gymnophthalmidae lizard from northeastern Brazil. (Squamata Gymnophthalmidae). *Zootaxa* 1873: 50-60.
- Rodrigues, M.T., J. Cassimiro, M.A. Freitas and T.F.S. Silva. 2009. A new microteiid lizard of genus *Acratosaura* (Squamata: Gymnophthalmidae) from Serra do Sincorá, State of Bahia, Brasil. *Zootaxa* 2013: 17-19.
- Santos, F.J.M., A.P. Penã and V.L.F. Luz. 2008. Considerações biogeográficas sobre a herpetofauna do Submédio e da Foz do Rio São Francisco, Brasil. *Estudos* 35 (1/2): 59-78.
- Prado, D. 2005. As caatingas da América do Sul; p. 3-73 In I.R. Leal, M. Tabarelli and J.M.C. Silva (ed.). *Ecologia e conservação da Caatinga*. Recife: Editora Universitária, Universidade Federal de Pernambuco.
- Queiroz, A.N. and T. Lema. 1996. Novo registro de *Colobosaura mentalis* Amaral, 1933, (Sauria: Teiidae) para o nordeste do Brasil. *Biociências* 4(1):87-90.
- Vicente, A., A.S. Ribeiro, E.A. Santos and C.R.P. Franco. 2005. Levantamento botânico. *Biologia Geral e Experimental* 300: 15-37.
- Vitt, L.J., J. Caldwell. 2009. *Herpetology: an introductory biology of amphibians and reptiles*. San Diego: Academic Press. 697p.

RECEIVED: November 2009

REVISED: June 2010

ACCEPTED: July 2010

PUBLISHED ONLINE: September 2010

EDITORIAL RESPONSIBILITY: Julio Cesar de Moura Leite