

# Amphibia, Anura, Leptodactylidae, *Leptodactylus paraensis* Heyer, 2005: Distribution extension, new state record, and geographic distribution map

Domingos J. Rodrigues<sup>1,2,3\*</sup>, Marcelo M. Lima<sup>4</sup>, Carla L. Velásquez<sup>3</sup> and Tamí Mott<sup>2,3,5</sup>

- 1 Universidade Federal de Mato Grosso, Instituto de Ciências Naturais Humanas e Sociais, Núcleo de Estudo da Biodiversidade da Amazônia Matogrossense. Avenida Alexandre Ferronato, 1200, Setor Industrial. CEP 78557-267. Sinop, MT, Brazil
  - 2 Instituto Nacional de Ciência e Tecnologia de Estudos Integrados da Biodiversidade Amazônica – INCT-CENBAM/CNPq/MCT
  - 3 Universidade Federal de Mato Grosso, Instituto de Biociências, Mestrado em Ecologia e Conservação da Biodiversidade. Avenida Fernando Corrêa da Costa, 2367, Boa Esperança. CEP 78060-900. Cuiabá, MT, Brazil
  - 4 Instituto Nacional de Pesquisas da Amazônia, Coordenação de Pesquisas em Ecologia. Avenida André Araújo 2936. CEP 69011-970. Manaus AM, Brazil
  - 5 Universidade Federal do Mato Grosso, Instituto de Biociências, Departamento de Biologia e Zoologia. CEP 78060-900. Cuiabá, MT, Brazil
- \*Corresponding author. E-mail: [djmingo23@gmail.com](mailto:djmingo23@gmail.com)


**ABSTRACT:** This note reports the presence of *Leptodactylus paraensis* at the municipality of Cláudia, state of Mato Grosso, central Brazil. This is the first recorded for this species in Mato Grosso.

The *Leptodactylus pentadactylus* species group is composed of 19 species of medium to large-sized frogs (Frost 2010). *Leptodactylus paraensis* Heyer 2005 is a large sized species that belongs to this group and is known only from its type-locality in Serra de Kukoinhokren (07°46' S, 51°57' W), state of Pará, Brazil (Heyer 2005). However, this species occurs also in Tapajós National Forest (A.P. Lima, pers. comm.), state of Pará, Brazil, approximately 650 km southeastern of type-locality. These localities are characterized as areas of closed canopy Amazonian rain forests (Heyer 2005). Here we report a new state record for *L. paraensis* extending its distribution over 500 km south in Amazonian forest in Brazil.

During field work from November to December 2009 and January 2010, we observed and collected several individuals of *L. paraensis* in native Amazonian forest with selective logging, in Continental Farm (two localities: module 1: 11°35'9.61" S, 55°16'10.86" W and module 2: 11°24'44.21" S, 55°19'25.20" W), and Iracema Farm (module 3: 11°38'20.40" S, 55° 5'25.43" W), all areas within the municipality of Cláudia, north of state of Mato Grosso, Brazil. The region constitutes a transitional area between the Amazonian moist forest and the Cerrado vegetation, forming the dry forest of Mato Grosso. The relief in the region is flat and gently dissected in tabular form, with elevations ranging from 400-500 m. The predominant soil in the area is Red-Yellow Dystrophic. The climate is tropical rainy with short period of drought, type Am (Köppen). The temperature annual average is 24 °C and rainfall around 2,000 mm/year. The rainy season occurs from December to February and the dry season from June to September.

Data were collected in December/2009 and, January and February/2010 in three modules with 32 plots (module 1 and 2 with 12 plots each, and module 3 with eight plots) systematically distributed over a 5 km<sup>2</sup> grid formed by 5 km long trails (Figure 1; more information is available from <http://ppbio.inpa.gov.br>). Each plot was at

least 1 km distant from any other. Plots were 250 m long and positioned to follow altitudinal contour lines, and thus minimized altitudinal and soil variation within each plot (Magnusson *et al.*, 2005). All plots were at least 600 m from the edge of the forest. We observed 22 individuals (two in module 1; seven in module 2 and 13 in module



**FIGURE 1.** Known distribution of *Leptodactylus paraensis* in Brazil (type-locality – star) and the updated current distribution: (square and triangle to module 1 and 2 at Continental farm respectively; triangle: module circle: module 3 at Iracema Farm. All these new records are located at the Municipality of Cláudia, state of Mato Grosso, Brazil).


3), but only two were collected (males 148.1 and 142.0 mm snout-vent length, respectively) by MML and DJR (collection permit # 10174-1, Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis – IBAMA). The two collected specimens and the other individuals only observed were found near a burrow (depth approximately 15 cm) in native forest with selective logging (Figure 2). *Leptodactylus paraensis* is abundant in native forests (Heyer, 2005) and forests that have been subjected to selective logging in Brazilian Amazon (A.P. Lima, pers. comm.), including the studied areas, where it was found in 11 of the 32 plots sampled, with four plots recording more of three individuals. Voucher specimens were deposited in the Herpetological Collection of the Universidade Federal de Mato Grosso, Campus Universitário of Sinop, under numbers UFMT 051 and UFMT 052.



**FIGURE 2.** Adult male of *Leptodactylus paraensis* in front of burrow (A) in Iracema Farm, municipality of Cláudia, state of Mato Grosso, Brazil.

Specimens of *Leptodactylus paraensis* herein reported exhibit all diagnostic characteristics described by Heyer (2005) including: lip pattern often with 1 or 2 dark elongate triangular marks approaching or entering the lower eye. Dorsal pattern often uniform light or dark; or well developed dark interorbital band/chevron and 2 moderate to large equally intense dark chevrons, second chevron in sacral region, chevrons confluent or not; or more than 2 dark, broad transverse bands of equal intensity in addition to interorbital band, confluent laterally or not. Belly pattern often mottled or uniform dark or dark with small light vermiculations. Posterior thigh pattern usually dark with small distinct light vermiculations or spots. Snout nearly rounded in dorsal and profile views, ventral region smooth, small tubercles in the dorsum, chest finely mottled, and anterior belly with large and light vermiculations on a slightly darker background, posterior belly with faint but distinct labyrinthine pattern.

*Leptodactylus paraensis* (Figure 3A) can be confused with *L. labyrinthicus* (Figure 3B) and *L. pentadactylus* (Figure 3C) in the surveyed region due to their morphological similarities. Accordingly to Heyer (2005), there is no consistent morphological feature that completely distinguishes *L. paraensis* from *L. labyrinthicus*. Nevertheless, the vocalization and the ecological requirements definitely differentiate them; *L. labyrinthicus* occurs in open formation habitats, while *L. paraensis* occurs in closed forest formations. Data of



**FIGURE 3.** Ventral, dorsal and head view of *Leptodactylus pentadactylus* species group. A) *L. paraensis*, B) *L. labyrinthicus*, C) *L. pentadactylus*. Photo A, B and C by D.J. Rodrigues

habitat usage for *L. paraensis* indicate that the species is limited to primary rain forests as *L. pentadactylus* (Heyer, 2005; Lima et al., 2006). A difference found in the present study is that *L. pentadactylus* was found always near to stream as showed by Hero and Galatti (1990), Lima et al. (2006); Menin et al. (2009), while *L. paraensis* was found in plateau, distant at least 500 m from the nearest stream. Morphological characters also distinguish *L. paraensis* from *L. pentadactylus* such as the extension of their dorsolateral folds. In *L. paraensis* these folds are interrupted and extend no further from the eye; dorsolateral folds of *L. pentadactylus* are either continuous or, if interrupted, extend beyond the sacrum (Hero and Galatti, 1990; Lima et al. 2006). *Leptodactylus pentadactylus* has a continuous line extending from eyes to tip of snout while *L. paraensis* present only part of this line (Hero and Galatti, 1990; Heyer, 2005).

*Leptodactylus paraensis* is classified as Least Concern in the IUCN Red List, but the biology and the distribution of this species is little-known, as the majority of newly

described species for which more information is still needed (IUCN 2010). The record of *L. paraensis* from municipality of Cláudia (modules 1, 2 and 3, Figure 1) represents the first record of this species in the state of Mato Grosso, expanding its distribution ca. 550 km south-southwest of type locality (Figure 1).

**ACKNOWLEDGMENTS:** Everton J. Almeida, Evandro F. dos Santos, and Ricardo Machiner for field work assistance. A anonymous reviewer for valuable comments and suggestions on manuscript; Continental and Iracema Farm for logistic support and Ministério da Ciência e Tecnologia / Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) CNPq nº 569382/2008-4 and 501408/2009-6 for financial support. Fellowships from CNPq to DJR and CAPES to CLV. This is publication 02 in the NEBAM technical series.

#### LITERATURE CITED

Frost, D.R. 2010. *Amphibian Species of the World: An online reference. Version 5.3*. American Museum of Natural History, New York. Electronic database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. Captured on 10 March 2010.

Hero, J-M. and U. Galatti. 1990. Characteristics distinguishing *Leptodactylus pentadactylus* and *L. knudseni* in the Central Amazon Rainforest. *Journal of Herpetology* 24: 226–228.

Heyer, W.R. 2005. Variaton and taxonomic clarification of the large species of the *Leptodactylus pentadactylus* species group (Amphibia: Leptodactylidae) from middle America, Northern South America and Amazonia. *Arquivos de Zoologia* 37(3): 269-348.

IUCN. 2010. *IUCN Red List of Threatened Species. Version 2009.2*. Eletronic database accessible at [www.iucnredlist.org](http://www.iucnredlist.org). Captured on 13 July of 2010.

Lima, A.P., W.E. Magnusson, M. Menin, L.K. Erdtmann, D.J. Rodrigues, C. Keller and W. Hödl. 2006. *Guia de Sapos da Reserva Adolpho Ducke, Amazônia Central (Guide to the frogs of Reserva Adolpho Ducke, Central Amazonia)*. Manaus: Áttema Design Editorial. 168 p.

Magnusson, W.E. A.P. Lima, R.C. Luizão, F. Luizão, F.R.C. Costa, C.V. Castilho and V.F. kinupp. 2005. RAPELD: uma modificação do método de Gentry para inventários de biodiversidade em sítios para pesquisa ecológica de longa duração. *Biota Neotropica* 5(2): <http://www.biotaneotropica.org.br/v5n2/pt/abstract?point-ofview+bn01005022005>.

Menin, M., A.P. Lima and D.J. Rodrigues. 2010. The tadpole of *Leptodactylus pentadactylus* (Anura: Leptodactylidae) from Central Amazonia. *Zootaxa* 2508: 65-68.

RECEIVED: May 2010

REVISED: July 2010

ACCEPTED: August 2010

PUBLISHED ONLINE: September 2010

EDITORIAL RESPONSIBILITY: Marcelo Nogueira de Carvalho Kokubum