

New distribution Record of *Phrynocephalus mystaceus* Pallas 1776 (Sauria: Agamidae) from Iran

Farahnaz Molavi¹, Tooba Mohammadian-kalat^{2*}, Bahram Hasanzade Kiabi³, Haji Gholi Kami⁴ and Morteza Yazdanpanahi³

- 1 Islamic Azad University, Mashhad Branch, Faculty of science, Department of Biology, Mashhad, Iran.
 - 2 Ferdowsi University of Mashhad, Faculty of Sciences, Department of Biology, Mashhad, Iran.
 - 3 Shahid Beheshti University, Faculty of Science, Department of Biology, Tehran, Iran.
 - 4 University of Agricultural Sciences and Natural Resources, Faculty of Science, Department of Biology, Gorgan, Iran.
- * Corresponding author. E-mail: tmohammadian@gmail.com

ABSTRACT: The Toad-headed agama, *Phrynocephalus mystaceus* Pallas 1776, was described in 1999 from eastern Khorasan by Anderson. Seven specimen of The Toad-headed agama were collected in Khar Turan National Park during fieldwork from June 2008 to June 2009. The new locality of the species is situated about 900 km west of the type locality. This record indicates a wider distribution of *Phrynocephalus mystaceus* on the Iranian plateau than previously thought. Information on morphological characters and habitat is presented.

Toad-headed agamas, *Phrynocephalus* Kaup, 1825, are distributed from southern Europe and southwest Asia (including the Arabian Peninsula and northern India) through central and Middle Asia to East Asia (northern China and Mongolia). This further demonstrates that the taxonomic diversity of *Phrynocephalus* in Iran is ambiguous. Currently there are approximately 40 species of Toad-headed Agamas of the genus *Phrynocephalus*, seven of which have been recorded from Iran (Anderson, 1999). However, Golubev & Sattorov (1992) recorded ten species of *Phrynocephalus* from Southwest and middle Asia. The best diagnostic character of *Phrynocephalus mystaceus* is a large fringed cutaneous fold at angle of mouth. Ananjeva (1986) refers this species to the monotypic genus *Megalochilus*, but Golubev argue that the differences are too slight to warrant a separate genus (Golubev and Sattorov, 1992). In this study, we report new

data on morphological characters and sexual dimorphism with a range extension for *P. mystaceus* in Iran.

Seven specimens of *P. mystaceus* were collected in June 2008–9 from Khar Turan National Park in Semnam province (Figure 1). This locality is located in a sandy area of eastern part of Ahmad Abad village (35°48' N, 56°42' E) 1000 m above sea level (Figure 2). *Haloxyylon* sp. and *Tamarix* sp. are dominant plant species of this locality.

Specimens were captured alive for study of their behavior in the laboratory, and later were preserved in 10% formalin. Characters used in this study are presented in Table 1. Morphometric characters were measured with vernier caliper (accuracy 0.02 mm) locality.

Calculation of mean, minimum, maximum, standard deviation; coefficient of variation in both male and female were implemented separately using SPSS Ver. 16.0.1 statistical package (SPSS Inc., Chigaco, IL, USA).



FIGURE 1. Habitat of *Phrynocephalu smystaceus*.




FIGURE 2. *Phrynocephalus mystaceus* type locality, square: previous localities (Anderson 1999), circle: new locality.

In order to investigate Sexual dimorphism, we examined distribution and variances of each character using Kolmogorov-Smirnov and Levene Tests and finally one-way ANOVA was employed to assess differences between male and female characters.

Results of statistical analysis of morphological characters are shown in Table 2. According to Table 2, all of the characters exhibited normal distribution and variances but none of characters could reveal sexual dimorphism.

Color pattern: Dorsum is sandy; with black and white dots and reticulations; a row of larger dark blotches on each side of vertebral Line; vent white; ventral aspect of tail tip black chin and throat with gray reticulation (Figure3).

In this species TL is about equal to the SVL and in largest specimen collected: SVL=97.48 and TL=101 mm. Although Anderson had found a specimen with SVL=122.7 and TL= 135.8 mm in Uzbekistan, the largest specimen that he had collected in Iran was a male with SVL=77.7 and TL=71 mm.

In all specimens studied, there is a greatly enlarged scale (equal to about four adjacent scale) on each side of thorax behind maxilla and occasionally an additional scale or two slightly smaller adjacent to it. This property



FIGURE 3. *Phrynocephalus mystaceus* in natural sandy habitat.

TABLE 1. The quantitative (metric and meristic) measured characters.

CHARACTERS	DEFINITION
SVL.TL	SVL / TL
SVL	Length of snout to vent (from tip of snout to anterior edge of cloacae)
HW	Wide of head (at the widest point of head)
HL	Length of head (from tip of snout to posterior edge of tympanum)
TL	Tail length (from posterior edge of cloacae to tip of tail)
NS	Number of nasal scales
AB	Number of scales around body
BE	Number of scales between eyes in the head width
BLEL	Number of scales between labial and eye (right)
BLER	Number of scales between labial and eye (left)
LLR	Number of lower labial scales (right)
LLL	Number of lower labial scales (left)
SLR	Number of supralabials (right)
FT4L	Total number of lamellae under the left forth toe
FT4R	Total number of lamellae under the right forth toe

TABLE 2. Results of statistical analysis of morphological characters of the *Phrynocephalus mystaceus* from Semnan province of Iran. Mean, Maximum and minimum values for different characters in both male (M) and female (F) and result of Kolmogorov-Smirnov test (K.S), Levene's test (Lev), One-Way Analysis of Variance (CRD) are summarized.

Variable	MEAN		SD		Min		Max		CV%		N		F		
	M	F	M	F	M	F	M	F	M	F	M	F	K.S	Lev	CRD
SVL/TL	0.80	1.01	0.05	0.30	0.93	0.97	1.04	1.03	5.10	2.90	3	4	0.929	0.385	0.487
NS	2.00	2.00	0.00	0.00	2.00	2.00	2.00	2.00	0.00	0.00	3	4	—	—	—
BLEL	3.00	3.00	0.00	0.00	3.00	3.00	3.00	3.00	0.00	0.00	3	4	—	—	—
BLER	3.33	3.00	0.58	0.00	3.00	3.00	4.00	3.00	17.00	0.00	3	4	0.056	0.05	0.285
LLL	9.00	9.50	1.41	1.29	8.00	3.00	10.00	11.90	15.00	13.00	2	4	0.955	0.116	0.247
SLL	15.00	11.75	0.00	4.19	15.00	6.00	15.00	16.00	0.00	35.00	3	4	0.65	0.110	0.665
LLR	8.50	11.75	2.12	3.59	7.00	9.00	10.00	17.00	24.00	30.00	2	4	0.677	0.481	0.317
SLR	15.33	16.25	1.53	2.22	14.00	14.00	17.00	19.00	9.00	13.00	3	4	0.78	0.361	0.569
FT4L	18.33	18.50	1.15	1.73	17.00	17.00	19.00	20.00	8.00	9.00	3	4	0.66	0.05	0.892
FT4R	17.67	19.00	0.85	1.41	17.00	17.00	18.00	20.00	3.000	7.40	3	4	0.93	0.310	0.191
HW	20.38	19.18	3.37	7.49	18.00	13.04	22.76	28.50	16.00	39.00	2	4	0.98	0.138	0.846
HL	24.25	20.20	2.47	6.86	22.50	13.48	26.00	29.00	10.00	33.00	2	4	0.97	0.192	0.483
BE	30.33	31.50	2.08	3.11	28.00	28.00	32.00	35.00	6.80	9.80	3	4	0.99	0.288	0.601
SVL	68.42	68.83	20.72	25.65	47.66	43.24	89.10	97.48	30.00	37.00	3	4	0.91	0.331	0.982
TL	69.07	68.91	17.42	27.26	51.20	42.88	86.00	101.00	25.00	39.00	3	4	0.88	0.179	0.393
AB	117.33	94.75	8.08	60.77	110.00	5.00	126.00	138.00	6.00	64.00	3	4	0.88	0.179	0.393

has reported by Anderson in other specimens of other regions; such scalation is absent in all other species of *Phrynocephalus* examined (all southwest Asian species and several Central Asian forms) (Anderson 1999). Therefore these two characters are added to former morphological key characters of this species.

Fringes on digits (numbered 17–18) are adaptation to its special habitat. Since the whole distribution of this Lizard is sand dunes.

Terentjev and Chernov (1949) recognize two subspecies from entire distribution range. Two subspecies of this species, *P. m. galli* and *P. m. mystaceus* are subject of debate. Anderson had found specimens from Iran appear to be intermediate between two taxa. They agree with *P. m. galli* in color pattern and fringes on the digits and show a *better fit to P. m. mystaceus* in the fold at the corners of mouth dose not reach the neck, and also that the tail is developed. Moreover, keels of breast and belly are said to be strongly developed in *P. m. galli* and less distinct in *P. m. mystaceus* (Anderson 1999). However, Anderson (1999) does not attribute Iranian specimens to any of these subspecies. Golubev and Sattorov (1992) regard *P. mystaceus* as a monotypic species based on study of morphological characters throughout the range. Present work revealed that morphology alone is probably not sufficient for fine resolution of subspecies. Combination of both morphological and molecular methods is recommended to demonstrate if *P. mystaceus* is a monotypic or polytypic species.

LITERATURE CITED

- Anderson, S.C. 1974. Preliminary key to the Turtles, Lizards and amphisbaenians of Iran. *Fieldiana: Zoology* 65(4): 27–44.
- Anderson, S.C. 1999. *The Lizards of Iran*, Ithaca, New York, USA: Society for the Study of Amphibians and Reptiles. 415 pp.
- Golubev, M.L. and T.S. Sattorov. 1992. On the intraspecific structure and extraspecific relations of the ear-folded toad agama *Phrynocephalus mystaceus* (Reptilia, Agamidae). *Vestnik Zoologii* 3: 26–32 [in Russian].
- SPSS, 2007. *SPSS 16.0.1 for Windows*. Chicago: SPSS Inc. Accessible at <http://www.spss.com>.
- Terentjev P.V. and S.A. Chernov. 1949. *Opredelitel Presmykayushchikhsya i Zemnovodnykh* (Guide to Reptile and Amphibians). Moscow and Leningrad: Soviet Science Prees. 340 pp. [In Russian].
- Tuck, Robert G., Jr. 1979. Notes on the Turan Biosphere Reserve herpetofauna, northwestern Iran. *Bulletin of the Maryland Herpetological Society* 15(4): 95–123.
- Welch K.R.G. 1983. *Herpetology of Europe and Southwest Asia: A checklist and Bibliography of the Order Amphisbaenia, Sauria and Serpentes*. Malabar, Florida: Robert E. Krieger Publishing Company. Viii+135 pp.
- Wermuth H. 1967. *Liste der rezenten Amphibien und Reptilien: Agamidae*. *Das Tierreich*. Berlin: Walter de Gruyter. vol. 86: xiv+127 pp.

RECEIVED: October 2013

ACCEPTED: February 2014

PUBLISHED ONLINE: May 2014

EDITORIAL RESPONSIBILITY: Perry Wood