





Notes on the butterfly theridiid *Episinus marignaci* (Lessert, 1933) (Araneae, Theridiidae) from South Africa

Anna S. Dippenaar-Schoeman¹, Stefan H. Foord^{1*}

¹ SARChI Chair in Biodiversity Value and Change and Centre for Invasion Biology, University of Venda, Thohoyandou, South Africa • ASD: dippenaaransie@gmail.com  <https://orcid.org/0000-0003-1532-1379> • SHF: stefan.foord@univen.ac.za  <https://orcid.org/0000-0002-9195-2562>

* Corresponding author

Abstract

We present the first records of the butterfly theridiid *Episinus marignaci* (Lessert, 1933) beyond its type locality, resulting in an extension of its geographic range from Angola to South Africa. Images of live specimens and a distribution map are provided. *Episinus marignaci* is rare locally but has a relatively large geographical distribution in South Africa.

Keywords

Angola, geographical distribution, South African National Survey of Arachnida

Academic editor: Gustavo Silva de Miranda | Received 21 July 2022 | Accepted 8 September 2022 | Published 26 October 2022

Citation: Dippenaar-Schoeman AS, Foord SH (2022) Notes on the butterfly theridiid *Episinus marignaci* (Lessert, 1933) (Araneae, Theridiidae) from South Africa. *Check List* 18 (5): 1177–1180. <https://doi.org/10.15560/18.5.1177>

Introduction

The genus *Episinus* Walckenaer, 1809 is represented by 66 species/subspecies with 10 species and one subspecies listed from Africa but only one species from South Africa (World Spider Catalog 2022). The South African *E. bilineatus* Simon, 1894 is categorized as Data Deficient, as the species was described based on a single juvenile specimen and the type locality was only given as Transvaal.

The *Episinus* specimens sampled during South African National Survey of Arachnida (SANSA) corresponded with the original description and images provided for *E. marignaci* (Lessert 1933) (Fig. 2A, C). The species was originally described as *Episinopsis marignaci* from Santo Amaro in Angola, but Levi and Levi (1962) regarded *Episinopsis* Simon, 1894 to be a synonym of *Episinus*.

Episinus marignaci is commonly known as butterfly spiders. It is a very colourful spider and has been sampled from several localities in South Africa. For the first time, images of live specimens are provided with notes on their behaviour, distribution, and conservation status in South Africa.

Methods

Extensive surveys were undertaken throughout the country during the South African National Survey of Arachnida (SANSA) and large numbers of theridiid specimens were sampled (Dippenaar-Schoeman et al. 2015). Requests for photographs for the SANSA Virtual Museum were also made (Dippenaar-Schoeman et al. 2012), and images of *E. marignaci* were received from

two localities in South Africa. The digital image of the epigyne was taken using the built-in camera of a Leica EZ4D microscope. Species-level determinations were done using Levi and Levi (1962). Voucher specimens of specimens collected are housed in the National Collection of Arachnida (NCA) at the Agricultural Research Council in Pretoria.

Results

Episinus marignaci (Lessert, 1933)

Episinopsis marignaci Lessert 1933: 103, figs. 17–19 (female).

Episinus marignaci—Dippenaar-Schoeman et al. 2013: 121; Dippenaar-Schoeman 2014: 144.

Figures 1, 2

Type locality. Angola, Santo Amaro.

New records. SOUTH AFRICA – **Eastern Cape** • Kei River Mouth; 32.68°S, 028.37°E; 3.III.2007; C. Haddad leg.; 1♀, NCA 2007/1425 • Addo Elephant National Park, Woody Cape; 33.88°S, 025.45°E; 2.II.2013; L. Wiese leg.; 1♀, NCA 2013/1109 • Addo Elephant National Park, Zuurberg; 33.20°S, 025.75°E; L. Wiese leg.; 1♀, NCA 2016/1796 – **KwaZulu-Natal** • iSimangaliso WP, uMkhuze Game Reserve; 27.63°S, 032.25°E; 5.II.2004; X. Combrink leg.; 1♀, NCA 2004/496 • Ndumo Game Reserve; 26.87°S, 032.24°E; 2.X.2004; C. Haddad leg.; 1♀, NCA 2004/919 • Umgeni Valley Nature Reserve; 29.29°S, 030.15°E; 5.x.2010; A.S. Dippenaar leg.; 1♀, NCA 2010/826 • Alverstone near Hillcrest; 29.77°S, 030.73°E; 28.XII.2016; Peter Webb leg.; 1♀, photographs – **Limpopo** • Blouberg Nature Reserve; 22.99°S, 029.04°E; 13.ii.2009; S. Foord leg.; 1♀, NCA 2009/2047 • Little Leigh, Western Soutpansberg; 22.9°S, 029.87°E; 5.III.2009; S. Foord leg.; 1♀, NCA 2009/2403 • Nylsvley Nature Reserve; 24.6°S, 028.67°E; 7.iii.1987; G. Ferreira leg.; 1♀, NCA 87/279 • Potgietersrus/Mokopane; 24.17°S, 028.09°E; 15.iv.2005; A.S. Dippenaar leg.; 1♀, NCA 2005/1051 • Mphaphuli Cycad Reserve; 22.42°S, 030.49°E; 18.III.2016; P. Webb leg.; 1♀, NCA 2016/3578 & photographs.

Identification. Total length of female 3–6 mm. Carapace slightly longer than wide; fawn with two dark, V-shaped longitudinal bands; eye region elevated projecting anteriorly (Fig. 1A, B); with a pair of tubercles between anterior and posterior median eyes; anterior eye row slightly recurved; posterior row straight or slightly recurved as seen from above; median eyes closer to laterals than to each other; clypeus low and flat, usually projecting anteriorly; fovea distinct and long. Abdomen reddish brown (Fig. 1D) with distinct triangular pattern dorsally (Fig. 1A–D); dorsal triangle consists of a yellow area bordered by thin reddish band (Fig. 1B); second smaller pink triangle on the yellow triangle, sometimes with yellow transverse bands. Leg formula 1423; legs II and III, same colour as carapace; legs I and IV brown, with dusky flecks and marks or bands. Epigyne with more or less semicircular openings in a depression with a



Figure 1. Female of *Episinus marignaci* from Mphaphuli Cycad Reserve. **A, B.** Dorsal view. **C, D.** Dorsolateral and lateral views. Photographs: Peter Webb.




Figure 2. Female of *Episinus marignaci*. **A, B.** Epigyne, ventral view. **C, D.** Habitus, dorsal view. A and C reproduced after Lessert (1933).

pair of seminal receptacles present posteriorly (Fig. 2A, B). Specimens lose their colour in alcohol (Fig. 2D).

Habitat. *Episinus marignaci* has been sampled from vegetation using a sweep net or by hand in the Savanna (Foord et al. 2011, 2019), Indian Ocean Coastal Belt, and Thicket biomes (Dippenaar-Schoeman et al. 2020; Fig. 3).

Behaviour. Little is known about the behaviour of this species. According to (Eberhard 1981), members of *Episinus* have been found at ground level between low vegetation and construct very simple H- or Y-shaped webs. In a study of the web of *E. angulatus* (Blackwall, 1836), Holm (1939) found that immature spiders in captivity construct a simple web to catch insects. The spiders sit between two threads. The lower portions of the two threads below the spider are viscid silk and breaks easily. Any insect caught by the strands breaks the thread and




Figure 3. Distribution map of *Episinus marignaci* in South Africa. Ecoregions (Olson et al. 2001) where they occur: ■ Drakensberg montane grasslands, woodlands, and forests, ■ Kwa-Zulu Cape coastal forest mosaic, ■ Lowland Fynbos and renosterveld, ■ Maputaland forest mosaic, ■ Southern African bushveld.

is pulled up and bitten by the spider. When disturbed, the spiders stretched their legs parallel to the body, and on further disturbance would fall to the ground. According to Archer (1946), *E. amoenus* Banks, 1911 preys on ants and lives on evergreen shrubs or in leaf litter in forest areas. Individuals hang with outstretched front legs, like *Tetragnatha* Latreille, 1804, underneath horizontal threads in low shrubs. *Episinus marignac* seems to be nocturnal and during the day they were found on leaves (Fig. 2D).

Discussion

Due to the wide geographical range, the species is listed as Least Concern. There are no significant threats to this species, and it is protected in the Ndumo Game Reserve (Haddad et al. 2006), Nylsvley Nature Reserve (Dippenaar-Schoeman and Prendini 2009), Addo Elephant National Park (Dippenaar-Schoeman et al. 2020), Blouberg Nature Reserve (Foord et al. 2019), uMkhuzi Game Reserve, and Umgeni Valley Nature Reserve.

Acknowledgements

This study was made possible by the financial support from Agricultural Research Council (ARC) and the

National Research Foundation of South Africa under grant number 87311. Comments and suggestions by two reviewers and the editor improved the manuscript significantly.

Authors' Contributions

Conceptualization: ASDS. Data Curation: ASDS. Investigation: ASDS, SHF. Visualization: SHF. Writing – original draft: ASDS. Writing – review and editing: SHF.

References

- Archer AF (1946) The Theridiidae or comb-footed spiders of Alabama. Alabama Museum of Natural History 22: 1–67.
- Dippenaar-Schoeman AS (2014) Field guide to the spiders of South Africa. Lapa, Midrand, South Africa, 432 pp.
- Dippenaar-Schoeman AS, Foord S, Haddad C (2013) Spiders of the savanna biome. Agricultural Research Council, Pretoria, South Africa, 134 pp.
- Dippenaar-Schoeman AS, Haddad CR, Foord SH, Lyle R, Lotz LN, Marais P (2015) South African National Survey of Arachnida (SANSa): review of current knowledge, constraints and future needs for documenting spider diversity (Arachnida: Araneae). Transactions of the Royal Society of South Africa 70: 245–275. <https://doi.org/10.1080/0035919X.2015.1088486>
- Dippenaar-Schoeman AS, Lyle R, Van den Berg A (2012) Bioinformatics on the spiders of South Africa. Serket 13: 121–127.

- Dippenaar-Schoeman AS, Prendini L (2009) Spiders and scorpions (Arachnida; Araneae, Scorpiones) of the Nylsvley Nature Reserve, South Africa. *Koedoe* 51: a161. <https://doi.org/10.4102/koedoe.v51i1.161>
- Dippenaar-Schoeman AS, Wiese L, Foord SH, Haddad CR (2020) A list of spider species found in the Addo Elephant National Park, Eastern Cape province, South Africa. *Koedoe* 62: a1578. <https://doi.org/10.4102/koedoe.v62i1.1578>
- Eberhard WG (1981) The single line web of *Phoroncidia studo* Levi (Araneae: Theridiidae): a prey attractant? *Journal of Arachnology* 9: 229–232.
- Foord S, Dippenaar-Schoeman AS, Haddad CR, Schoeman C, Hahn N, Lyle R (2019) Spider checklist for the Blouberg, in the Vhembe Biosphere Reserve, South Africa. *Bothalia* 49: a2455. <https://doi.org/10.4102/abc.v49i1.2455>
- Foord SH, Dippenaar-Schoeman AS, Haddad CR, Lotz L, Lyle R (2011) The faunistic diversity of spiders (Arachnida: Araneae) of the Savanna Biome in South Africa. *Transactions of the Royal Society of South Africa* 66: 170–201. <https://doi.org/10.1080/0035919X.2011.639406>
- Haddad CR, Dippenaar-Schoeman AS, Wesołowska W (2006) A checklist of the non-acarine arachnids (Chelicerata: Arachnida) of the Ndumo Game Reserve, Maputoland, South Africa. *Koedoe* 49: a116. <https://doi.org/10.4102/koedoe.v49i2.116>
- Holm A (1939) Beitrage zur Biologie der Theridiiden. *Festschrift zum 60 Geburtstag von Professor Dr. Embrik Strand* 5: 56–57.
- Lessert, R (1933) Araignées d'Angola. Resultats de la Mission scientifique suisse en Angola 1928–1929. *Revue Suisse de Zoologie* 404: 85–59.
- Levi H, Levi L (1962) The genera of the spider family Theridiidae. *Bulletin of the Museum of Comparative Zoology* 127: 1–71.
- Olson DM, Dinnerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, D'amico JA, Itoua I, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Ricketts TH, Kura Y, Lamoreux JF, Wettengel WW, Hedao P, Kassem KR (2001) Terrestrial ecoregions of the world: a new map of life on earth. *Bioscience* 51: 933–938. <https://doi.org/c635xt>
- World Spider Catalog (2022) World Spider Catalog. Version 23.5. Natural History Museum, Bern. <https://wsc.nmbe.ch>. Accessed on: 2022-9-12.