




Distribution of *Nemipterus marginatus* (Valenciennes, 1830) (Nemipteridae, Perciformes) in the estuary and coastal areas of North Central, Vietnam

Hoang Ngoc Thao^{1*}, Trinh Thi Thu¹, Le Thi Ha², Nguyen Thi Ly³, Dao Thi Hong⁴

¹ Laboratory of Zoology, Department of Biology, Faculty of Nature Science, Hong Duc University, Thanh Hoa Province, Vietnam • HNT: hoangngocthao@hdu.edu.vn  <https://orcid.org/0000-0001-9305-5518> • TTT: thukxs@gmail.com

² Tinh Gia 3 High School, Thanh Hoa Thanh Hoa Province, Vietnam • LTH: tranhaan932010@gmail.com

³ Le Van Huu High School, Thanh Hoa Thanh Hoa Province, Vietnam • NTL: lyhieuvanha@gmail.com

⁴ Dong Son 1 High School, Thanh Hoa Thanh Hoa Province, Vietnam • DTH: hongdt.thptdongson1@thanhhoa.edu.vn

* Corresponding author

Abstract

We provide data on morphological characteristics and the distribution of the nemipterid fish *Nemipterus marginatus* (Valenciennes, 1830) in the estuary and coastal area of Thanh Hoa province, North Central Vietnam. This is also a clear location record for the distribution of this species in the North Central region compared to previous studies, and a new record for the study area.

Keywords

Nemipterid fish, new distribution, Thanh Hoa province

Academic editor: Hudson Tercio Pinheiro | Received 6 March 2022 | Accepted 8 October 2022 | Published 18 October 2022

Citation: Hoang NT, Trinh TT, Le TH, Nguyen TL, Dao TH (2022) Distribution of *Nemipterus marginatus* (Valenciennes, 1830) (Nemipteridae, Perciformes) in the estuary and coastal areas of North Central, Vietnam. Check List 18 (5): 1127–1130. <https://doi.org/10.15560/18.5.1127>

Introduction

Species of the genus *Nemipterus* (Nemipteridae) compose an important group of commercial fishes in Vietnam. About 25 species of *Nemipterus* have been described (FishBase 2021), of which 12 species have been recognized in Vietnam coastal waters (Le et al. 2012), including: *Nemipterus bathybius* (Snyder, 1911), *N. celebicus* (Bleeker, 1854), *N. furcosus* (Valenciennes, 1830), *N. hexodon* (Quoy & Gaimard, 1824), *N. japonicus* (Bloch, 1791), *N. marginatus* (Valenciennes, 1830), *N. nematophorus* (Bleeker, 1853), *N. nemurus* (Bleeker, 1857), *N. peronii* (Valenciennes, 1830), *N. tambuloides* (Bleeker, 1853), *N. virgatus* (Houttuyn, 1782), and *N. zysron* (Bleeker, 1857). In Vietnam, *N. marginatus* has been

recorded in the South and South-Central regions (Dinh and Phan 2010; Le et al. 2012). Nguyen et al. (1995) also confirmed its distribution in the north but gave no specific location. During specimen collection in the estuary and coastal areas of Thanh Hoa province, we found specimens of *N. marginatus*, which represent records for the study area.

Methods

Fieldwork was conducted in August 2020 in the coastal area of Hai Binh commune, Nghi Son town, and October 2021 in Hoi estuary of the Ma river, Sam Son town,

Thanh Hoa province, Vietnam (Fig. 1). Specimens were preserved and deposited at the Animal Laboratory, Hong Duc University; 16 specimens were examined. The technical terms and measurements follow the description by Russell (1990). The abbreviations: total length (TL), standard length (SL), pre dorsal length (PDL), post dorsal length (PDL1), head length (HL), post orbital length (POL), snout length (SnL), eye diameter (ED), Interorbital distance (ID), Body depth (BD), caudal peduncle length (CPL); dorsal fin rays (DF), pectoral fin rays (PF), ventral fin rays (VF), anal fin rays (AF), caudal fin rays (CF), and number lateral-line scale (L). Field numbers are abbreviated: LB (for specimens collected in Lach Bang estuary, Hai Binh commune), CH (for specimens collected in Hoi estuary).

Results

Nemipterus marginatus (Valenciennes, 1830)

New records. VIETNAM – Thanh Hoa Province • Nghi Son town, Hai Binh commune, Hai Binh fishing port; 19°24'N, 105°46'E; 10.VIII.2020; Le Thi Ha, Nguyen Thi Ly, specimens collected from fishermen's boats; 15 spec., sex indeterminate, LB.001, LB.002, LB.003, LB.004, LB.005, LB.006, LB.007, LB.008, LB.009, LB.010, LB.011 (Fig. 2), LB.012, LB.013, LB.014, LB.015 • Sam Son town, Hoi estuary of the Ma river, Hoi fishing port; 19°46'N, 105°53'E; 22.X.2021; Dao Thi Hong, specimen collected from fishermen's boats fishing in the estuary area; 1 spec., sex indeterminate, CH.75.

Identification. Our specimens agree well with the description of the species by Russell (1990): body depth 3.43–3.91 in standard length; mouth oblique, maxillary extending to the anterior border of eye, snout length 0.63–0.92 in diameter of eye; eye medium-sized, diameter of eye 3.3–4.06 in head length, and 1.09–1.59 in snout length; lower margin of eye above a line from tip of snout to upper of pectoral fin; interorbital width 1.08–1.96 in eye; suborbital depth 1.7–1.9 in eye diameter; pectoral fins long, 0.86–1.33 in head length, extending to between level of anus and origin of anal fin; pelvic reaching to first or second anal rays; caudal fin forked, upper lobe produced into a short filament, filament length 11.6–33.0 mm; teeth in both the lower and upper jaw, 4 or 5 pairs of small canines in front of the upper jaw.

Morphometrics: dorsal fin X, 9; anal fin III, 7; pectoral fin 2, 15–16; pelvic fin I, 5; caudal fin 17–18; lateral-line scales 50–53. Ratios: SL/BD = 3.64 (3.43–3.91), SL/HL = 3.70 (3.33–4.06), SL/PDL = 3.06 (2.69–3.33), SL/PDL1 = 1.20 (1.15–1.23), SL/CPL = 6.98 (6.50–8.65). HL/SnL = 3.03 (2.52–3.36), HL/ED = 3.78 (3.3–4.32), HL/POL = 1.62 (1.44–1.83), HL/ID = 5.79 (3.92–7.2) ID. SnL/POL = 0.54 (0.47–0.61), SnL/ED = 1.26 (1.09–1.59). ID/ED = 0.68 (0.51–0.93).

Colour: body rosy dorsally and on upper half of flanks, belly silvery; head with a narrow yellow stripe from below nostrils to eye and another yellow stripe from the front part of upper jaw to eye; a broad yellow-orange band divided above the pectoral fin, along body from below origin of lateral line to the upper part of the




Figure 1. Distribution of *Nemipterus marginatus*. The yellow area is distribution of this species according to IUCN (2019). The red square is new records.



Figure 2. *Nemipterus marginatus* from the coast of North Central Vietnam (LB.005). Scale bar = 1 cm.

caudal peduncle; a second broad yellow band from above the base of the pectoral fin to the lower part of caudal peduncle; dorsal fin with a yellow to orange margin, and broad yellow median band; anal fin with two narrow yellow bands; pelvic fins pink; pectoral fins translucent; caudal fin pink, including filament on the upper lobe.

Discussion

The specimens collected from Thanh Hoa province, Vietnam agree well with the description of *Nemipterus marginatus* by Russell (1990) and Ping et al. (2011) (Table 1).

Russell (1990, 2001) reported maximum size of *Nemipterus marginatus* as 150 mm SL, commonly 130 mm SL; but specimens we collected ranged in size to 180 mm SL (TL 245 mm, LB.001).

Nemipterus marginatus is distributed in the western

Pacific, including the South China Sea, Strait of Malacca, southern coasts of Sumatra and Java, northern Australia, the Gulf of Papua, and the Solomon Islands (Russell 1990, 2001; IUCN 2019). Ping et al. (2011) recorded this species on the southern coast of China. In Vietnam, a report by Nguyen et al. (2014) in the Hoi estuary of Ma river, Thanh Hoa Province, recorded 115 fish species belonging to 38 families and 13 orders; similarly, Duong Quang Ngoc (2007) recorded 263 fish species, 58 families, and 14 orders in research on the fish fauna of the Ma river basin in Vietnam. However, no species of the family Nemipteridae were recorded in either of these studies. Thus, the research results here have added a new record on the distribution of *N. marginatus* for the estuary and coastal areas of Thanh Hoa province, Vietnam.

Acknowledgements

We are very grateful to the fishermen in the Hoi estuary (Sam Son town) and Lach Bang estuary (Hai Binh commune) who supported us during field surveys and the collection of specimens. We would also like to thank Dr. Hudson Tercio Pinheiro and Dr. Robert Forsyth for support in the review process and English language editing; Dr. Barry Russell provided helpful comments on the manuscript.

Authors' Contributions

Data curation: HNT. Formal analysis: TTT, LTH, NTL, DTH. Methodology: HNT, TTT. Supervision: HNT. Writing – original draft: TTT. Writing – review and editing: HNT.

Table 1. Comparison of *Nemipterus marginatus* specimens from Vietnam with the descriptions by Russell (1990) and Ping et al. (2011)

Character	Vietnam	Russell 1990	Ping et al. 2011
Dorsal fin count	X, 9	X, 9	X, 9
Anal fin count	III, 7	III, 7	III, 7
Canines	Small, recurved canines anteriorly in upper jaw		
Standard length/body depth	3.4–3.9	3.1–3.8	3.5–3.9
Head length/eye	3.3–4.3	3.4–4.3	3.3–3.6
Head length/pectoral fin	0.9–1.3	0.9–1.3	0.9–1.0
Head length/pelvic fin	0.9–1.5	0.9–1.3	1.1–1.3
Eye/interorbital width	1.1–2.0	1.5–2.2	1.1–1.7
Eye/suborbital width	1.7–1.9	1.0–2.6	1.0–1.3
Eye/snout length	0.6–0.9	0.8–1.4	0.9–1.0

References

- Dinh PA, Phan TH (2010) Fish category in the Southern Sea of Sontra Peninsula, Danang city. *Science and Technology Journal, Da Nang University*, No 1(36): 56–64 [in Vietnamese].
- Duong QN (2007) Contributing to the study of fish in the Ma river basin in Vietnam. PhD thesis, The University of Pedagogy, Ha Noi, Vietnam, 159 pp [in Vietnamese].
- Froese R and Pauly D (Editors) (2021) FishBase. World Wide Web electronic publication. <http://www.fishbase.org>. Accessed: 2022-3-09.
- IUCN Marine Biodiversity Unit/GMSA (2019) *Nemipterus marginatus*. The IUCN Red List of threatened species, version 2021-3. <https://www.iucnredlist.org/species/75553587/75553596>. Accessed: 2022-3-09.
- Le TTT, Vo VQ, Nguyen PUV (2012) A checklist of the Mojarras, threadfin breams and grunters (family: Gerreidae, Nemipteridae, Terapontidae) in the Vietnamese seawaters. *Anthology of marine research* 18: 119–126 [in Vietnamese].
- Nguyen HP (ed), Le TP, Nguyen NT, Nguyen PD, Do TTN, Nguyen VL (1995) List of marine fish in Vietnam. Vol. 3. Perciformes, Percoidae and Echeneoidei. Science and Technology Publishing House, Ha Noi, Vietnam, 607 pp [in Vietnamese].
- Nguyen XH, Nguyen TN, Le DG, Vu TT (2014) Biodiversity of fish in the coastal area of Cua Hoi, Thanh Hoa province. In: *Proceedings of the national scientific conference on marine biology and sustainable development (2nd)*, Ha Noi, Vietnam, 79–85 [in Vietnamese].
- Russell BC (1990) FAO species catalogue. Volume 12. Nemipterid fishes of the world. Rome, FAO, 149 pp. + viii pls.
- Russell BC (2001) Nemipteridae. In: Carpenter KE, Niem VH (Eds.) *The living marine resources of the Western Central Pacific. Volume 5. FAO species identification guide for fishery purposes*. FAO, Rome, Italy, 2791–3380.
- Ping N, Rennie W, Jing L (2011) First record of Red Filment Threadfin Bream, *Nemipterus marginatus* (Valenciennes, 1830) (Perciformes, Nemipteridae), from Chinese waters. *Chinese Journal of Oceanology and Limnology* 29 (6): 1306–1308. <https://doi.org/10.1007/s00343-011-0312-3>