

Conference Abstract

Desktop or remote knowledge base management systems for taxonomic data and identification keys: Xper2 and Xper3

Régine Vignes-Lebbe[‡], Sylvain Bouquin[‡], Adeline Kerner[§], Estelle Bourdon[|]

[‡] Institut de Systématique, Évolution, Biodiversité ISYEB - UMR 7205 – CNRS, MNHN, UPMC, EPHE UPMC Univ. Paris 06, Sorbonne Universités 57 rue Cuvier, CP48 F-75005, Paris, France, Paris, France

[§] CNRS UMR 7207, MNHN, Paris, France
[|] MNHN, Paris, France

Corresponding author: Régine Vignes-Lebbe (regine.vignes_lebbe@upmc.fr)

Received: 30 Jul 2017 | Published: 07 Aug 2017

Citation: Vignes-Lebbe R, Bouquin S, Kerner A, Bourdon E (2017) Desktop or remote knowledge base management systems for taxonomic data and identification keys: Xper2 and Xper3. Proceedings of TDWG 1: e19911. <https://doi.org/10.3897/tdwgproceedings.1.19911>

Abstract

Storing, sharing, managing taxonomic data is crucial for research and systematic outreach. In this context, Xper2 (Ung et al. 2010) and Xper3 (Vignes Lebbe et al. 2015) are two platforms dedicated to taxonomic descriptions and computer-aided identification. These freeware provide a sleek and easy to use interface, and do not require computer skills. Xper2 is available on Windows, MacOS or Linux. Download the software on [the website](#). Xper3 gives users the ability to work simultaneously and remotely on the same knowledge base. Create your own account on [the online platform](#).

The numerous knowledge bases (over one thousand active KB) already created and published with the two platforms prove their relevance and efficiency (See for example Mathieu et al. 2012, Kerner 2012, Corvez and Grand 2014, Martin et al. 2015, Engel et al. 2016, Saïd et al. 2017). Scientists can share their data on phenotypes (structured descriptions, documented with images, videos, and text including bibliography and external links), compare phenotypes, and import or export partial or total content in various standard formats such as SDD (Structured Descriptive Data, a TDWG data standard), CSV (Comma Separated Values, used in Spreadsheet), and NEXUS (used in

phylogenetic analyses, Maddison et al. 1997) for external analyses. KB can be exchanged between Xper2 and Xper3 via SDD files.

We propose demos of the desktop version (Xper2) and online version (Xper3), illustrated with various contents on fossils, botany, marine species and arthropods. The TDWG 2017 participants are invited to come with their own data (list of taxa, identification keys, character-taxon matrices). They will learn and test how to import existing content into Xper KB, and how to build and publish keys (free access keys and single access keys) automatically.

Keywords

Xper2, Xper3, knowledge base, identification keys, online platform, taxonomy

Presenting author

Régine Vignes-Lebbe

References

- Corvez A, Grand A (2014) Enabling comparisons of characters using an Xper2 based knowledge-base of fern morphology. *Phytotaxa* 183 (3): 145-158. <https://doi.org/10.11646/phytotaxa.183.3.2>
- Engel J, Brousseau L, Baraloto C (2016) GuiaTreeKey, a multi-access electronic key to identify tree genera in French Guiana. *PhytoKeys* 68: 27-44. <https://doi.org/10.3897/phytokeys.68.8707>
- Kerner A (2012) Archaeocyatha, a computer-aided identification of genera. *Geologica Belgica* 15 (4): 209-214.
- Maddison DR, Swofford DL, Maddison WP (1997) Nexus: An Extensible File Format for Systematic Information. *Systematic Biology* 46 (4): 590-621. <https://doi.org/10.1093/sysbio/46.4.590>
- Martin P, Schmelz R, Dole-Olivier M (2015) Groundwater oligochaetes (Annelida, Clitellata) from the Mercantour National Park (France), with the descriptions of one new genus and two new stygobiont species. *Zoosystema* 37 (4): 551-570. <https://doi.org/10.5252/z2015n4a2>
- Mathieu B, Cêtre-Sossah C, Garros C, Chavernac D, Balenghien T, Carpenter S, Setier-Rio M, Vignes-Lebbe R, Ung V, Candolfi E, Delécolle J (2012) Development and validation of IIKC: an interactive identification key for Culicoides (Diptera: Ceratopogonidae) females from the Western Palaearctic region. *Parasites & Vectors* 5 (1): 137. <https://doi.org/10.1186/1756-3305-5-137>
- Saïd A, Hennequin S, Rouhan G, Dubuisson J (2017) Disentangling the diversity and taxonomy of Hymenophyllaceae (Hymenophyllales, Polypodiidae) in the Comoros. *European Journal of Taxonomy* 313: 1-53. <https://doi.org/10.5852/ejt.2017.313>

- Ung V, Dubus G, Zaragüeta-Bagils R, Vignes-Lebbe R (2010) Xper²: introducing e-Taxonomy. *Bioinformatics* 26 (5): 703-704. <https://doi.org/10.1093/bioinformatics/btp715>
- Vignes Lebbe R, Chesselet P, Diep Thi MH (2015) Xper3: new tools for collaborating, training and transmitting knowledge on botanical phenotypes. In: Rakotoarisoa NR, Blackmore S, Riéra B (Eds) *Botanists of the 21st century*. 11 pp. [ISBN 978-92-3-100120-8].