

Number and location of rDNA clusters in the superfamilies Tenthredinoidea and Cynipoidea (Hymenoptera): an update

Vladimir E. Gokhman¹, Valentina G. Kuznetsova², Boris A. Anokhin²

1 *Russian Entomological Society, Moscow, Russia* **2** *Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia*

Corresponding authors: Vladimir E. Gokhman (vegokhman@hotmail.com);

Valentina G. Kuznetsova (valentina.kuznetsova@zin.ru)

Academic editor: Snezana Grozeva | Received 20 November 2024 | Accepted 23 November 2024 | Published 4 December 2024

<https://zoobank.org/983F7C54-D9E4-45A1-A296-80D8AC5742BD>

Citation: Gokhman VE, Kuznetsova VG, Anokhin BA (2024) Number and location of rDNA clusters in the superfamilies Tenthredinoidea and Cynipoidea (Hymenoptera): an update. *Comparative Cytogenetics* 18: 239–246. <https://doi.org/10.3897/compcytogen.18.142301>

Abstract

To identify nucleolus organizing regions (NORs), fluorescence *in situ* hybridization (FISH) with 18S rDNA probe was performed on chromosomes of *Tenthredo campestris* Linnaeus, 1758 (Tenthredinidae), *Arge ciliaris* (Linnaeus, 1767) (Argidae) ($n = 10$ in both) and *Aulacidea hieracii* (Bouché, 1834) (Cynipidae) ($2n = 20$). In all these species, a single pericentromeric rDNA cluster per haploid karyotype was detected. This number of NORs is confirmed as ancestral for the order Hymenoptera.

Keywords

Apocrita, Argidae, chromosomes, Cynipidae, Hymenoptera, rDNA-FISH, Symphyta, Tenthredinidae

Hymenoptera represent one of the largest insect orders, with the approximate number of described species far exceeding 150,000 (Huber 2017). However, the overwhelming majority of this taxonomic diversity belongs to the suborder Apocrita, or higher Hymenoptera (see, e.g., Forbes et al. 2018), whereas the substantially less speciose Symphyta (= lower Hymenoptera) harbor less than nine thousand members (Huber 2017).

Nevertheless, Symphyta include the least advanced Hymenoptera, and therefore studying these taxa is necessary to reconstruct ancestral character states for the order in general. Among other characteristics, this also applies to different karyotype features of Hymenoptera (Gokhman 2023a). For example, the ancestral nature of the canonical “insect-type” telomeric repeat in this order, TTAGG, was proven only when telomere structure of certain sawflies has been studied (Gokhman and Kuznetsova 2018; Lukhtanov and Pazhenkova 2023). However, other characteristics of symphytan chromosome sets, i.e., the number and location of clusters of 45S ribosomal DNA (rDNA), were studied in just a few members of the superfamily Tenthredinoidea, which belong to the families Tenthredinidae, Athaliidae and Diprionidae (Rousselet et al. 1999, 2000; Kuznetsova et al. 2001; Matsumoto et al. 2002). During these studies, different techniques for revealing 45S rDNA sites, which represent nucleolus organizing regions (NORs), i.e., AgNOR-banding, staining GC-enriched chromosome segments with chromomycin A₃ (CMA₃) and *in situ* hybridization with rDNA probes, including that using fluorescent dyes (FISH), were employed (Gokhman and Kuznetsova 2024). Nevertheless, the ancestral number and location of these rDNA clusters in the Symphyta remain ambiguous, since haploid karyotypes of two members of the genus *Tenthredo* Linnaeus, 1758, *T. velox* Fabricius, 1798 and *T. arcuata* Forster, 1771, were shown to have one and two 45S rDNA sites, respectively (Kuznetsova et al. 2001). Moreover, another species of the family Tenthredinidae, *Rhogogaster viridis* (Linnaeus, 1758), as well as *Diprion pini* (Linnaeus, 1758) and *Neodiprion abietis* (Harris, 1841) (Diprionidae) also have single 45S rDNA clusters (Rousselet et al. 1999, 2000; Kuznetsova et al. 2001), but the haploid karyotype of *Athalia rosae* (Linnaeus, 1758), which is now placed in a separate family Athaliidae (Wutke et al. 2024), carries four sites of that kind (Matsumoto et al. 2002).

Although much more is now known about the number and location of NORs in Apocrita (Gokhman and Kuznetsova 2024), reconstruction of the ancestral number and location of these parameters in this suborder is also far from straightforward. Specifically, both the number and location of these sites vary substantially across parasitoids as well as across aculeate Hymenoptera. Whilst the only 45S rDNA cluster per haploid karyotype is a widespread condition in Apocrita (see, e.g., Gokhman et al. 2024), chromosome sets of these insects can harbor up to six and even 15 NORs in parasitic wasps and Aculeata, respectively (Gokhman and Kuznetsova 2024). FISH also visualized single 45S rDNA clusters in all studied members of the superfamily Cynipoidea, i.e., in *Diplolepis rosae* (Linnaeus, 1758) (Cynipidae) (Gokhman et al. 2014) as well as in four species of the family Figitidae (Gokhman et al. 2016). In addition, karyotypes of three other members of Cynipidae, *Aulacidea hieracii* (Bouché, 1834), *Isocolus jaceae* (Schenck, 1863) and *I. scabiosae* (Giraud, 1859), have been recently examined using CMA₃ (Gokhman 2021). Although single CMA₃-positive sites per haploid chromosome set were detected in all these species, these results could be corroborated using FISH, since CMA₃-positive chromosome segments do not always represent NORs (see, e.g., Gromicho et al. 2005; Gokhman et al. 2017). We have therefore undertaken the present study to further identify the number and location of rDNA clusters in the superfamilies Tenthredinoidea and Cynipoidea. The results of this study are given below.

Material and methods

Origin of insects

Adult sawflies and galls containing immature stages of *A. hieracii* were collected in 2022 by the first author near Ozhigovo, Russia (about 60 km SW Moscow) as well as by M.I. Nikelshparg (Saratov State University, Saratov, Russia) near the city of Saratov, Russia (about 730 km SE Moscow), respectively. All sawflies were identified by S.A. Basov (Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia).

Preparation of chromosomes

Chromosome preparations were obtained according to the guidelines provided by Naito (1982) and Imai et al. (1988) with a few modifications. Adult female sawflies were dissected in small Petri dishes in distilled water, unfertilized mature eggs were extracted from their bodies, placed into the dishes on a filter paper soaked with distilled water, and then incubated for 3–4 days at room temperature (RT). Haploid embryos and cerebral ganglia were extracted from the eggs and gall wasp prepupae, respectively, and dissected in 0.5% hypotonic sodium citrate solution containing 0.005% colchicine. The embryos and ganglia were then transferred to a fresh portion of hypotonic solution and incubated for about 30 min at room temperature. The material was transferred onto a pre-cleaned microscope slide using a Pasteur pipette and then gently flushed with Fixative I (glacial acetic acid: absolute ethanol: distilled water 3:3:4). The tissues were disrupted using dissecting needles in an additional drop of Fixative I. A drop of Fixative II (glacial acetic acid: absolute ethanol 1:1) was applied to the center of the area, and the more aqueous phase was blotted off the edges of the slide. The slides were then dried and stored at RT for a few weeks.

FISH procedure

Genomic DNA from a male *Pyrrhocoris apterus* (Linnaeus, 1758) (Hemiptera, Heteroptera, Pyrrhocoridae) was isolated using CTAB extraction method. FISH with the 18S rRNA gene probe was carried out on chromosomes of all studied species. The target 18S rRNA gene was PCR amplified (see Grozeva et al. 2011 for the table of primers) from the genomic DNA of *P. apterus*, and labeled by PCR with biotin.

In situ hybridization was performed as described by Schwarzacher and Heslop-Harrison (2000) with modifications. Chromosome preparations were dehydrated through 70, 80 and 96% ethanol at RT and treated with 100 µg/ml RNase A (Sigma) for 30 min at 37 °C in a humid chamber; washed three times in 2× SSC (5 min each) at RT; dehydrated through 70, 80 and 96% ethanol at RT; incubated in 5 mg/ml pepsin in 0.01 N HCl for 10 min at 37 °C; washed sequentially in PBS,

in PBS containing 0.05 M MgCl₂ for 5 min each, in 1% PFA in PBS containing 0.05 M MgCl₂ for 15 min, in PBS for 5 min, in PBS containing 0.05 M MgCl₂ for 5 min at RT each; dehydrated through 70, 80 and 96% ethanol at RT and finally, dried. After pretreatment, 6.5 µl hybridization mixture containing about 100 ng of labeled probe, 50% formamide, 2× SSC, 10% (w/v) dextran sulfate, 1% (w/v) Tween 20 and 10 µg salmon-sperm DNA was added on preparations. Slides were mounted using glass coverslips and rubber cement. The slides were denatured for 5 min at 75 °C. The chromosome slides were then incubated for 42–44 h at 37 °C. Following hybridization, the slides were washed in 2× SSC for 3 min at 45 °C, then in 50% formamide in 2× SSC for 10 min at 45 °C, two times in 2× SSC (10 min each) at 45 °C, blocked in 4× SSC containing 1.5% (w/v) BSA and 0.1% Tween 20 for 30 min at 37 °C in a humid chamber. 18S rRNA gene probe was detected with 5 µg/ml Avidin-FITC (Invitrogen). Detection was performed in 4× SSC containing 1.5% BSA and 0.1% Tween 20 for 1 h at 37 °C. Slides were washed three times in 4× SSC containing 0.02% Tween 20 (10 min each) at 45 °C and dehydrated through 70, 80 and 96% ethanol at RT. Chromosomes were mounted in an antifade medium (ProLong Gold antifade reagent with DAPI, Invitrogen) and covered with a glass coverslip.

Image acquisition and analysis

Metaphase plates and interphase nuclei were analyzed under a Leica DM 6000B microscope with a 100× objective. Fluorescence images were taken with a Leica DFC 345 FX camera using Leica Application Suite 4.5.0 software with an Image Overlay module. To prepare illustrations, the resulting images were arranged and enhanced using GIMP 2.10. Chromosomes were classified according to the guidelines provided by Levan et al. (1964).

Results

Superfamily Tenthredinoidea

Family Tenthredinidae

Tenthredo campestris Linnaeus, 1758

A detailed description of the karyotype of this species can be found in Gokhman (2024). The haploid set of *T. campestris* includes 10 biarmed chromosomes. In the karyotype of this species, the first chromosome is about 1.3 times longer than the second one, which is, in turn, also 1.3 times longer than the third chromosome. The remaining chromosomes more or less gradually decrease in size. The haploid karyotype of *T. campestris* harbors the only pericentromeric rDNA cluster on a particular medium-sized chromosome (Fig. 1A).




Figure 1. FISH with 18S rDNA probe on Hymenoptera chromosomes. *Tenthredo campestris*, $n = 10$ (A), *Arge ciliaris*, $n = 10$ (B), *Aulacidea hieracii*, $2n = 20$ (C). Probe signals are indicated in green. Scale bar: 10 μm .

Family Argidae

Arge ciliaris (Linnaeus, 1767)

A detailed karyotypic description of this species is given by Gokhman (2023b). The haploid set of *A. ciliaris* harbors 10 biarmed chromosomes. The first chromosome is about 1.8 times longer than the second one; the remaining chromosomes gradually decrease in size. On some metaphase plates, the second metacentric bears an obvious pericentromeric secondary constriction on its longer arm (see Gokhman 2023b). It is therefore not surprising that the only 18S rDNA site revealed in this species by FISH is apparently co-localized with this constriction (Fig. 1B).

Superfamily Cynipoidea

Family Cynipidae

Aulacidea hieracii (Bouché, 1834)

A detailed description of the karyotype of this species, including results of CMA₃ staining can be found in Gokhman (2021). The diploid set of *A. hieracii* includes 20 chromosomes, which gradually decrease in size, with a single pericentromeric CMA₃-positive band located on the largest metacentric. FISH with the 18S rDNA probe revealed the only paired site in place of the CMA₃-positive band (see Gokhman 2021), thus suggesting that both used techniques visualized the same rDNA cluster (Fig. 1C).

Discussion

Haploid karyotypes of both studied members of Tenthredinoidea, *T. campestris* (Tenthredinidae) and *A. ciliaris* (Argidae), harbor single rDNA clusters. Moreover, this character state predominates within Tenthredinoidea (see above), and this therefore sug-

gests that the single rDNA site per haploid chromosome set is ancestral at least for the whole superfamily. This is further corroborated by the fact that *A. ciliaris* remains the only examined member of the Argidae + Pergidae clade, a sister one to the remaining Tenthredinoidea except Blasticotomidae (Wutke et al. 2024), the most basal family which chromosomes are totally unknown. If this is true, then higher numbers of rDNA clusters found in *Tenthredo arcuata* and *Athalia rosae* (Kuznetsova et al. 2001; Matsumoto et al. 2002) apparently represent apomorphic conditions. Anyway, studying these clusters in other superfamilies of the Symphyta could either confirm or reject this assumption.

In turn, Tenthredinoidea also represent a sister clade to all other Hymenoptera, except for the family Xyelidae (Wutke et al. 2024), in which the structure and location of rDNA sites are still unknown (see Gokhman 2023a). Nevertheless, a single 45S rDNA cluster per haploid karyotype can be found in the majority of parasitoid Hymenoptera including all known Cynipoidea (Gokhman et al. 2014, 2016; Gokhman 2022; present paper). Furthermore, the same character state is considered ancestral for Aculeata, despite wide variation in the number and location of these clusters in aculeate Hymenoptera (Menezes et al. 2021). Taken together, all this information suggests that a single rDNA site per haploid genome is also ancestral for the whole order Hymenoptera. Again, this does not seem surprising, since a similar pattern is apparently characteristic of the class Insecta in general (Gokhman and Kuznetsova 2024).

Acknowledgements

The authors are grateful to S.A. Basov (Zoological Institute, Russian Academy of Sciences) for identifying the sawflies as well as to M.I. Nikelshparg (Saratov State University, Saratov, Russia) for providing living material on *A. hieracii*.

This research was funded within the framework of the State Research Projects Nos 122031100272-3 (V.G.K.) and 122031100275-4 (B.A.A.).

References

- Forbes AA, Bagley RK, Beer MA, Hippee AC, Widmayer HA (2018) Quantifying the unquantifiable: why Hymenoptera, not Coleoptera, is the most speciose animal order. *BMC Ecology* 18: 21. <https://doi.org/10.1186/s12898-018-0176-x>
- Gokhman VE (2021) Chromosomes of three gall wasps of the tribe Aylacini (Hymenoptera, Cynipidae). *Comparative Cytogenetics* 15(2): 171–178. <https://doi.org/10.3897/compcytogen.v15.i2.66781>
- Gokhman VE (2022) Comparative karyotype analysis of parasitoid Hymenoptera (Insecta): major approaches, techniques, and results. *Genes* 13: 751. <https://doi.org/10.3390/genes13050751>
- Gokhman VE (2023a) Chromosomes of Symphyta (Hymenoptera): current state and perspectives of research. *Russian Entomological Journal* 32(4): 375–382. <https://doi.org/10.15298/rusentj.32.4.02>

- Gokhman VE (2023b) Chromosomes of the genus *Arge* Schrank, 1802 (Hymenoptera, Argeidae): new data and review. *Comparative Cytogenetics* 17: 327–333. <https://doi.org/10.3897/compcytogen.17.115485>
- Gokhman VE (2024) Morphometric analysis of chromosomes of sawflies of the genera *Tenthredo* Linnaeus, 1758 and *Athalia* Leach, 1817 (Hymenoptera, Tenthredinidae, Athaliidae). *Russian Entomological Journal* 33(4).
- Gokhman VE, Kuznetsova VG (2018) Presence of the canonical TTAGG insect telomeric repeat in the Tenthredinidae (Symphyta) suggests its ancestral nature in the order Hymenoptera. *Genetica* 146: 341–344. <https://doi.org/10.1007/s10709-018-0019-x>
- Gokhman VE, Kuznetsova VG (2024) Structure and evolution of ribosomal genes of insect chromosomes. *Insects* 15: 593. <https://doi.org/10.3390/insects15080593>
- Gokhman VE, Anokhin BA, Kuznetsova VG (2014) Distribution of 18S rDNA sites and absence of the canonical TTAGG insect telomeric repeat in parasitoid Hymenoptera. *Genetica* 142: 317–322. <https://doi.org/10.1007/s10709-014-9776-3>
- Gokhman VE, Bolsheva NL, Govind S, Muravenko OV (2016) A comparative cytogenetic study of *Drosophila* parasitoids (Hymenoptera, Figitidae) using DNA-binding fluorochromes and FISH with 45S rDNA probe. *Genetica* 144: 335–339. <https://doi.org/10.1007/s10709-016-9902-5>
- Gokhman VE, Luna MG, Vallina C, Bressa MG (2024) Chromosomes of *Pseudapanteles dignus* (Muesebeck, 1938) and a review of known karyotypes of the subfamily Microgastriinae (Hymenoptera, Braconidae). *Comparative Cytogenetics* 183: 199–211. <https://doi.org/10.3897/compcytogen.18.133534>
- Gokhman VE, Pereira FF, Costa MA (2017) A cytogenetic study of three parasitic wasp species (Hymenoptera, Chalcidoidea, Eulophidae, Trichogrammatidae) from Brazil using chromosome morphometrics and base-specific fluorochrome staining. *Comparative Cytogenetics* 11(1): 179–188. <https://doi.org/10.3897/compcytogen.v11i1.11706>
- Gromicho M, Ozouf-Costaz C, Collares-Pereira MJ (2005) Lack of correspondence between CMA₃-, Ag-positive signals and 28S rDNA loci in two Iberian minnows (Teleostei, Cyprinidae) evidenced by sequential banding. *Cytogenetic and Genome Research* 109: 507–511. <https://doi.org/10.1159/000084211>
- Grozeva S, Kuznetsova VG, Anokhin BA (2011) Karyotypes, male meiosis and comparative FISH mapping of 18S ribosomal DNA and telomeric (TTAGG)_n repeat in eight species of true bugs (Hemiptera, Heteroptera). *Comparative Cytogenetics* 5(4): 335–374. <https://doi.org/10.3897/compcytogen.v5i4.2307>
- Huber JT (2017) Biodiversity of Hymenoptera. In: *Insect Biodiversity: Science and Society* 1, 2nd edn. Footitt RG, Adler PH, 419–462. Oxford, UK, Wiley Blackwell. <https://doi.org/10.1002/9781118945568.ch12>
- Imai HT, Taylor RW, Crosland MWJ, Crozier RH (1988) Modes of spontaneous chromosomal mutation and karyotype evolution in ants with reference to the minimum interaction hypothesis. *Japanese Journal of Genetics* 63: 159–185. <https://doi.org/10.1266/jjg.63.159>
- Kuznetsova VG, Westendorff M, Nokkala S (2001) Patterns of chromosome banding in the sawfly family Tenthredinidae (Hymenoptera, Symphyta). *Caryologia* 54: 227–233. <https://doi.org/10.1080/00087114.2001.10589230>

- Levan A, Fredga K, Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201–220. <https://doi.org/10.1111/j.1601-5223.1964.tb01953.x>
- Lukhtanov VA, Pazhenkova EA (2023) Diversity and evolution of telomeric motifs and telomere DNA organization in insects. *Biological Journal of the Linnean Society* 140(4): 536–555. <https://doi.org/10.1093/biolinnean/blad068>
- Matsumoto K, Yamamoto DS, Sumitani M, Lee JM, Hatakeyama M, Oishi K (2002) Detection of a single copy on a mitotic metaphase chromosome by fluorescence in situ hybridization (FISH) in the sawfly, *Athalia rosae* (Hymenoptera). *Archive of Insect Biochemistry and Physiology* 49: 34–40. <https://doi.org/10.1002/arch.10005>
- Menezes RST, Cabral-de-Mello DC, Milani D, Bardella VB, Almeida EAB (2021) The relevance of chromosome fissions for major ribosomal DNA dispersion in hymenopteran insects. *Journal of Evolutionary Biology* 34: 1466–1476. <https://doi.org/10.1111/jeb.13909>
- Naito T (1982) Chromosome number differentiation in sawflies and its systematic implication (Hymenoptera, Tenthredinidae). *Kontyû* 50(4): 569–587.
- Rousselet J, Chaminade N, Géri C, Hewitt GM, Lemeunier F (1999) Chromosomal location of rRNA genes in *Diprion pini* detected by in situ hybridization. *Heredity* 81: 573–578. [https://doi.org/10.1016/s0764-4469\(99\)80095-x](https://doi.org/10.1016/s0764-4469(99)80095-x)
- Rousselet J, Monti L, Auger-Rozenberg M-A, Parker JS, Lemeunier F (2000) Chromosome fission associated with growth of ribosomal DNA in *Neodiprion abietis* (Hymenoptera: Diprionidae). *Proceedings of the Royal Society of London B* 267: 1819–1823. <https://doi.org/10.1098/rspb.2000.1216>
- Schwarzacher T, Heslop-Harrison P (2000) *Practical in Situ Hybridization*. Oxford, UK, BIOS Scientific Publishers Ltd, xii + 203 pp.
- Wutke S, Blank SM, Boevé J-L, Faircloth BC, Koch F, Linnen CR, Malm T, Niu G, Prous M, Schiff NM, Schmidt S, Taeger A, Vilhelmsen L, Wahlberg N, Wei M, Nyman T (2024) Phylogenomics and biogeography of sawflies and woodwasps (Hymenoptera, Symphyta). *Molecular Phylogenetics and Evolution* 199: 108144. <https://doi.org/10.1016/j.ympev.2024.108144>

ORCID

Vladimir E. Gokhman <https://orcid.org/0000-0001-9909-7559>

Valentina G. Kuznetsova <https://orcid.org/0000-0001-8386-5453>

Boris A. Anokhin <https://orcid.org/0000-0002-4110-6704>