



Conference Abstract

New trends in bioassessment of aquatic ecosystems: from organisms to DNA-based metrics

Jan Pawlowski[‡], Maria Kahlert[§]

[‡] University of Geneva, Geneva, Switzerland

[§] Swedish University of Agricultural Sciences, 750 07 Uppsala, Sweden

Corresponding author: Jan Pawlowski (jan.pawlowski@unige.ch)

Received: 02 Mar 2021 | Published: 04 Mar 2021

Citation: Pawlowski J, Kahlert M (2021) New trends in bioassessment of aquatic ecosystems: from organisms to DNA-based metrics. ARPHA Conference Abstracts 4: e65382. <https://doi.org/10.3897/aca.4.e65382>

Abstract

Traditionally, the biological quality of aquatic ecosystems is assessed using selected groups of organisms that can be identified morphologically. Recent advances in high-throughput genomic approaches offered new opportunities to monitor biodiversity and assess ecological status using DNA barcoding and metabarcoding. The DNA-based tools have been used in three different ways: (1) to replace morphological identification of biological quality elements in existing biotic indices, (2) to develop new molecular indices based on morphologically inconspicuous groups of potential environmental indicators, and (3) to predict biotic indices from environmental DNA datasets using machine learning methods (Pawlowski et al. 2018). The next steps need to take advantages and challenges of these different approaches into account in view of their future application in routine bioassessment. The Working Group 2 of DNAqua-Net, Biotic Indices & Metrics, has worked with several task forces tackling different organism groups (fish, macroinvertebrates, diatoms, bacteria, protists, meiofauna), because challenges have been shown to be quite different dependent on the target organisms Kahlert et al. 2019. For the fish the eDNA-metabarcoding methods are well developed and give very good results in terms of species detection. The important question is to see if the semi-quantitative data retrieved from the metabarcoding (proportion in eDNA sequences) could be translated to proportions in biomass/numbers that are now used in many indices. The fish researchers are trying to fit these data in, but some correction factors might be needed to correct for differences between molecular and conventional methods. Regarding the macroinvertebrates, much

discussion regarding index development was focusing on the importance of abundance measurements, and it was tested how existing indices would perform if barcoding data would be used instead of morphological data. Still discussion is ongoing on several technical issues, including the use of preservative for DNA extraction from bulk samples, the choice of primers for PCR amplification and the incompleteness of reference databases which impedes the correct assignment of eDNA sequences. Also minimum standards for routine operation are still missing. The diatom group has worked much on practical issues, starting a large initiative to compare diatom metabarcoding protocols used in routine freshwater biomonitoring for standardization (Bailet et al. 2019, Keck et al. 2018, Vasselon et al. 2017). With diatoms, all three approaches to develop molecular indices have been tested and seem promising, i.e. using existing indices with taxa names derived by matching sequences with reference databases, developing new indices based on molecular data only with traditional fixed scores, and using machine-learning techniques (Bailet et al. 2020, Vasselon et al. 2018, Tapolczai et al. 2019, Keck et al. 2018) The micro-and meiobiota group has worked towards an inclusion of microorganisms into aquatic assessment, because the microbial community dynamics are a missing link important for our understanding of rapid changes in the structure and function of aquatic ecosystems, and should be addressed in the future environmental monitoring of freshwater ecosystems (Sagova-Mareckova et al. 2021). Another focus was on how sediment DNA analysis can be integrated into stated goals of routine monitoring applications. It has been an interesting journey, and we WG2 coordinators would like to thank all the people for their engagement! Keep up the good work!

Presenting author

Jan Pawlowski

Presented at

1st DNAQUA International Conference (March 9-11, 2021)

References

- Bailet B, Bouchez A, Franc A, Frigerio J, Keck F, Karjalainen S, Rimet F, Schneider S, Kahlert M (2019) Molecular versus morphological data for benthic diatoms biomonitoring in Northern Europe freshwater and consequences for ecological status. *Metabarcoding and Metagenomics* 3 <https://doi.org/10.3897/mbmg.3.34002>
- Bailet B, Apothéloz-Perret-Gentil L, Baričević A, Chonova T, Franc A, Frigerio J, Kelly M, Mora D, Pfannkuchen M, Proft S, Ramon M, Vasselon V, Zimmermann J, Kahlert M (2020) Diatom DNA metabarcoding for ecological assessment: Comparison among bioinformatics pipelines used in six European countries reveals the need for standardization. *Science of The Total Environment* 745 <https://doi.org/10.1016/j.scitotenv.2020.140948>

- Kahlert M, Alfjorden A, Apunte-Ramos K, Baillet B, Pérez Burillo J, Carrera Gonzalez AG, Castro D, Di Bernardi C, Dully V, Fekete J, Frühe L, González R, Gratsia E, Hanjalić J, Kamberović J, Kelly A, Meriggi C, Nousiainen I, Ørberg S, Orr J, Quintana C, Papatheodoulou A, Sargac J, Shahbaz M, Tapolczai K, Tosic K, Wallin I, Zupančič M, Bohman P, Buttigieg PL, Häubner N, Leese F, Macher J, Peura S, Roslin T, Strand M, Terenius O, Vasselon V, Weigand A (2019) New molecular methods to assess biodiversity. Potentials and pitfalls of DNA metabarcoding: a workshop report. *Research Ideas and Outcomes* 5 <https://doi.org/10.3897/rio.5.e38915>
- Keck F, Vasselon V, Rimet F, Bouchez A, Kahlert M (2018) Boosting DNA metabarcoding for biomonitoring with phylogenetic estimation of operational taxonomic units' ecological profiles. *Molecular Ecology Resources* 18 (6): 1299-1309. <https://doi.org/10.1111/1755-0998.12919>
- Pawlowski J, Kelly-Quinn M, Altermatt F, Apothéloz-Perret-Gentil L, Beja P, Boggero A, Borja A, Bouchez A, Cordier T, Domaizon I, Feio MJ, Filipe AF, Fornaroli R, Graf W, Herder J, van der Hoorn B, Iwan Jones J, Sagova-Mareckova M, Moritz C, Barquín J, Piggott J, Pinna M, Rimet F, Rinkevich B, Sousa-Santos C, Specchia V, Trobajo R, Vasselon V, Vitecek S, Zimmerman J, Weigand A, Leese F, Kahlert M (2018) The future of biotic indices in the ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems. *Science of The Total Environment* 1295-1310. <https://doi.org/10.1016/j.scitotenv.2018.05.002>
- Sagova-Mareckova M, Boenigk J, Bouchez A, Cermakova K, Chonova T, Cordier T, Eisendle U, Elersek T, Fazi S, Fleituch T, Frühe L, Gajdosova M, Graupner N, Haegerbaeumer A, Kelly A-, Kopecky J, Leese F, Nöges P, Orlic S, Panksep K, Pawlowski J, Petrussek A, Piggott JJ, Rusch JC, Salis R, Schenk J, Simek K, Stovicek A, Strand DA, Vasquez MI, Vrålstad T, Zlatkovic S, Zupancic M, Stoeck T (2021) Expanding ecological assessment by integrating microorganisms into routine freshwater biomonitoring. *Water Research* 191 <https://doi.org/10.1016/j.watres.2020.116767>
- Tapolczai K, Keck F, Bouchez A, Rimet F, Kahlert M, Vasselon V (2019) Diatom DNA Metabarcoding for Biomonitoring: Strategies to Avoid Major Taxonomical and Bioinformatical Biases Limiting Molecular Indices Capacities. *Frontiers in Ecology and Evolution* 7 <https://doi.org/10.3389/fevo.2019.00409>
- Vasselon V, Rimet F, Tapolczai K, Bouchez A (2017) Assessing ecological status with diatoms DNA metabarcoding: Scaling-up on a WFD monitoring network (Mayotte island, France). *Ecological Indicators* 82: 1-12. <https://doi.org/10.1016/j.ecolind.2017.06.024>
- Vasselon V, Bouchez A, Rimet F, Jacquet S, Trobajo R, Corniquel M, Tapolczai K, Domaizon I (2018) Avoiding quantification bias in metabarcoding: Application of a cell biovolume correction factor in diatom molecular biomonitoring. *Methods in Ecology and Evolution* 9 (4): 1060-1069. <https://doi.org/10.1111/2041-210x.12960>