



Conference Abstract

# Nitrate-Reducing Fe(II)-Oxidizing microorganisms: linking Fe, C and N Cycles in subsurface environments

Cristina Escudero Parada<sup>‡</sup>, Andreas Kappler<sup>‡</sup>

<sup>‡</sup> University of Tuebingen, Tuebingen, Germany

Corresponding author: Cristina Escudero Parada ([c.escudero@uni-tuebingen.de](mailto:c.escudero@uni-tuebingen.de))

Received: 16 Jun 2023 | Published: 13 Oct 2023

Citation: Escudero Parada C, Kappler A (2023) Nitrate-Reducing Fe(II)-Oxidizing microorganisms: linking Fe, C and N Cycles in subsurface environments. ARPHA Conference Abstracts 6: e108089.

<https://doi.org/10.3897/aca.6.e108089>

## Abstract

The discovery of the Nitrate-Reducing Fe(II)-Oxidizing (NRFeOx) microbial metabolism, which couples the oxidation of Fe(II) to the reduction of nitrate ( $\text{NO}_3^-$ ) using organic matter or carbon dioxide ( $\text{CO}_2$ ) as carbon source, was a major milestone in microbial ecology (Straub et al. 1996). NRFeOx microorganisms play an essential role on a global scale in three of the most important biogeochemical cycles: iron (Fe), carbon (C) and nitrogen (N) (Kappler et al. 2021, Huang et al. 2021). In addition, these organisms participate in the mobilization or stabilization of organic carbon, as well as in  $\text{CO}_2$  fixation, thus contributing to the reduction of atmospheric  $\text{CO}_2$  (Kappler et al. 2021). Finally, the activity of these microorganisms is key to remove the pollutant  $\text{NO}_3^-$  from aquifers, which is one of the major worldwide environmental issues since many environments exceed the maximum regulatory concentration ( $50 \text{ mg L}^{-1}$ ) (Kazakis et al. 2020).

A plethora of NRFeOx microorganisms have been described in the last decades. However, most of these microorganisms have been reclassified as chemodenitrifiers. That is to say, Fe(II) is not enzymatically oxidized but indirectly by the reactive nitrogen species produced during denitrification (Fig. 1). In fact, only in three cultures so far, named KS, BP and AG, has the presence of true NRFeOx metabolism been unequivocally demonstrated (Straub et al. 1996, Huang et al. 2021b, Jakus et al. 2021b).




Figure 1. [doi](#)

Overview of the three different types of interaction between nitrate-reducing bacteria and Fe(II). (A) Autotrophic NRFeOx obtain carbon from  $\text{CO}_2$  and oxidize Fe(II) enzymatically. (B) Mixotrophic NRFeOx require additional organic carbon as a carbon source, and Fe(II) oxidation has some enzymatic component (although there may also be some abiotic component). (C) Chemodenitrifiers require organic carbon and have no enzymatic component of Fe(II) oxidation. The position of the minerals (orange) relative to cells (black) indicates whether or not cell encrustation is expected. Image from Bryce et al. (2018).

Cultures KS, BP and AG have been studied thoroughly in the past years, analyzing the rate and mechanism by which these communities carry out autotrophic NRFeOx. Different omics studies have revealed that cultures KS, BP and AG consist of a mixture of bacterial species, which collaborate in order to grow under autotrophic NRFeOx conditions. Each culture is dominated by a novel candidate species of the genus *Ferrigenium* (Huang et al. 2022) capable of fixing  $\text{CO}_2$  and oxidizing Fe(II), but which requires flanking species to complete denitrification (Huang et al. 2021b, He et al. 2016, Huang et al. 2021a).

Interestingly, these communities not only carry out NRFeOx using dissolved Fe(II) as energy source (Straub et al. 1996, Huang et al. 2021b, Jakus et al. 2021b), but they can also oxidize Fe(II) minerals, the main form in which Fe(II) can be found in the Earth's crust (Huang et al. 2021). In fact, Fe(II)-bearing minerals are thought to be the main drivers of  $\text{NO}_3^-$  reduction in subterranean environments (Huang et al. 2021), which has additional ecological consequences. NRFeOx microorganisms can trigger the turnover of the Fe(II)-bearing minerals, resulting in the mobilization of mineral structural elements such as S, P, C or contaminant heavy metals and the precipitation of Fe(III) minerals at circumneutral pH (Weber et al. 2001, Jakus et al. 2021a).

Here, we will present a review of the insights learned from the three NRFeOx autotrophic cultures and discuss their ecological role, their importance in biogeochemical cycles, and their potential biotechnological applications.

## Keywords

geomicrobiology, biogeochemical cycles, bioremediation

## Presenting author

Cristina Escudero Parada

## Presented at

ISEB-ISSM 2023

## Funding program

Teach@Tübingen program

## Hosting institution

Eberhard-Karls-University of Tübingen

## Conflicts of interest

The authors have declared that no competing interests exist.

## References

- Bryce C, Blackwell N, Schmidt C, Otte J, Huang Y, Kleindienst S, Tomaszewski E, Schad M, Warter V, Peng C, Byrne J, Kappler A (2018) Microbial anaerobic Fe(II) oxidation - Ecology, mechanisms and environmental implications. *Environmental Microbiology* 20 (10): 3462-3483. <https://doi.org/10.1111/1462-2920.14328>
- He S, Tominski C, Kappler A, Behrens S, Roden E (2016) Metagenomic Analyses of the Autotrophic Fe(II)-Oxidizing, Nitrate-Reducing Enrichment Culture KS. *Applied and Environmental Microbiology* 82 (9): 2656-2668. <https://doi.org/10.1128/aem.03493-15>
- Huang J, Jones A, Waite TD, Chen Y, Huang X, Rosso K, Kappler A, Mansor M, Tratnyek P, Zhang H (2021) Fe(II) Redox Chemistry in the Environment. *Chemical Reviews* 121 (13): 8161-8233. <https://doi.org/10.1021/acs.chemrev.0c01286>
- Huang Y, Straub D, Blackwell N, Kappler A, Kleindienst S (2021a) Meta-omics Reveal *Gallionellaceae* and *Rhodanobacter* Species as Interdependent Key Players for Fe(II) Oxidation and Nitrate Reduction in the Autotrophic Enrichment Culture KS. *Applied and Environmental Microbiology* 87 (15). <https://doi.org/10.1128/aem.00496-21>
- Huang Y, Straub D, Kappler A, Smith N, Blackwell N, Kleindienst S (2021b) A Novel Enrichment Culture Highlights Core Features of Microbial Networks Contributing to Autotrophic Fe(II) Oxidation Coupled to Nitrate Reduction. *Microbial Physiology* 31 (3): 280-295. <https://doi.org/10.1159/000517083>
- Huang Y, Jakus N, Straub D, Konstantinidis K, Blackwell N, Kappler A, Kleindienst S (2022) 'Candidatus ferrigenium straubiae' sp. nov., 'Candidatus ferrigenium bremense'

sp. nov., 'Candidatus ferrigenium altingense' sp. nov., are autotrophic Fe(II)-oxidizing bacteria of the family Gallionellaceae. *Systematic and Applied Microbiology* 45 (3). <https://doi.org/10.1016/j.syapm.2022.126306>

- Jakus N, Mellage A, Hörschen C, Maisch M, Byrne J, Mueller C, Grathwohl P, Kappler A (2021a) Anaerobic Neutrophilic Pyrite Oxidation by a Chemolithoautotrophic Nitrate-Reducing Iron(II)-Oxidizing Culture Enriched from a Fractured Aquifer. *Environmental Science & Technology* 55 (14): 9876-9884. <https://doi.org/10.1021/acs.est.1c02049>
- Jakus N, Blackwell N, Osenbrück K, Straub D, Byrne J, Wang Z, Glöckler D, Elsner M, Lueders T, Grathwohl P, Kleindienst S, Kappler A (2021b) Nitrate Removal by a Novel Lithoautotrophic Nitrate-Reducing, Iron(II)-Oxidizing Culture Enriched from a Pyrite-Rich Limestone Aquifer. *Applied and Environmental Microbiology* 87 (16). <https://doi.org/10.1128/aem.00460-21>
- Kappler A, Bryce C, Mansor M, Lueder U, Byrne J, Swanner E (2021) An evolving view on biogeochemical cycling of iron. *Nature Reviews Microbiology* 19 (6): 360-374. <https://doi.org/10.1038/s41579-020-00502-7>
- Kazakis N, Matiatos I, Ntona M, Bannenberg M, Kalaitzidou K, Kaprara E, Mitrakas M, Ioannidou A, Vargemezis G, Voudouris K (2020) Origin, implications and management strategies for nitrate pollution in surface and ground waters of Anthemountas basin based on a  $\delta^{15}\text{N-NO}_3^-$  and  $\delta^{18}\text{O-NO}_3^-$  isotope approach. *Science of The Total Environment* 724 <https://doi.org/10.1016/j.scitotenv.2020.138211>
- Straub KL, Benz M, Schink B, Widdel F (1996) Anaerobic, nitrate-dependent microbial oxidation of ferrous iron. *Applied and Environmental Microbiology* 62 (4): 1458-1460. <https://doi.org/10.1128/aem.62.4.1458-1460.1996>
- Weber K, Picardal F, Roden E (2001) Microbially Catalyzed Nitrate-Dependent Oxidation of Biogenic Solid-Phase Fe(II) Compounds. *Environmental Science & Technology* 35 (8): 1644-1650. <https://doi.org/10.1021/es0016598>