



Conference Abstract

Genetic assessment of the Forest Dormouse *Dryomys nitedula* (Pallas, 1773) in Latvia

Dainis Ruņģis[‡], Inese Gavarāne[§], Linda Bankovska[§], Valdis Pilāts[|], Zanda Segliņa[|], Digna Pilāte[‡]

[‡] Latvian State Forest Research Institute "Silava", Salaspils, Latvia

[§] Daugavpils University, Daugavpils, Latvia

[|] Nature Conservation Agency, Sigulda, Latvia

Corresponding author: Zanda Segliņa (zanda.seglina@daba.gov.lv)

Received: 06 Apr 2022 | Published: 15 Apr 2022

Citation: Ruņģis D, Gavarāne I, Bankovska L, Pilāts V, Segliņa Z, Pilāte D (2022) Genetic assessment of the Forest Dormouse *Dryomys nitedula* (Pallas, 1773) in Latvia. ARPHA Conference Abstracts 5: e84997.

<https://doi.org/10.3897/aca.5.e84997>

Abstract

The Latvian population of the Forest Dormouse is small and isolated, situated far from the main distribution range of this species. The population is located in the south-eastern region of Latvia, close to the Belarusian border. Sixty-three nest-boxes were located throughout this distribution area and captured forest dormouse individuals were sampled for genetic analysis. Animal age, sex, and coordinates from place of capture were recorded. DNA was extracted from 120 individuals using the QIAamp DNA Mini Kit (Qiagen, Germany). RAPD-PCR polymorphisms at 65 loci were used to examine the genetic diversity and differentiation of the Forest Dormouse population in Latvia. 89% of loci were polymorphic, average expected heterozygosity over all loci was 0.321, average polymorphism information content (PIC) was 0.246. The program 'Structure' was used to perform Bayesian assignment of individuals to a predefined number of clusters. The deltaK method identified that the most likely number of clusters within the Latvian Forest Dormouse population was three. Assignment of individuals to a particular cluster mostly corresponded with the location where the individual was captured. Analysis of molecular variance (AMOVA) indicated that there was significant genetic differentiation between the three clusters with pairwise F_{st} (a modification of F_{st}) values ranging from 0.118 to 0.180.

Keywords

Forest Dormouse, genetic, clusters

Presenting author

Zanda Segliņa

Presented at

Poster presentation at the 11th International Dormice Conference (May 9-13, 2022)