

Invasive alien plants in Sergipe, north-eastern Brazil

Juliano Ricardo Fabricante¹, Kelianne Carolina Targino de Araújo¹,
Thieres Santos Almeida¹, João Paulo Bispo Santos², Daniel Oliveira Reis¹

1 *Laboratório de Ecologia e Conservação da Biodiversidade, Departamento de Biociências, Universidade Federal de Sergipe, Campus Universitário Prof. Alberto Carvalho, 49.510–200, Itabaiana, SE, Brazil*

2 *Instituto de Biologia, Universidade Estadual de Campinas, 13.056–405, Campinas, SP, Brazil*

Corresponding author: João Paulo Bispo Santos (joaobiotanica@hotmail.com)

Academic editor: A.M. Leal-Zanchet | Received 12 July 2020 | Accepted 8 January 2021 | Published 29 January 2021

Citation: Fabricante JR, de Araújo KCT, Almeida TS, Santos JPB, Reis DO (2021) Invasive alien plants in Sergipe, north-eastern Brazil. *Neotropical Biology and Conservation* 16(1): 89–104. <https://doi.org/10.3897/neotropical.16.e56427>

Abstract

Biological invasions are considered one of the greatest threats to global biodiversity. In addition, they cause substantial economic impacts. However, studies about the subject in Brazil are still scarce. The aim of the present study was to prepare an inventory of non-native flora with invasive potential from Sergipe, Brazil. The inventory was carried out along the entire length of the sites. The species with potential invaders were grouped according to the biome/ecosystem and classified according to their habit and origin. Eighty-five species with invasive potential were sampled, 43 in the Caatinga, 75 in the Atlantic Forest, 36 in Sandbank and 22 in Mangrove. From these species, 17 were inventoried in all the biomes/ecosystems and 36 were observed in only one of them, six in the Caatinga, 27 in the Atlantic Forest and three in Sandbank. The number of potentially-invasive species sampled in Sergipe is alarming. The present study showed nearly twice the species listed by other authors for the entire northeast Brazil. This high number of taxa may be a consequence of facilitating the transfer of these species and the conservation conditions of the ecosystems studied in Sergipe. Another very worrying factor is that many of the species sampled are extremely aggressive and cause a series of impacts.

Keywords

Checklist, environmental impacts, floristic survey, invasive exotic

Introduction

Biological Invasions (BI) are considered one of the greatest threats to global biodiversity (Williamson 1996; Westbrooks 1998; IUCN 1999; Ziller 2001; Andrade et al. 2010; Zenni and Ziller 2011; Fabricante and Siqueira-Filho 2012). The BIs occur when a non-native species is able to adapt, reproduce and disperse to different locations in the area of initial introduction, establishing and invading new regions (Richardson et al. 2000; Moro et al. 2012). The invasive exotic species have the ability to alter the structure of communities, inhibit the regeneration of native species, cause interference in the nutrient cycling, water balance, productivity, succession processes and biological diversity (Pujadas 2001; Lockwood et al. 2007; Ziller and Zalba 2007; Andrade et al. 2009; Oliveira-Costa and Souza 2015; Santos and Calafate 2018).

There are some hypotheses that attempt to explain the success of invasive exotic species. One of them is that degraded environments are facilitators of BI (Williamson 1996). In addition, the production of large amounts of propagules, the rapid growth of the root system, adaptation to fire, the ability to interfere with the growth of neighbouring plants, the morphological and physiological similarity with native species, self-pollination and the ability to fix nitrogen by symbiotic bacteria, are characteristics that should contribute to the success of BI processes (Cronk and Fuller 1995). The absence of natural enemies is also another contributing factor, since exotic organisms, in general, are free from predators, competitors and parasites from their region of origin (Ziller 2001).

Knowing the threats that BI can cause to biodiversity and to productive sectors of society, the topic was discussed globally for the first time at ECO-Rio 92 (CBD 1992; ONUBR 2020). Years later, the Aichi Goals were created, where it was established by Goal 9 that, by 2020, the signatory countries of the Agreement, should establish measures to prevent the introduction of exotic invaders and that already introduced species should be identified, prioritised and controlled or eradicated (CBD 2020).

Following this movement, in 1996, the Inter-American Biodiversity Information Network (IABIN) was created, which comprises 18 signatory countries, including Brazil (I3N 2020). In addition to the online database that provides a list of its invasive exotic species, here in Brazil, the first federative document directed to the topic was created in 2009. Resolution No. 5 of the National Biodiversity Commission (Brasil 2009), updated in 2018 by Resolution No. 7 (Brasil 2018), outlines the National Strategy on Invasive Exotic Species. The document aims to “Guide the implementation of measures to prevent the introduction and dispersion and significantly reduce the impact of invasive exotic species on Brazilian biodiversity and ecosystem services, to control or eradicate invasive alien species” (Brasil 2018). This document is divided into six components: (1) Legislation, intersectoral articulation and international cooperation; (2) Prevention, early detection and rapid response; (3) Eradication, control and mitigation of impacts; (4) Scientific research; (5) Technical training; (6) Environmental Education and Communication.

According to Objective 2.1 of component 2, it is necessary to update the survey and identify the invasive exotic species present in the country (Brasil 2018). Still, according to this objective, it is only after the identification of these species that it is possible to define preventative measures against them. Thus, inventories of these taxa should be performed in order to identify their occurrence in different ecosystems, since these can not only provide subsidies for projects aimed at prevention of environmental impacts caused by exotic species, but also for conservation plans (Ziller 2016). Despite the evident importance of exotic species lists, only Paraná, Santa Catarina, Rio Grande do Sul and São Paulo published official lists (MMA 2020). However, it is worth noting the growth in research related to the theme in recent years (Ziller 2001; Nunes et al. 2018).

In front of the impacts that these taxa can cause and the absence of a species list for Sergipe, the present study sought to answer the following questions: I. What are the non-native species with potential invasion that occur in Sergipe; II. What is the habit of these species; III. What is the origin of these species; and IV. In which biomes/ecosystems can they be found? Thus, this work aimed to carry out the floristic survey of non-native plants with potential for invasion to Sergipe, Brazil.

Material and methods

The State of Sergipe, located in north-eastern Brazil, has an area of 21,910 km² (Prata et al. 2013). Its vegetation is composed of Atlantic Forest and Caatinga biomes, both intensely fragmented and degraded (Ferrari 2005; Prata et al. 2013). The State has three climatic types according to the Köppen-Geiger classification: tropical with dry summer (As) in northern coast and central State, tropical monsoon (Am) in the south coast and semi-arid (BSh) in the west (Alvares et al. 2013).

For this study, 100 expeditions were carried out to sites under the domains of Caatinga, Atlantic Forest, Sandbank and Mangrove in 23 municipalities in the State of Sergipe (Fig. 1). The average duration of each expedition was 4 hours. The studied sites are represented by conservation units (PARNA Serra de Itaiana, APA Morro do Urubu, APA Litoral Sul, APA Litoral Norte and MONA Grota do Angico), fragments of native properties vegetation agricultural environments (areas of cultivation and pastures), vacant lots and margins of highways and roads.

The species inventory was carried out by means of walks along the entire extension of the places (active search) where samples of fertile material from the plants were collected according to Mori et al. (1989). After collection, this material was herborised and deposited at the Herbarium ASE (Thiers 2020) of the Federal University of Sergipe, São Cristóvão, SE. With this, each plant deposited in the herbarium ASE was registered by receiving a voucher. The taxonomic categories are in accordance with APG IV (2016) and the spelling of the authors' names according to the IPNI – International Plant Names Index (2020).




Figure 1. Municipalities where expeditions were carried out to survey plant species with potential to invade the State of Sergipe, Brazil. 1 – Aracaju; 2 – Areia Branca; 3 – Barra dos Coqueiros; 4 – Campo do Brito; 5 – Canindé de São Francisco; 6 – Capela; 7 – Carira; 8 – Estância; 9 – Frei Paulo; 10 – Indiaroba; 11 – Itabaiana; 12 – Itaporanga D’ajuda; 13 – Lagarto; 14 – Laranjeiras; 15 – Macambira; 16 – Malhador; 17 – Nossa Senhora da Glória; 18 – Pacatuba; 19 – Pirambu; 20 – Propriá; 21 – Ribeirópolis; 22 – São Cristóvão; 23 – São Domingos.

The habit (herbs, lianas, palms, shrub and trees) of the plants was defined through observations made in the field and by consulting Flora do Brasil (2020). The origins of the inventoried species were determined by consulting specialised databases (BioNET-EAFRINET 2020; CABI 2020; I3N 2020; ISSG 2020) and scientific articles.

With the list of species, those categorised as non-native, were separated and those with invasive potential (naturalised and exotic invaders) were used to prepare the present study. The definition of naturalised and exotic invasive species used here is based on the work of Richardson et al. (2000), where they consider non-native species that reproduce and form stable populations only in the place where they were introduced and exotic invaders, those that reproduce, disperse and establish themselves in places far from the place of introduction. The classification of the species in naturalised and exotic was made by in situ observations and by consulting the specialised database (BioNET-EAFRINET 2020; CABI 2020; I3N 2020; ISSG 2020). Bar graphs were made using “ggplot2” package (Wickham 2016) and the Venn Dia-

gram was created using “VennDiagram” package (Chen and Boutros 2011), both figures being made using the software R (R Core Team 2020). The data underpinning the analysis reported in this paper are deposited at GBIF, the Global Biodiversity Information Facility, and are available at <https://doi.org/10.15468/6mdt5d>.

Results

Altogether, 83 non-native species with invasion potential were sampled in the State of Sergipe, of which 73 were found in the Atlantic Forest domains, 43 in the Caatinga, 36 in Sandbank and 22 in the Mangrove (Table 1). From these species, 17 were inventoried in all the biomes/ecosystems studied and 35 were observed in only one of them, 26 in the Atlantic Forest, six in the Caatinga and three in Sandbank (Fig. 2).

The sampled species belonged to 29 botanical families. The families with the highest number of records were Poaceae with 20 species (23.4%), followed by Fabaceae with six (7.1%), Asparagaceae, Cucurbitaceae and Myrtaceae with five (5.9%), Apocynaceae and Asteraceae with four (4.7%). The other families presented three, two or just one recorded species (Fig. 3B).

Regarding the origin of the species, 22 were African species, 12 Asian and 15 species originated in the continents of Africa and Asia, six species originated in North America, while Central America, South America and Oceania presented only three species each. In addition, 19 other species had two or more continents as their origin (Fig. 3B). Regarding the habit of species, the most abundant group was the herbs with 43 (51.2%) representatives, followed by trees with 18 (21.42%), shrubs with 12 (14.28%), lianas with nine (10.71%) and palm trees with only one species (1.19%) (Fig. 3C; Table 2).




Figure 2. Venn Diagram showing the number of invasive alien species shared amongst the biomes/ecosystems of the State of Sergipe in northeast Brazil.

Table 1. List of alien species sampled by biome/ecosystem in Sergipe, Brazil. 0: absence; 1: presence. Af = Atlantic Forest; Ca = Caatinga; Ma = Mangrove; Sa = Sandbank. Vouchers of herbarium ASE, Federal University of Sergipe, São Cristóvão, SE. * = Species not collected due to lack of fertile material.

Family/Species	Ca	Af	Sa	Ma	Voucher
Acanthaceae Juss.					
<i>Asystasia gangetica</i> (L.) T. Anderson	0	1	0	1	37616
<i>Dyschoriste depressa</i> Nees	0	1	0	0	39688
Amaranthaceae A. Juss.					
<i>Amaranthus viridis</i> L.	1	1	1	1	37621
<i>Chenopodium murale</i> L.	1	1	0	0	*
Anacardiaceae R. Br.					
<i>Mangifera indica</i> L.	0	1	1	0	36980
Apocynaceae Juss.					
<i>Calotropis procera</i> (Aiton) W.T. Aiton	1	0	1	0	36968
<i>Catharanthus roseus</i> (L.) G. Don	1	1	0	0	41924
<i>Cryptostegia madagascariensis</i> Bojer	1	1	1	0	39879
<i>Thevetia peruviana</i> (Pers.) K. Schum.	0	1	0	0	37380
Araceae Juss.					
<i>Dieffenbachia seguine</i> (Jacq.) Schott	0	1	0	0	*
<i>Syngonium angustatum</i> Schott	0	1	0	0	37382
<i>Xanthosoma sagittifolium</i> (L.) Schott	0	1	0	0	37373
Araceae Schultz Sch.					
<i>Elaeis guineensis</i> Jacq.	0	1	0	0	*
Asparagaceae Juss.					
<i>Agave angustifolia</i> Haw.	1	1	0	0	*
<i>Agave sisalana</i> Perrine ex Engelm.	1	1	1	0	38464
<i>Dracaena fragrans</i> (L.) Ker Gawl.	0	1	0	0	37376
<i>Furcraea foetida</i> (L.) Haw.	1	0	0	0	*
<i>Sansevieria trifasciata</i> Prain	0	1	0	0	37374
Asteraceae Bercht. & J. Presl					
<i>Cosmos caudatus</i> Kunth	0	1	0	0	*
<i>Cosmos sulphureus</i> Cav.	0	1	0	0	*
<i>Sonchus asper</i> (L.) Hill	0	1	0	0	*
<i>Tithonia diversifolia</i> (Hemsl.) A. Gray	0	1	0	0	*
Cactaceae Juss.					
<i>Opuntia stricta</i> (Haw.) Haw.	0	0	1	0	37623
Casuarinaceae R.Br					
<i>Casuarina equisetifolia</i> L.	0	1	1	0	36982
Combretaceae R.Br.					
<i>Terminalia catappa</i> L.	1	1	1	1	37617
Commelinaceae Mirb.					
<i>Commelina benghalensis</i> L.	1	1	1	1	36971
<i>Tradescantia zebrina</i> Heynh. ex Bosse	0	1	0	0	*
Convolvulaceae Juss.					
<i>Ipomoea nil</i> (L.) Roth	0	1	0	0	39694
Crassulaceae J.St.-Hil.					
<i>Kalanchoe delagoensis</i> Eckl. & Zeyh.	0	1	0	0	*
<i>Kalanchoe mortagei</i> Raym. -Hamet & H. Perrier	0	1	0	0	*
<i>Kalanchoe pinnata</i> (Lam.) Pers.	0	1	0	0	*
Cucurbitaceae Juss.					
<i>Citrullus lanatus</i> (Thunb.) Matsum. & Nakai	1	1	0	0	*
<i>Cucumis anguria</i> L.	1	1	0	0	*
<i>Luffa cylindrica</i> (L.) M. Roem.	1	1	1	0	38450
<i>Momordica charantia</i> L.	1	1	1	1	36979
Cyperaceae Juss.					
<i>Cyperus rotundus</i> L.	1	1	1	1	36975
Euphorbiaceae Juss.					
<i>Euphorbia tirucalli</i> L.	1	0	0	0	*
<i>Jatropha gossypifolia</i> L.	1	1	1	1	37622

Family/Species	Ca	Af	Sa	Ma	Voucher
<i>Ricinus communis</i> L.	1	1	1	1	37619
Fabaceae Lindl.					
<i>Acacia mangium</i> Willd.	0	1	1	0	*
<i>Crotalaria pallida</i> Aiton	0	1	1	0	*
<i>Crotalaria retusa</i> L.	0	1	1	0	36999
<i>Leucaena leucocephala</i> (Lam.) de Wit	1	1	1	1	37618
<i>Pithecellobium dulce</i> (Roxb.) Benth.	0	1	1	1	39699
<i>Prosopis</i> spp.	1	0	0	0	36978
Lamiaceae Martinov					
<i>Leonotis nepetifolia</i> (L.) R. Br.	1	1	1	0	*
Malvaceae Juss.					
<i>Talipariti tiliaceum</i> (L.) Fryxell	0	0	1	0	39807
<i>Thespesia populnea</i> (L.) Sol. ex Corrêa	0	1	0	1	36977
Meliaceae A. Juss.					
<i>Azadirachta indica</i> A. Juss.	1	1	1	1	37620
Moraceae Gaudich.					
<i>Artocarpus heterophyllus</i> Lam.	0	1	0	0	*
Myrtaceae Juss.					
<i>Eucalyptus</i> spp.	0	1	0	0	*
<i>Psidium guajava</i> L.	0	1	1	0	*
<i>Syzygium cumini</i> (L.) Skeels	0	1	0	1	38454
<i>Syzygium jambos</i> (L.) Alston	0	1	0	0	39885
<i>Syzygium malaccense</i> (L.) Merr. & L.M. Perry	0	1	0	0	*
Nyctaginaceae Juss.					
<i>Boerhavia diffusa</i> L.	1	1	1	1	36974
<i>Mirabilis jalapa</i> L.	0	1	0	0	*
Papaveraceae Juss.					
<i>Argemone mexicana</i> L.	1	1	0	0	*
Poaceae Barnhart					
<i>Aristida adscensionis</i> L.	1	0	0	0	38453
<i>Arundo donax</i> L.	0	1	1	1	*
<i>Cenchrus ciliaris</i> L.	1	1	0	0	*
<i>Cenchrus echinatus</i> L.	1	1	1	1	36969
<i>Cenchrus polystachios</i> (L.) Morrone	1	1	0	0	*
<i>Cenchrus purpureus</i> (Schumach.) Morrone	0	1	0	0	*
<i>Cynodon dactylon</i> (L.) Pers.	1	1	1	1	37624
<i>Dactyloctenium aegyptium</i> (L.) Willd.	1	1	1	1	36973
<i>Echinochloa colona</i> (L.) Link	1	1	0	0	41904
<i>Eleusine indica</i> (L.) Gaertn.	1	1	1	0	41940
<i>Eragrostis ciliaris</i> (L.) R. Br.	1	1	1	0	37630
<i>Eragrostis pilosa</i> (L.) P. Beauv.	1	1	1	1	41908
<i>Eragrostis tenella</i> (L.) P. Beauv. ex Roem. & Schult.	1	1	1	1	36981
<i>Megathyrsus maximus</i> (Jacq.) B.K. Simon & S.W.L. Jacobs	1	1	1	1	39883
<i>Melinis minutiflora</i> P. Beauv.	0	1	0	0	42075
<i>Melinis repens</i> (Willd.) Zizka	1	1	1	1	39703
<i>Sorghum bicolor</i> subsp. <i>arundinaceum</i> (Desv.) de Wet & J.R. Harlan	1	1	1	0	*
<i>Tragus berteronianus</i> Schult.	1	0	0	0	*
<i>Urochloa brizantha</i> (Hochst. ex A. Rich.) R.D. Webster	0	1	0	0	37629
<i>Urochloa mosambicensis</i> (Hack.) Dandy	1	1	0	0	37628
Polygonaceae A. Juss.					
<i>Antigonon leptopus</i> Hook. & Arn.	0	1	0	0	39695
Rubiaceae Juss.					
<i>Morinda citrifolia</i> L.	0	0	1	0	*
Solanaceae A. Juss.					
<i>Nicandra physalodes</i> (L.) Gaertn.	1	1	0	0	41880
<i>Nicotiana glauca</i> Graham	1	0	0	0	42083
<i>Physalis angulata</i> L.	1	1	0	0	36997
Total	43	73	36	22	




Figure 3. Records of invasive exotic species in the State of Sergipe, Brazil: number of species per family (A), number of species by origin (B), number of species recorded for each habit (C). In (A), the term “others” indicates families with only one recorded species; in (C), “other” indicates that the species has two or more continents as origin (see Table 2).

Table 2. Habit and origin of alien species sampled in Sergipe, Brazil. H = herb; L = Liana; P = palm; S = shrub; T = tree.

Specie	Habit	Origin
<i>Acacia mangium</i> Willd.	T	Africa; Asia
<i>Agave angustifolia</i> Haw.	H	Mexico
<i>Agave sisalana</i> Perrine ex Engelm.	H	Mexico
<i>Amaranthus viridis</i> L.	H	Central America
<i>Antigonon leptopus</i> Hook. & Arn.	L	Mexico
<i>Argemone mexicana</i> L.	H	Mexico; Central America
<i>Aristida adscensionis</i> L.	H	Canary Islands
<i>Artocarpus heterophyllus</i> Lam.	T	India; Malaysia
<i>Arundo donax</i> L.	H	Asia
<i>Asystasia gangetica</i> (L.) T. Anderson	L	India; Malaysia; Asia; Africa
<i>Azadirachta indica</i> A. Juss.	T	India
<i>Boerhavia diffusa</i> L.	H	India
<i>Calotropis procera</i> (Aiton) W.T. Aiton	S	Africa; Asia
<i>Casuarina equisetifolia</i> L.	T	Asia
<i>Catharanthus roseus</i> (L.) G. Don	H	Madagascar
<i>Cenchrus ciliaris</i> L.	H	Africa; Asia
<i>Cenchrus echinatus</i> L.	H	Central America
<i>Cenchrus polystachios</i> (L.) Morrone	H	Africa; Asia
<i>Cenchrus purpureus</i> (Schumach.) Morrone	H	Africa
<i>Chenopodium murale</i> L.	H	Europa; Asia; Africa
<i>Citrullus lanatus</i> (Thunb.) Matsum. & Nakai	L	Africa
<i>Commelina benghalensis</i> L.	H	Africa; Asia
<i>Cosmos caudatus</i> Kunth	H	Central America

Specie	Habit	Origin
<i>Cosmos sulphureus</i> Cav.	H	Madagascar
<i>Crotalaria pallida</i> Aiton	S	Africa
<i>Crotalaria retusa</i> L.	S	Africa; Asia
<i>Cryptostegia madagascariensis</i> Bojer	L	Madagascar
<i>Cucumis anguria</i> L.	L	Africa
<i>Cynodon dactylon</i> (L.) Pers.	H	Africa; Asia; Europa
<i>Cyperus rotundus</i> L.	H	India; Africa
<i>Dactyloctenium aegyptium</i> (L.) Willd.	H	Africa; Asia
<i>Dieffenbachia seguine</i> (Jacq.) Schott	S	Americas
<i>Dracaena fragrans</i> (L.) Ker Gawl.	S	Western Africa
<i>Dyschoriste depressa</i> Nees	S	South Africa
<i>Echinochloa colona</i> (L.) Link	H	Asia; Europa
<i>Elaeis guineensis</i> Jacq.	P	Africa
<i>Eleusine indica</i> (L.) Gaertn.	H	Africa; Asia
<i>Eragrostis ciliaris</i> (L.) R. Br.	H	Africa
<i>Eragrostis pilosa</i> (L.) P. Beauv.	H	Europa; Africa
<i>Eragrostis tenella</i> (L.) P. Beauv. ex Roem. & Schult	H	Africa; Asia
<i>Eucalyptus</i> spp.	T	Australia
<i>Euphorbia tirucalli</i> L.	S	Africa; India
<i>Furcraea foetida</i> (L.) Haw.	H	Central America; North of South America
<i>Ipomoea nil</i> (L.) Roth.	L	Asia
<i>Jatropha gossypifolia</i> L.	S	Tropical America
<i>Kalanchoe delagoensis</i> Eckl. & Zeyh.	H	Madagascar
<i>Kalanchoe mortagei</i> Raym. -Hamet & H. Perrier	H	Madagascar
<i>Kalanchoe pinnata</i> (Lam.) Pers.	H	Africa
<i>Leonotis nepetifolia</i> (L.) R. Br.	H	Africa; Asia
<i>Leucaena leucocephala</i> (Lam.) de Wit	T	Mexico
<i>Luffa cylindrica</i> (L.) M.Roem.	L	Africa; Asia
<i>Mangifera indica</i> L.	T	India; Asia
<i>Megathyrsus maximus</i> (Jacq.) B.K. Simon & S.W.L. Jacobs	H	Africa
<i>Melinis minutiflora</i> P. Beauv.	H	Africa
<i>Melinis repens</i> (Willd.) Zizka	H	Africa
<i>Mirabilis jalapa</i> L.	S	Tropical America
<i>Momordica charantia</i> L.	L	Asia
<i>Morinda citrifolia</i> L.	T	Asia; Australia
<i>Nicandra physalodes</i> (L.) Gaertn.	H	Peru
<i>Nicotiana glauca</i> Graham	S	Argentina; Bolivia
<i>Opuntia stricta</i> (Haw.) Haw.	S	Central America
<i>Physalis angulata</i> L.	H	Australia; Tropical America
<i>Pithecellobium dulce</i> (Roxb.) Benth.	T	Mexico
<i>Prosopis</i> spp.	T	Americas
<i>Psidium guajava</i> L.	T	North America
<i>Ricinus communis</i> L.	S	Africa; Asia
<i>Sansevieria trifasciata</i> Prain	H	Africa
<i>Sonchus asper</i> (L.) Hill	H	Oceania
<i>Sorghum bicolor</i> subsp. <i>arundinaceum</i> (Desv.) de Wet & J.R. Harlan	H	Africa
<i>Syngonium angustatum</i> Schott	L	Central America
<i>Syzygium cumini</i> (L.) Skeels	T	Asia
<i>Syzygium jambos</i> (L.) Alston	T	Asia
<i>Syzygium malaccense</i> (L.) Merr. & L.M. Perry	T	Asia
<i>Talipariti tiliaceum</i> (L.) Fryxell	T	Polynesia
<i>Terminalia catappa</i> L.	T	India; Malaysia; Asia
<i>Thespesia populnea</i> (L.) Sol. ex Corrêa	T	Asia
<i>Thevetia peruviana</i> (Pers.) K. Schum.	T	Central and North America; India
<i>Tithonia diversifolia</i> (Hemsl.) A. Gray	H	Mexico; Central America
<i>Tradescantia zebrina</i> Heynh. ex Bosse	H	Mexico; Central America
<i>Tragus berteronianus</i> Schult.	H	Asia; Africa
<i>Urochloa brizantha</i> (Hochst. ex A. Rich.) R.D. Webster	H	Africa
<i>Urochloa mosambicensis</i> (Hack.) Dandy	H	Africa
<i>Xanthosoma sagittifolium</i> (L.) Schott	H	South America

Discussion

The number of plant species with invasive potential sampled in Sergipe is alarming, because it exceeds the values found in states with much larger territories, such as Santa Catarina (39 spp.), Rio Grande do Sul (44 spp.) and Paraná (72 spp.) (Brasil 2013; Instituto Hórus 2016; Ziller 2016). In addition, our list had almost twice as many species when compared to the study carried out by Leão et al. (2011) throughout north-eastern Brazil. It should also be noted that 52 species on our list were not mentioned in former studies.

In addition to our high sampling effort, the storage conditions of the biomes/ecosystems in Sergipe should explain the large number of taxa sampled here. Sergipe is considered the most degraded State in north-eastern Brazil (Ferrari 2005). Until the beginning of the 21st century, about 41% of the State had plant cover (Campos 1912), whereas, currently, there is only 13% natural forest cover (SEMARH-SE 2014).

The highest abundance of non-native invasive species with potential in the Atlantic Forest may be the result of strong environmental degradation which conditioned such biome (Myers et al. 2000; Tabarelli et al. 2005; INPE 2017). Degraded environments are more susceptible to the invasion process (Azevedo et al. 2010) and because of this, they tend to have a high dominance of exotic species (Huston 2004). According to Davis (2009), environmental degradation facilitates the invasion process, reducing competition and increasing the availability of resources.

Most of the species sampled by us are r-strategists, that is, they make pioneering use of the soil, have a short germination time, high growth rates and small size (Rejmánek and Richardson 1996). These attributes seem to be very important for the success of invasive alien species, since the same has also been observed in other studies (Horowitz et al. 2014; Fabricante et al. 2015; Galasso et al. 2018; Alves and Fabricante 2019; Silva and Fabricante 2019).

In Sergipe, the Atlantic Forest has been reduced to just 8% of its original area (Santos 2009), which is quite worrying since this biome is considered a priority area for biodiversity conservation – a hotspot (Myers et al. 2000). The condition of the Caatinga biome in the State is in a similar situation. It has been drastically reduced due to the exploration and use of its areas for crops and pastures (Arruda 2001).

The predominance of species belonging to the botanical family Poaceae and Fabaceae is due to the intrinsic characteristics of the taxa of these groups. Many of these species have an efficient dispersion system, form large seed-banks and have allelochemicals in their tissues (GISP 2005; Leão et al. 2011; Fabricante 2014). In addition, several of these species have been introduced massively to serve as fodder, especially in semi-arid regions (GISP 2005; Leão et al. 2011). Species, such as *Prosopis juliflora* (Sw.) DC. and *Leucaena leucocephala* (Lam.) de Wit, for example, exhibit drought resistance, high production of propagules and palatable parts, that is, characteristics that increase the invasive potential of species (Williams and Baruch 2000; Arriaga et al. 2004; CABI 2020).

Other floristic surveys of non-native species carried out in Caatinga sites also demonstrated a predominance of herbaceous species (Fabricante et al. 2015; Alves

and Fabricante 2019; Silva and Fabricante 2019). In the Atlantic Forest, woody species predominated (Souza et al. 2011; Mielke et al. 2015; Petri et al. 2018). Both situations corroborate our results. The edaphoclimatic conditions of biomes must favour certain extracts of vegetation (IBGE 2012; IFN 2017). Many herbaceous species were introduced into Brazil for human or animal food (Zenni 2014). Thus, it is expected that this habit will be prevalent in large-scale studies, such as this one.

The facilitation in the transfer of species that has occurred in recent years due to globalisation and advances in transport technologies (GISP 2005; Lockwood et al. 2007), associated with the market and people's interest in non-native species, may explain the existence of plants from different parts of the world found in our study. Most of these species were intentionally introduced into Brazil and other parts of the world, as they have different attributes, such as ornamentals, food, fuel production, fodder, amongst others (GISP 2005; Leão et al. 2011; Sampaio and Schmidt 2014; Zenni 2014).

Moreover, the high number of species of African origin can be justified by the climatic and biophysical similarities with Brazilian regions (GISP 2005; Herrera et al. 2016). According to Pérez et al. (2006) and Facon et al. (2006), the invasive potential of a species is intensified when the place of introduction has environmental conditions similar to its place of origin. In addition, a huge number of studies have already demonstrated through niche modelling that several invasive exotic species originating in Africa have great potential for invasion in Brazilian regions, including Sergipe (Araújo et al. 2013; Fabricante 2013; Alves et al. 2014; Herrera et al. 2016; Santos and Fabricante 2018).

The results presented here indicated a large number of non-native species with invasive potential and the existence of recognised aggressive species (e.g. *Prosopis juliflora* (Sw.) DC., *Leucaena leucocephala* (Lam.) de Wit, *Artocarpus heterophyllus* Lam., *Ricinus communis* L., *Calotropis procera* (Aiton) WT Aiton, *Terminalia catappa* L., *Boerhavia diffusa* L., *Cyperus rotundus* L., *Azadirachta indica* A. Juss., *Psidium guajava* L.). Such results may stimulate new studies, as well as serve as an aid to management, decision-making and the creation of public policies in the State.

References

- 13N [Invasives Information Network] (2020) Base de dados. <http://bd.institutohorus.org.br/www> [accessed on 27 October 2020]
- Alves CA, Stape JL, Sentelhas PC, de Moraes G, Leonardo J, Sparovek G (2013) Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift* 22(6): 711–728. <https://doi.org/10.1127/0941-2948/2013/0507>
- Alves JS, Fabricante JR (2019) Exotic invasive flora evaluation on different environments and preservation conditions from an area belonging to Caatinga, Petrolina, PE. *Gaia Scientia* 13(1): 40–52. <https://doi.org/10.22478/ufpb.1981-1268.2019v13n1.39803>
- Alves JS, Reis LBO, Silva EKC, Fabricante JR, Filho JAS (2014) *Leucaena leucocephala*. In: Fabricante JR (Ed.) *Plantas Exóticas e Exóticas Invasoras da Caatinga* (Vol. 4). Editora Bookess, Florianópolis, 13–20.

- Andrade LA, Fabricante JR, Oliveira FX (2009) Invasão biológica por *Prosopis juliflora* (Sw.) DC.: Impactos sobre a diversidade e a estrutura do componente arbustivo-arbóreo da caatinga no estado do Rio Grande do Norte, Brasil. *Acta Botanica Brasílica* 23(4): 935–946. <https://doi.org/10.1590/S0102-33062009000400004>
- Andrade LA, Fabricante JR, Oliveira FX (2010) Impactos da invasão de *Prosopis juliflora* (Sw.) DC. (Fabaceae) sobre o estrato arbustivo-arbóreo em áreas de Caatinga no Estado da Paraíba, Brasil. *Acta Scientiarum. Biological Sciences* 32(3): 249–255. <https://doi.org/10.4025/actascibiolsoci.v32i3.4535>
- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181(1): 1–20. <https://doi.org/10.1111/boj.12385>
- Araújo KCT, Fabricante JR, Castro RA, Furtado MDG, Filho JAS (2013) *Cenchrus echinatus*. In: Fabricante JR (Ed.) *Plantas Exóticas e Exóticas Invasoras da Caatinga* (Vol. 2). Editora Bookess, Florianópolis, 20–25.
- Arriaga L, Castellanos AE, Moreno E, Alarcón J (2004) Potential ecological distribution of alien invasive species and risk assessment: A case study of buffel Grass in arid regions of México. *Conservation Biology* 18(6): 1504–1514. <https://doi.org/10.1111/j.1523-1739.2004.00166.x>
- Arruda MB (2001) *Ecosistemas Brasileiros*. Edições IBAMA, Brasília, 49 pp.
- Azevedo C, Dechoum M, Zenni R, Ziller S, Zalba S (2010) *Espécies Exóticas Invasoras* (Vol. 3). *Cadernos da Mata Ciliar*, São Paulo, 35 PP.
- BioNET-EAFRINET (2020) Invasive plants. [http://keys.lucidcentral.org/keys/v3/eafrinet/weeds/key/weeds/Media/Html/Cynodon_dactylon_\(Couch_Grass\).htm](http://keys.lucidcentral.org/keys/v3/eafrinet/weeds/key/weeds/Media/Html/Cynodon_dactylon_(Couch_Grass).htm) [Accessed on 30 January 2020]
- Brasil (2009) Resolução n. 5, de 21 de Outubro de 2009. https://www.mma.gov.br/estruturas/conabio/_arquivos/resolucao_conabio05_estrategia_32a_nacional_especies_exticas_invasoras_15.pdf [Accessed on 27 October 2020]
- Brasil (2013) Portaria SEMA n° 79 de 31 de Outubro de 2013. http://www.institutohorus.org.br/download/marcos_legais/Portaria%20SEMA%20RS%2079%20-%202013%20Lista%20invasoras.pdf [Accessed on 30 January 2020]
- Brasil (2018) Resolução n. 7, de 29 de Maio de 2018. <https://www.mma.gov.br/images/arquivo/80049/Conabio/Resolucoes/RESOLUCAO%20N%207%20DE%2029%20DE%20MAIO%20DE%202018%20%20Diario%20Oficial%20da%20Uniao%20-%20Imprensa%20Nacional.pdf> [Accessed on 30 January 2020]
- CABI (2020) *Invasive Species Compendium*. <https://www.cabi.org/isc/> [Accessed on 26 April 2020]
- Campos LD (1912) *Mapa Florestal do Brasil*. Escala, São Paulo, 102 pp.
- CBD (1992) *Convention on Biological Diversity*. <https://www.cbd.int/convention/text/default.shtml> [Accessed on 17 October 2018]
- CBD (2020) *Aichi Biodiversity Targets*. <https://www.cbd.int/sp/targets/> [Accessed on 03 February 2020]
- Chen H, Boutros PC (2011) VennDiagram: A package for the generation of highly-customizable Venn and Euler diagrams in R. *BMC Bioinformatics* 12(1): 1–7. <https://doi.org/10.1186/1471-2105-12-35>

- Cronk QCB, Fuller JL (1995) *Plant Invaders: the threat to natural ecosystems*. Routledge, London, 241 pp.
- Davis MA (2009) *Invasion Biology*. Oxford University Press, New York, 241 pp.
- Fabricante JR (2013) *Plantas Exóticas e Exóticas Invasoras da Caatinga*. Bookess, Florianópolis, 51 pp.
- Fabricante JR (2014) *Plantas Exóticas e Exóticas Invasoras da Caatinga*. Bookess, Florianópolis, 50 pp.
- Fabricante JR, Siqueira-Filho JA (2012) Flora das Caatingas do Rio São Francisco. In: Siqueira-Filho JA (Org.) *Plantas Exóticas e Invasoras das Caatingas do Rio São Francisco (Vol. 1)*. Ed. Editora Andrea Jakobsson Estúdio, Petrolina, 367–394.
- Fabricante JR, Ziller SR, Araújo KCT, Furtado MDG, Basso FA (2015) Non-native and invasive alien plants on fluvial islands in the São Francisco River, northeastern Brazil. *Check List* 11(1): 1–7. <https://doi.org/10.15560/11.1.1535>
- Facon B, Genton BJ, Shykoff J, Jarne P, Estoup A, David P (2006) A general eco-evolutionary framework for understanding bioinvasions. *Trends in Ecology & Evolution* 21(3): 130–135. <https://doi.org/10.1016/j.tree.2005.10.012>
- Ferrari SF (2005) Parque Nacional Serra de Itabaiana: O Futuro. In: Carvalho CM, Vilar JC (Coords) *Parque Nacional Serra de Itabaiana-Levantamento da Biota (Vol. 10)*. Editora IBAMA, São Cristóvão, 121–131.
- Flora do Brasil (2020) Lista de Espécies da Flora do Brasil. <http://floradobrasil.jbrj.gov.br/reflora/listaBrasil/ConsultaPublicaUC/ConsultaPublicaUC.do#CondicaoTaxonCP> [Accessed on 27 October 2020]
- Galasso G, Conti F, Peruzzi L, Ardenghi NMG, Banfi E, Celesti-Grappo L, Albano A, Alessandrini A, Bacchetta G, Ballelli S, Bandini Mazzanti M, Barberis G, Bernardo L, Blasi C, Bouvet D, Bovio M, Cecchi L, Del Guacchio E, Domina G, Fascetti S, Gallo L, Gubellini L, Guiggi A, Iamónico D, Iberite M, Jiménez-Mejías P, Lattanzi E, Marchetti D, Martinetto E, Masin RR, Medagli P, Passalacqua NG, Peccenini S, Pennesi R, Pierini B, Podda L, Poldini L, Prosser F, Raimondo FM, Roma-Marzio F, Rosati L, Santangelo A, Scoppola A, Scortegagna S, Selvaggi A, Selvi F, Soldano A, Stinca A, Wagensommer RP, Wilhelm T, Bartolucci F (2018) An updated checklist of the vascular flora alien to Italy. *Plant Biosystems – An International Journal Dealing with all Aspects of Plant Biology* 152(3): 556–592. <https://doi.org/10.1080/11263504.2018.1441197>
- GISP (2005) *América do Sul invadida: a crescente ameaça das espécies exóticas invasoras*. Global Invasive Species Programme, Nairobi, 80 pp.
- Herrera I, Golçalves E, Pauchard A, Bustamante RO (2016) *Manual de Plantas Invasoras de Sudamérica*. Instituto de Ecología y Biodiversidad, Chile, 116 pp.
- Horowitz C, Martins CR, Walter BMT (2014) Flora exótica no Parque Nacional de Brasília: Levantamento e classificação das espécies. *Biodiversidade Brasileira* 3(2): 50–73.
- Huston MA (2004) Management strategies for plant invasions: Manipulating productivity, disturbance, and competition. *Diversity & Distributions* 10(3): 167–178. <https://doi.org/10.1111/j.1366-9516.2004.00083.x>
- IBGE (2012) *Manual técnico da vegetação brasileira*. Fundação Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro, 275 pp.

- IFN (2017) Inventário Florestal Nacional – Sergipe: principais resultados. Ministério do Meio Ambiente, Brasília, 87 pp.
- INPE (2017) Atlas dos remanescentes florestais da Mata Atlântica período 2015–2016. Instituto Nacional das Pesquisas Espaciais, São Paulo, 69 pp.
- Instituto Hórus (2016) Estratégias e políticas públicas para o controle das espécies exóticas invasoras. Secretaria do Ambiente e Desenvolvimento Sustentável, Porto Alegre, 52 pp.
- IPNI [International Plant Names Index] (2020) IPNI – International Plant Names Index. <https://www.ipni.org/> [Accessed 26 Oct 2020]
- ISSG (2020) Invasive Species Specialist Group. <http://www.issg.org/> [Access 07 June 2020]
- IUCN (1999) IUCN guidelines for the prevention of biodiversity loss due to biological invasion. *Species* 31(1): 28–42.
- Leão TCC, Almeida WR, Dechoum MS, Ziller SR (2011) Espécies Exótica Invasoras no Nordeste do Brasil: Contextualização, Manejo e Políticas Públicas. Centro de Pesquisas Ambientais do Nordeste e Instituto Hórus de Desenvolvimento e Conservação Ambiental, Recife, 99 pp.
- Lockwood JL, Hoopes MF, Marchetti MP (2007) *Invasion Ecology*. Blackwell Publishing, Oxford, 312 pp.
- Mielke EC, Negrelle RRB, Cuquel FL, Lima WP (2015) Espécies exóticas invasoras arbóreas no Parque da Barreirinha em Curitiba: Registro e implicações. *Ciência Florestal* 25(2): 327–336. <https://doi.org/10.5902/1980509818451>
- MMA (2020) Estratégia Nacional Para Espécies Exóticas Invasoras. <http://www.mma.gov.br/component/k2/item/7504> [Access on 02 February 2020]
- Mori SA, Silva LA, Lisboa G, Coradin L (1989) *Manual de Manejo do Herbário Fanerogâmico*. Editora Ilhéus, Ilhéus, 104 pp.
- Moro MF, Souza VC, Oliveira-Filho AT, Queiroz LPD, Fraga CND, Rodal MJN, Araújo FS, Martins FR (2012) Alienígenas na sala: O que fazer com espécies exóticas em trabalhos de taxonomia, florística e fitossociologia? *Acta Botanica Brasílica* 26(4): 991–999. <https://doi.org/10.1590/S0102-33062012000400029>
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GA, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403(6772): 853–858. <https://doi.org/10.1038/35002501>
- Nunes ADS, Higuchi P, Silva ACD, Kilca RDV, Silva MAFD, Larsen JG (2018) *Ligustrum lucidum* as an opportunist invasive species in an Araucaria Forest in South Brazil. *Rodriguésia* 69(2): 351–362. <https://doi.org/10.1590/2175-7860201869207>
- Oliveira-Costa J, Souza R (2015) Organização geossistêmica e invasões por *Acacia* sp. (Fabaceae: Mimosoideae) na Bacia Hidrográfica do Rio Arouce. Uma visão das invasões biológicas à escala do geossistema. *Revista de Geografia e Ordenamento do Território* 8(1): 145–169. <https://doi.org/10.17127/got/2015.8.008>
- ONUBR (2019) A ONU e o meio ambiente. <https://nacoesunidas.org/acao/meio-ambiente/> [Accessed on 02 February 2020]
- Pérez JE, Nirchio M, Alfonsi C, Muñoz C (2006) The biology of invasions: The genetic adaptation paradox. *Biological Invasions* 8(5): 1115–1121. <https://doi.org/10.1007/s10530-005-8281-0>

- Petri L, Aragaki S, Gomes EPC (2018) Management priorities for exotic plants in an urban Atlantic Forest reserve. *Acta Botanica Brasílica* 34(4): 631–641. <https://doi.org/10.1590/0102-33062017abb0317>
- Prata APN, Amaral MCE, Farias MCV, Alves MV (2013) *Flora de Sergipe* (Vol. 1). Gráfica e Editora Triunfo, Aracaju, 592 pp.
- Pujadas J (2001) Land-use and socio-economic correlates of plant invasions in European and North African countries. *Biological Conservation* 100(3): 397–401. [https://doi.org/10.1016/S0006-3207\(01\)00047-7](https://doi.org/10.1016/S0006-3207(01)00047-7)
- R Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. Version 4.0.0. <https://www.R-project.org/> [Accessed on 03 June 2020]
- Rejmánek M, Richardson DM (1996) What attributes make some plant species more invasive? *Ecology* 77(6): 1655–1661. <https://doi.org/10.2307/2265768>
- Richardson DM, Pyšek P, Rejmánek M, Barbour MG, Panetta FD, West CJ (2000) Naturalization and invasion of alien plants: Concepts and definitions. *Diversity & Distributions* 6(2): 93–107. <https://doi.org/10.1046/j.1472-4642.2000.00083.x>
- Sampaio AB, Schmidt IB (2014) Espécies exóticas invasoras em unidades de conservação federais do Brasil. *Biodiversidade Brasileira* 2(1): 32–49.
- Santos ALC (2009) Diagnóstico dos fragmentos de mata atlântica de Sergipe através de sensoriamento remoto. Dissertação de mestrado. São Cristóvão, Sergipe: Universidade Federal de Sergipe.
- Santos AI, Calafate L (2018) Espécies Invasoras. *Revista de Ciência Elementar* 6(1): 1–5. <https://doi.org/10.24927/rce2018.004>
- Santos JPB, Fabricante JR (2018) Biological invasion by *Thespesia populnea* in sites under fluvio-marine influence. *Neotropical Biology and Conservation* 13(1): 356–360. <https://doi.org/10.4013/nbc.2018.134.11>
- SEMARH-SE (2014) Diagnóstico Florestal de Sergipe. Secretaria de Estado de Meio Ambiente e dos Recursos Hídricos, Aracaju, 200 pp.
- Silva FO, Fabricante JR (2019) Invasão Biológica no Parque Nacional do Catimbau, Pernambuco, Brasil. *Revista de Ciências Ambientais* 13(1): 17–26. <https://doi.org/10.18316/rca.v13i2.5576>
- Souza AL, Ferreira RA, Mello AA, Plácido DDR, Santos CZAD, Graça DASD, Almeida Júnior PP, Barretto SSB, Dantas JDM, Paula JWA, Silva TL, Gomes LPS (2011) Diagnóstico quantitativo e qualitativo da arborização das praças de Aracaju, se. *Revista Árvore* 35(6): 1253–1263. <https://doi.org/10.1590/S0100-67622011000700012>
- Tabarelli M, Pinto LP, Silva JMC, Hirota MM, Bedê LC (2005) Desafios e oportunidades para a conservação da biodiversidade na Mata Atlântica brasileira. *Megadiversidade* 1(1): 132–138.
- Thiers B (2020) Index Herbariorum. Part I: The herbaria of the world. New York Botanical Garden. <http://sweetgum.nybg.org/science/ih/> [Accessed on 26 October 2020]
- Westbrooks R (1998) *Invasive Plants: Changing the Landscape of America*. Fact Book, Utah, 490 pp.

- Wickham H (2016) *ggplot2: elegant graphics for data analysis*. Springer. <https://doi.org/10.1007/978-3-319-24277-4>
- Williams DG, Baruch Z (2000) African grass invasion in the Americas: Ecosystem consequences and the role of ecophysiology. *Biological Invasions* 2(1): 123–140. <https://doi.org/10.1023/A:1010040524588>
- Williamson M (1996) *Biological Invasions*. Springer, London, 245 pp.
- Zenni RD (2014) Analysis of introduction history of invasive plants in Brazil reveals patterns of association between biogeographical origin and reason for introduction. *Austral Ecology* 39(4): 401–407. <https://doi.org/10.1111/aec.12097>
- Zenni RD, Ziller SR (2011) An overview of invasive plants in Brazil. *Revista Brasileira de Botânica. Brazilian Journal of Botany* 34(3): 431–446. <https://doi.org/10.1590/S0100-84042011000300016>
- Ziller SR (2001) Os processos de degradação ambiental originados por plantas exóticas invasoras. *Ciência Hoje* 30(1): 1–6.
- Ziller SR (2016) *Lista comentada de espécies exóticas invasoras do estado de Santa Catarina*. Fundação do Meio Ambiente, Santa Catarina, 85 pp.
- Ziller SR, Zalba S (2007) Propostas de ação para prevenção e controle de espécies exóticas invasoras. *Natureza & Conservação* 5(1): 8–15.