

# Phylogeography of the Central american red brocket deer, *Mazama temama* (Artiodactyla, Cervidae) in southeastern Mexico

Ricardo Serna-Lagunes<sup>1</sup>, Dayana Kristel Romero-Ramos<sup>1</sup>,  
Christian Alejandro Delfín-Alfonso<sup>2</sup>, Juan Salazar-Ortiz<sup>3</sup>

- 1 *Laboratorio de Bioinformática y Bioestadística. Facultad de Ciencias Biológicas y Agropecuarias, región Orizaba-Córdoba. Universidad Veracruzana. Calle Josefa Ortiz de Domínguez, sin number, Col. Peñuela, C. P. 94950, Amatlán de los Reyes, Veracruz, México*
- 2 *Instituto de Investigaciones Biológicas, Universidad Veracruzana, Av. Luis Castelazo Ayala, sin number, Col. Industrial Ánimas, C. P. 91190, Xalapa, Veracruz, México*
- 3 *Maestría en Paisaje y Turismo Rural. Colegio de Postgraduados, Campus Córdoba. Km. 348 Carretera Federal Córdoba-Veracruz, Congregación Manuel León, C. P. 94946, Amatlán de Los Reyes, Veracruz, México*

Corresponding author: Juan Salazar-Ortiz (salazar@colpos.mx)

---

Academic editor: A. M. Leal-Zanchet | Received 29 August 2020 | Accepted 27 May 2021 | Published 11 June 2021

---

Citation: Serna-Lagunes R, Romero-Ramos DK, Delfín-Alfonso CA, Salazar-Ortiz J (2021) Phylogeography of the Central american red brocketdeer, *Mazama temama* (Artiodactyla, Cervidae) in southeastern Mexico. *Neotropical Biology and Conservation* 16(2): 369–382. <https://doi.org/10.3897/neotropical.16.e58110>

---

## Abstract

Anthropogenic threats have increasingly isolated the populations of *Mazama temama* (Erxleben, 1777) and limited the gene flow in this species. Knowledge of the phylogeographic structure of this species is therefore essential for its conservation. Thus, in this study, we describe the phylogeographic structure of two *M. temama* populations of Veracruz and Oaxaca, Mexico. We sequenced the D-Loop region of the mitochondrial DNA of 16 individuals, in order to estimate the diversity and genetic differentiation ( $F_{ST}$ ), Tajima's D index, "Mismatch distribution" test; a phylogram and a haplotype network was constructed and we performed multidimensional scaling analysis to test the hypothesis of association between geographic distance and genetic diversity. The haplotypic and nucleotide diversity was high, indicating divergent populations ( $F_{ST} = 0.223$ ), while the Tajima's D index (-1,03300;  $P > 0.10$ ) determined disequilibrium in the D-Loop region, derived from a population expansion that was evidenced in the "Mismatch distribution" test and confirmed with the haplotype network in the form of a star. Four lineages were identified in the phylogram (Veracruz  $n = 3$ , Oaxaca  $n = 1$ ), evidencing geographic

and reproductive isolation between the two populations. This was confirmed by the multidimensional scaling analysis, which evidenced recent evolutionary divergence between the populations analyzed, which are considered evolutionary units of conservation.

### Keywords

Central American red brocket deer, lineages, nucleotide diversity, phylogram

## Introduction

The Central American Red Brocket, *Mazama temama* (Erxleben, 1777) is a deer species that has been targeted by illegal hunters for local commercial purposes or as a source of protein (Leopold 1959; Gallina and Mandujano 2009), as well as for its habitat (Gallina and Mandujano 2009; Bello-Gutiérrez et al. 2010). Even so, this deer species is considered Data Deficient by IUCN (Bello et al. 2016). This category is due to uncertainties about the threats that the species has suffered (Bello et al. 2016). Although threats such as those reported above are known, the degree of danger posed to the species is not clear (Bello et al. 2016). Thus, basic information about the species is essential to support conservation strategies.

Since certain populations of *M. temama* are found isolated in remnants of vegetation, it is important to study the phylogeography of this species in order to understand the micro-evolutionary processes these populations have undergone historically.

Phylogeography is a multidiscipline that has been used as a management tool for the conservation of species, and involves the study of micro and macroevolutionary processes that govern the geographic structure of genealogical lineages of species (Avice et al. 1987; Arbogast and Kenagy 2001; Vázquez-Domínguez 2002; Vázquez-Domínguez 2007) and features two important components: demographic structure and genetic structure (Avice 1992). An approach in the phylogeographic context can thus help to clarify changes in genealogical lineages based on their geographic distribution (Avice and Hamrick 1996; Avice 2000).

On the American continent, particularly in Mexico, phylogeography has been used to study populations of wild fauna, especially endemic mammals, or those sought by hunters, or those included in some category of risk, leading to important implications for their conservation (Vázquez-Domínguez 2005; Vega et al. 2007; Vázquez-Domínguez et al. 2009; Escobedo-Morales et al. 2016). There is evidence that phylogeographic studies have had a considerable impact on species management (Vázquez-Domínguez and Vega 2002), especially in the ungulates, describing cervid lineages in order to define particular strategies of management (Serna-Lagunes et al. 2021), identifying bottlenecks and low gene flow in species of interest to hunters (Castillo-Rodríguez et al. 2020) and evidence of genetic divergence among lineages of the same species (Fuentes-Hurtado et al. 2011). In this sense, in this study we describe the phylogeographic structure of two populations of *M. temama*.

## Material and methods

### Collection of biological material

We were able to generate molecular data from 10 specimens of *M. temama*, among two neighboring populations in the states of Veracruz and Oaxaca, in Mexico (Table 1, Fig. 1). Eight live specimens were confiscated by the environmental authority (Procuraduría Federal de Protección al Ambiente-PROFEPA), which authorized the collection of blood and skin samples. The specimens were seized in five localities of the Sierra de Zongolica, Zongolica, Tehuipango, Tequila and Tezonapa in Las Montañas region, in the center of Veracruz, where the main threats to populations of this species are the fragmentation of their habitat, the isolation of their populations in secondary vegetation, as well as the hunting and illegal possession of these deer as pets (Salazar-Ortiz et al. 2020). The specimens are currently being kept in the wildlife management unit (UMA, by its Spanish acronym) “El Pochote”, in Ixtaczoquitlán, Veracruz. The two other skin samples come from two localities in the state of Oaxaca and were provided by the Colección Nacional de Mastozoología of the Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional (CIIDIR), Oaxaca unit, of the Instituto Politécnico Nacional, with registry of origin in Oaxaca (Table 1).


### Extraction of DNA, PCR and sequencing

The blood samples were dehydrated at ambient temperature and the skin samples fragmented in order to extract the DNA in accordance with instructions from the DNeasy Blood and Tissue kit of QIAGEN. DNA extraction was verified by applying electrophoresis in a horizontal 1% agarose gel with ethidium bromide at 70V for 40 min in 1X buffer TAE. The bands of the extracted DNA were visualized in a transilluminator and documented with the program GelCapture.

From each extracted DNA sample, 2.5  $\mu$ L were taken to perform the Polymerase Chain Reaction (PCR) at the final phase or end-point. From this DNA, the Con-

**Table 1.** Geographic data of the collection sites, localities in Veracruz and Oaxaca, in Mexico, including the registry code for the nucleotide sequence database GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>). Note: \* sample collected in the UMA “El Pochote”, Ixtaczoquitlán, Veracruz.

State	Sequence GenBank	Sample type	Coordinates	Locality
Veracruz	MG910312	Skin	18°44'08.99"N, 96°55'14.00"W	La Pila, Zongolica*
	MG910321		18°40'00.65"N, 97°00'00.44"W	Zongolica*
	MG910314		18°42'47.00"N, 96°56'43.00"W	Totolacatla, Zongolica*
	MG910315		18°42'54.00"N, 96°58'40.01"W	Laguna Ixpaluca, Zongolica*
	MG910316		18°32'29.00"N, 96°46'44.00"W	Rancho Nuevo, Tezonapa*
	MG910317	Blood	18°40'00.65"N, 97°00'00.44"W	Zongolica*
	MG910318		18°40'00.65"N, 97°00'00.44"W	Zongolica*
	MG910319		18°40'00.65"N, 97°00'00.44"W	Zongolica*
	MG910320		17°28'06.06"N, 96°02'09.44"W	San Juan Petlapa
Oaxaca	MG910313	Skin (cat. OAX.MA 4988)	17°15'24.58"N, 96°01'44.22"W	Totontepec, Villa de Morelos



**Figure 1.** Geographic location of the *M. temama* samples collection sites in Oaxaca and Veracruz, Mexico.

trol or D-Loop region of the DNA<sub>mt</sub> was analyzed using the universal primers: DL-HI16498 (5' CCTGAAGTACTAGGAACCCAGATG 3') and Thr-L15910 (5' GAATTCC-CCGGTCTTGTA 3') (Vilà et al. 1999). The reactions were performed in a final volume of 25 µl and contained the following: 2.5 µl of DNA (< 250 ng), 1.25 µl of the Forward and 1.25 µl of the Reverse primers, both at a final concentration of

10  $\mu\text{M}$ , 12.5  $\mu\text{l}$  of 2X Promega PCR Master Mix (25mM Tris-HCl pH 9, 25mM NaCl, 2.5mM  $\text{MgCl}_2$ , 100  $\mu\text{M}$  each of dATP, dGTP, dCTP, dTTP, 0.5 U of Taq DNA polymerase, 0.05 mg/ml BSA) and 7.5  $\mu\text{l}$  of nuclease-free water. The PCR was conducted in a thermocycler, using the following program: initial denaturalization at 95 °C for 1 min, followed by 35 cycles of amplification: 94 °C for 1 min, annealing at 51 °C for 2 min, extension at 72 °C for 1 min 30 sec and a final extension step at 72 °C for 7 min. This procedure was conducted in the Laboratorio Cordobés de Diagnóstico Pecuario, S.C., located in Córdoba, Veracruz, Mexico. The PCR Products were verified using Green DNA Dye. Verification of the PCR products was corroborated using the same DNA extraction method.

The PCR products were sent to the company MacroGen (Korea) for purification and sequencing. MacroGen employs the Sanger technique, with capillary technology and an Applied Biosystems 3730XLs sequencer in order to perform simple sequencing of nucleotides ([www.macrogen.com](http://www.macrogen.com)).

### Statistical analysis

A database was generated with the obtained sequences, grouping them according to population. The sequences were aligned with the algorithm Clustal W, installed in the program MEGA v. 6 (Tamura et al. 2013). For each population, the nucleotide composition (proportion of each nitrogenous base) of the sequences was estimated. The number of polymorphic sites ( $S$ ), number of identified haplotypes ( $H_i$ ), number of haplotypes shared between populations ( $H_s$ ), haplotypic diversity ( $H_d$ ) and nucleotide diversity ( $\pi$ ) were calculated. The index of genetic differentiation ( $F_{ST}$ ) was also calculated as a measure with which to distinguish between the populations under study. The Tajima's D neutrality test (Tajima 1989) was applied as an indicator of historic changes in population expansion. The "Mismatch distribution" test was performed to measure the effect on population size of the historic demographic changes that molded the genetic diversity observed. This analysis is represented in a graph that compares the observed and expected nucleotide frequencies (Rogers et al. 1996). The fit of this test was verified with the values of  $R^2$ , initial theta ( $\theta$ ) and tau ( $\tau$ ) (Ramos-Onsins and Rozas 2002). These analyses were performed with the software DnaSP v.5 (Librado and Rozas 2009).

The model of nucleotide substitution that best fitted the sequences of *M. temama* was identified (Nei and Kumar 2000) using the software MEGA v.6 (Tamura et al. 2013). The model of Kimura 2 (K2) was that which best described the pattern of nucleotide substitution by presenting a lower number of parameters (18) under the coefficient of the Bayesian Information Criterion (BIC = 3394.84), with a mean value of the Akaike Information Criterion (AICc = 3301.260) and the highest value of the parameter of Maximum Likelihood (lnL = -1632.379) (Nei and Kumar 2000; Tamura et al. 2013). The most parsimonious phylogram was constructed with Maximum Parsimony (MP), based on the Branch and Bound algorithm (Nei and Kumar 2000). The fit of the phylogram branches was evaluated with 1000 Bootstrap

replicates (Felsenstein 1985) using the software MEGA v.6 (Tamura et al. 2013), following the protocols of Hall (2004 and 2013). A haplotype network was constructed with Median-Joining Network and MP. This determines the mutational steps between haplotypes and was generated with the software Network v.5 (Fluxus Technology Ltd 1999–2016).

Finally, nucleotide distances were estimated between pairs of sequences with the software DNAsp (Librado and Rozas 2009), in order to determine the degree of current genetic similarity or divergence, where short genetic distances indicate close relationships while long distances suggest distant relationships. Likewise, geographic distances (km: between collection localities) were obtained between pairs of sequences using ArcMap v.10.3. With these distances (nucleotide and geographic, see Table 4), a multidimensional scaling analysis of two dimensions (MDS; Borg and Groenen 2003) was conducted to determine the similarity (or dissimilarity) among the sequences. This analysis is a test of interdependence that calculates a matrix of paired Euclidean distances, producing a graph with a geometric space of two dimensions (Lessa 1990), on which are represented the proximities that exist between the nucleotide and geographic distances of each sequence; the fit of the graph was evaluated with the values of “Stress” (>0.2 good or >0.7 very good) and  $R^2$  (>0.3 good fit or >0.7 very good fit). In addition, a principal component analysis (PCA) was applied to the matrix of nucleotide and geographic distances with the Varimax rotation method (Yakubu et al. 2011). Both analyses were conducted with the software SPSS 15.0 (SPSS 2004).

## Results


The analysis of the nucleotide composition of the obtained sequences indicated a higher average frequency of Cytosine bases in the *M. temama* population of Veracruz, while the Oaxaca population presented a greater proportion of Adenine bases (Table 2).

**Table 2.** Nucleotide composition of the *M. temama* sequences from Veracruz and Oaxaca, Mexico. MV = *Mazama* of Veracruz; MO = *Mazama* of Oaxaca.

State (Population)	Sequence	ID GenBank sequence	T (%)	C (%)	A (%)	G (%)	Total	Locality
Veracruz	MG910312	1_MV	18.9	22.2	32.2	26.7	180	La Pila, Zongolica
	MG910321	8_MV	33.7	30.6	15.8	19.9	196	Zongolica
	MG910313	20_MV	20.3	24.6	32.9	22.2	207	Totolacatla, Zongolica
	MG910314	21_MV	33.3	27.8	19.7	19.2	198	Laguna Ixpaluca, Zongolica
	MG910315	39_MV	24.5	25.5	24.5	25.5	200	Rancho Nuevo, Tezonapa
	MG910316	41_MV	24.3	28.2	28.6	18.9	206	Zongolica
	MG910317	42_MV	31.1	21.4	27.2	20.4	206	Zongolica
	MG910318	45_MV	21.7	32.2	24.4	21.7	180	Zongolica
Mean (%)=			26	26.6	25.7	21.8	196.6	
Oaxaca	MG910320	5_MO	21.4	20.8	35.8	22	159	Totontepec, Villa de Morelos
	MG910312	13_MO	28.8	21.5	26.2	23.6	191	San Juan Petlapa
	Mean (%)=		25.1	21.1	31	22.8	175	

**Table 3.** Genetic diversity of two *M. temama* populations of Veracruz and Oaxaca, in Mexico. n, number of sequences;  $H_d$ , haplotypic diversity;  $H_i$ , number of haplotypes identified;  $\pi$ , nucleotide diversity; S, number of polymorphic sites.

Population	n	S	$H_i$	$H_d$	$\pi$
Oaxaca	2	48	2	1.00 ± 0.5	0.307 ± 0.21
Veracruz	8	149	8	1.00 ± 0.063	0.653 ± 0.167




**Figure 2.** "Mismatch distribution" for the *M. temama* populations from Oaxaca (left) and Veracruz (right), in Mexico.


**Table 4.** Matrix of nucleotide (below the diagonal) and geographic (above the diagonal, in km) distance between *M. temama* haplotypes from Veracruz and Oaxaca, in Mexico. MV = *Mazama* of Veracruz; MO = *Mazama* of Oaxaca.

ID sequence register in GenBank	1_MV	5_MO	8_MV	13_MO	20_MV	21_MV	39_MV	41_MV	42_MV	45_MV
1_MV	-	190	12	169	4	7	26	12	12	12
5_MO	1.183	-	189	24	191	190	164	188	188	188
8_MV	0.883	1.118	-	167	8	7	28	2	2	2
13_MO	1.375	0.365	0.948	-	168	171	142	167	167	167
20_MV	1.818	1.193	1.589	1.447	-	4	26	8	8	8
21_MV	2.116	1.42	2.198	1.431	2.156	-	28	6	6	6
39_MV	1.818	1.143	1.268	1.38	0.097	1.948	-	27	27	27
41_MV	0	1.587	0	1.565	1.529	2.482	1.529	-	0	0
42_MV	0	2.103	1.682	1.458	2.262	2.622	1.791	2.103	-	0
45_MV	0	1.713	1.589	1.762	2.393	0	2.023	2.262	0	-

The two populations of *M. temama* showed differences in genetic diversity. In the population of Oaxaca, there was a lower number of polymorphic sites compared to the population of Veracruz, which explains why the number of haplotypes was different. That is, each sample analyzed was found to be an independent haplotype, and haplotypes were not shared between the populations. Haplotypic and nucleotide diversity was high in both populations, indicating that the sampled individuals of each population were heterogeneous in terms of the D-loop region (Table 3). This is consistent with the results of the index of genetic differentiation ( $F_{ST}$ ) = 0.223, which indicated genetic differences between the populations of Oaxaca and Veracruz, as a result of not sharing haplotypes ( $H_i$  = 0).



**Figure 3.** Phylogram of genealogical relationships between the two *M. temama* populations of Oaxaca and Veracruz, in Mexico. Values on the branches represent the fit of the Bootstrap test.



**Figure 4.** Haplotype network of the two *M. temama* populations of Oaxaca and Veracruz, in Mexico.

The Tajima's D index was negative (-1,03300;  $P > 0.10$ ), indicating the presence of deletions or mutations, such that the populations were not found in neutral equilibrium. The "Mismatch distribution" test fitted to the model of a population of constant size (Oaxaca:  $R^2 = 0.500$ , initial theta ( $\theta$ )= 14.750, tau ( $\tau$ )= 48; Veracruz:  $R^2 = 0.217$ , initial theta ( $\theta$ )= 14.75, tau ( $\tau$ )= 82.571), evidencing a possible population expansion by the variation shown in the graphs of this test (Fig. 2).

The phylogram [(consistency index (CI)= 0.631579 and, retention index (RI)= 0.427673)], showed paraphyletic relationships between the samples, indicating that there is no geographic structure. For example, some Veracruz samples were closer to Oaxaca samples than other Veracruz populations (Fig. 3).


The haplotype network presented a structure in the form of a star (Fig. 4), with 351 total non-segregating sites and 233 mutations. This is commonly observed in populations that have undergone a process of demographic expansion and was consistent with the results of the Tajima's D index and the "Mismatch distribution" test.

According to the MDS (Fig. 5), the *M. temama* individuals most related according to nucleotide distance were those found in southern Veracruz and Oaxaca (upper right extreme of the dimensional plane), while the least related were those of central and southern Veracruz (extreme lower left of the dimensional plane), which explains the genetic relationship based on geographic distance; the stress value was good (0.14), with a value of  $R^2 = 0.87$ , indicating a significant relationship between the pairs of sequences of the matrix of nucleotide and geographic distances (Fig. 5). Finally, the PCA determined that the greatest proportion of the variance was accommodated in the first two principal components (>80%). Principal component 1 explained 63.8 of the variances (Table 5), while principal component 2 explained 16.3% and together these components explained 80.3% of the total accumulated variation (Fig. 6).

## Discussion

In this study we performed our first phylogeographic analysis on the species *M. temama*. Our results show that the populations analyzed underwent a genetic bottleneck and rapid population expansion.


Although we have studied a small sample of the population, these populations studied here do not share haplotypes due to the geographical distance between the two populations. On the other hand, it is possible that the reproductive behavior



**Figure 5.** Multidimensional scaling analysis applied to the matrix of nucleotide and geographic distances of pairs of *M. temama* sequences.

**Table 5.** Total explained variance of the Principal Components Analysis (APC) applied to the matrix of nucleotide and geographic distances of pairs of *M. temama* sequences, showing the values in at least the first two components.

Component	Extraction sums of squared loadings			Rotation sums of squared loadings		
	Total	Variance proportion	% accumulated	Total	Variance proportion	% accumulated
I	6.380	63.803	63.803	6.347	63.467	63.467
II	1.658	16.578	80.381	1.691	16.913	80.381



**Figure 6.** Principal components analysis applied to the matrix of nucleotide and geographic distances of pairs of *M. temama* sequences.

of this cervid means that the frequency of mtDNA transfer decreases, due to the philopatry of the females (Purdue et al. 2000; Purdue et al. 2006). It is only the males that migrate great distances to mate and thus the lineages of the females are presented in greater proportion with proximity to their area of birth (Nelson and Mech 1992). The lack of shared haplotypes between the two studied *M. temama* populations is evidence of the geographic and reproductive isolation.

The results of the "Mismatch distribution" test show an evident multimodal form, reflecting processes of sudden population expansion associated with a stochastic process of lineage extinction through genetic drift, recorded in the genetic structure of the populations (Rogers and Harpending 1992). This may reflect the threats that the species suffers in Mexico; *M. temama* is of hunting interest and subject to constant pressures of exploitation, habitat fragmentation, poaching, auto-consumption by local communities and predation by carnivores (Gallina 2005). This has led to a reduction in the effective population size where only certain lineages pass to the next generation, while others are extinguished or are present in lower frequency in the population. These results are to be expected considering that *M. temama* is a species that suffers from poaching, since a similar phenomenon has been reported in the cervid *Pantholops hodgsonii* in China, which is at risk of extinction due to the loss of genetic diversity (Xiang-Dong et al. 2005).

The populations analyzed have different evolutionary histories, as shown by the string haplotypic and nucleotide relationship between the populations, evidencing differentiated lineages between the two populations (Grant and Bowen 1998). It is concerning that the populations studied present low values of gene flow, as occurs in some other species of Neotropical cervids. This is associated with the fact that *M. temama* inhabits fragmented ecosystems with no biological corridors to allow migration of individuals for the purposes of mating. Phylogeographic studies must

be conducted in more populations of *M. temama* in order to evaluate their status of genetic diversity and to generate a program of conservation of their diversity, since their ecological niche is restricted to vulnerable ecosystems due to the reduction and fragmentation of their habitat (Serna-Lagunes et al. 2014).

## Acknowledgements

The authors offer their sincere gratitude to the reviewers of our manuscript, whose observations and comments enabled the improvement of this research. Thanks go to Linda Elvira Rodríguez Vera, Jorge Noriega Loyo and the Área de Biología Molecular of the Laboratorio Cordobés de Diagnóstico Pecuario for validation of the PCR and provision of materials and equipment. Thanks also to the UMA El Pochote and the Colección Nacional de Mastozoología (CIIDIR), for the tissue samples. To the project “Characterization of animal genetic resources of the high mountains, Veracruz: application of phylogeography and ecological modeling (PRODEP: 511-6/18-9245/PTC-896) for their support in finalizing this publication.

## References

- Arbogast BS, Kenagy GJ (2001) Comparative phylogeography as an integrative approach to historical biogeography. *Journal of Biogeography* 28(7): 819–825. <https://doi.org/10.1046/j.1365-2699.2001.00594.x>
- Avice JC (1992) Molecular population structure and the biogeographic history of a regional fauna: a case history with lessons for conservation biology. *Oikos* 63: 62–76. <https://doi.org/10.2307/3545516>
- Avice JC (2000) *Phylogeography: the history and formation of species*. Harvard University Press, Cambridge, Massachusetts, 447 pp. <https://doi.org/10.2307/j.ctv1nznfjg7>
- Avice JC, Hamrick JL (1996) *Conservation genetics: case histories from nature*. Chapman & Hall, Springer, EEUU, 512 pp. <https://doi.org/10.1007/978-1-4757-2504-9>
- Avice JC, Arnold J, Ball RM, Bermingham E, Lamb T, Neigel JE, Reeb CA, Saunders NC (1987) Intraspecific phylogeography: The mitochondrial DNA bridge between population genetics and systematics. *Annual Review of Ecology and Systematics* 18(1): 489–522. <https://doi.org/10.1146/annurev.es.18.110187.002421>
- Bello J, Reyna R, Schipper J (2016) *Mazama temama*. The IUCN Red List of Threatened Species 2016: e.T136290A22164644. <https://doi.org/10.2305/IUCN.UK.2016-2.RLTS.T136290A22164644.en>
- Bello-Gutiérrez J, Reyna-Hurtado R, Wilham J (2010) Central American red brocket deer *Mazama temama* (Kerr, 1992). In: Duarte JMB, González S (Eds) *Neotropical Cervidology: biology and medicine of Latin American deer*. Funep, Jaboticabal, Brasil/IUCN, Gland, 166–171.
- Borg I, Groenen P (2003) Modern multidimensional scaling: Theory and applications. *Journal of Educational Measurement* 40(3): 277–280. <https://doi.org/10.1111/j.1745-3984.2003.tb01108.x>

- Castillo-Rodríguez RG, Serna-Lagunes R, Cruz-Romero A, Núñez-Pastrana R, Rojas-Avelizapa LI, Llarena-Hernández RC, Dávila JA (2020) Characterization of the genetic diversity of a population of *Odocoileus virginianus veraecrucis* in captivity using microsatellite markers. *Neotropical Biology and Conservation* 15(1): 29–41. <https://doi.org/10.3897/neotropical.15.e47262>
- Escobedo-Morales LA, Mandujano S, Eguiarte LE, Rodríguez-Rodríguez MA, Maldonado JE (2016) First phylogenetic analysis of Mesoamerican brocket deer *Mazama pandora* and *Mazama temama* (Cetartiodactyla: Cervidae) based on mitochondrial sequences: Implications for Neotropical deer evolution. *Mammalian Biology* 81(3): 303–313. <https://doi.org/10.1016/j.mambio.2016.02.003>
- Felsenstein J (1985) Confidence limits on phylogenies: An approach using the Bootstrap. *Evolution; International Journal of Organic Evolution* 39(4): 783–791. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>
- Fluxus Technology Ltd (1999–2016) Network 4.5.1.0. User Guide. <http://www.fluxus-engineering.com/sharenet.htm>
- Fuentes-Hurtado M, Marin JC, Gonzalez-Acuna D, Verdugo C, Vidal F, Vianna JA (2011) Molecular divergence between insular and continental Pudu deer (*Pudu puda*) populations in the Chilean Patagonia. *Studies on Neotropical Fauna and Environment* 46(1): 23–33. <https://doi.org/10.1080/01650521.2010.537906>
- Gallina S (2005) Temazate. In: Ceballos G, Oliva G (Eds) *Los mamíferos silvestres de México*. CONABIO-Fondo de cultura Económica, México, 512–513.
- Gallina S, Mandujano S (2009) Research on ecology, conservation and management of wild ungulates in Mexico. *Tropical Conservation Science* 2(2): 116–127. <https://doi.org/10.1177/194008290900200201>
- Grant WS, Bowen BW (1998) Shallow population histories in deep evolutionary lineages of marine fishes: Insights from sardines and anchovies and lesson for conservation. *The Journal of Heredity* 89(5): 415–426. <https://doi.org/10.1093/jhered/89.5.415>
- Hall BG (2004) *Phylogenetic trees made easy: A how-to manual*. Sinauer Associate, Sunderland, EU.
- Hall BG (2013) Building phylogenetic trees from molecular data with MEGA. *Molecular Biology and Evolution* 30(5): 1229–1235. <https://doi.org/10.1093/molbev/mst012>
- Leopold AS (1959) *Wildlife of Mexico, the game birds and mammals*. University California Press, Berkeley, EUA. <https://doi.org/10.1525/9780520326422>
- Librado P, Rozas J (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics (Oxford, England)* 25(11): 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Nei M, Kumar S (2000) *Molecular evolution and phylogenetic*. Oxford University Press, New York.
- Nelson ME, Mech LD (1992) Dispersal in female White-Tailed deer. *Journal of Mammalogy* 73(4): 891–994. <https://doi.org/10.2307/1382212>
- Purdue JR, Smith MH, Patton JC (2000) Female philopatry and extreme spatial genetic heterogeneity in White-Tailed Deer. *Journal of Mammalogy* 81(1): 179–185. [https://doi.org/10.1644/1545-1542\(2000\)081%3C0179:FPAESG%3E2.0.CO;2](https://doi.org/10.1644/1545-1542(2000)081%3C0179:FPAESG%3E2.0.CO;2)

- Purdue JR, Oleksyk TK, Smith MH (2006) Independent occurrences of multiple repeats in the control region of mitochondrial DNA of White-Tailed Deer. *The Journal of Heredity* 97(3): 235–243. <https://doi.org/10.1093/jhered/esj032>
- Ramos-Onsins SE, Rozas J (2002) Statistical properties of new neutrality tests against population growth. *Molecular Biology and Evolution* 19(12): 2092–2100. <https://doi.org/10.1093/oxfordjournals.molbev.a004034>
- Rogers A, Harpending H (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution* 9: 552–569. <https://doi.org/10.1093/oxfordjournals.molbev.a040727>
- Rogers AR, Fraley AE, Bamshad MJ, Watkins WS, Jorde LB (1996) Mitochondrial Mismatch analysis is insensitive to the mutational process. *Molecular Biology and Evolution* 13(7): 895–902. <https://doi.org/10.1093/molbev/13.7.895>
- Salazar-Ortiz J, Barrera-Perales MG, Ramírez-Ramírez Y, Serna-Lagunes R (2020) Diversidad de mamíferos del municipio de Tequila, Veracruz, México. *Revista Abanico Veterinario* 10: 1–18. <https://doi.org/10.21929/abavet2020.30>
- Serna-Lagunes R, López-Mata L, Cervantes-Serna LJ, Gallegos-Sánchez J, Cortez-Romero C, Zalazar-Marcial E, Sánchez-Páez R, Olguín CA, Salazar-Ortiz J (2014) Nicho ecológico de *Mazama temama* en el centro de Veracruz, México: Implicaciones para su manejo. *Agroproductividad* 7: 51–67.
- Serna-Lagunes R, Pastor-Medina EJ, Castillo-Rodríguez RG, Cruz-Romero A, Salazar-Ortiz J, Del Rosario-Arellano JL, García-Martínez MA, Mora-Collado N, Andrés-Meza P (2021) Genetic Diversity of *Odocoileus virginianus veraecrucis* (Goldman & Kellog, 1940) and Other's Subspecies in Mexico: Implications for Its Genetic Conservation. In: Nardelli M, Túnez JI (Eds) *Molecular Ecology and Conservation Genetics of Neotropical Mammals*. Springer, Cham. Springer Nature Switzerland AG, 149–170. [https://doi.org/10.1007/978-3-030-65606-5\\_7](https://doi.org/10.1007/978-3-030-65606-5_7)
- Tajima F (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123(3): 585–595. <https://doi.org/10.1093/genetics/123.3.585>
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S (2013) MEGA 6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30(12): 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Vázquez-Domínguez E (2002) Phylogeography, historical patterns and conservation of natural areas. In: Nelson G, Day JC, Sportza LM, Loucky J, Vásquez C (Eds) *Protected areas and the regional planning imperative in North America*. University of Calgary Press, Canada, 369–378. <https://doi.org/10.2307/j.ctv6cfq65.29>
- Vázquez-Domínguez E (2005) ¿Cuál es el enfoque de la ecología molecular? Ejemplos de un ratón espinoso. In: Sánchez-Rojas G, Rojas-Martínez AE (Eds) *Tópicos en sistemática, biogeografía, ecología y conservación de mamíferos*. Centro de Investigaciones Biológicas, Universidad Autónoma del Estado de Hidalgo, México, 59–70.
- Vázquez-Domínguez E (2007) Filogeografía y vertebrados. In: Eguiarte LE, Souza V, Aguirre X (Eds) *Ecología Molecular*. Secretaría de Medio Ambiente y Recursos Naturales. Instituto Nacional de Ecología. Universidad Nacional Autónoma de México, Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México, 441–466.

- Vázquez-Domínguez E, Vega R (2002) ¿Dónde estamos y hacia dónde vamos en el conocimiento genético de los mamíferos mexicanos? Departamento de Ecología de la Biodiversidad, Instituto de Ecología, Universidad Nacional Autónoma de México.
- Vázquez-Domínguez E, Castañeda-Rico S, Garrido-Garduño T, Gutiérrez-García TA (2009) Avances metodológicos para el estudio conjunto de la información genética, genealógica y geográfica en análisis evolutivos y de distribución. *Revista Chilena de Historia Natural* 82(2): 277–297. <https://doi.org/10.4067/S0716-078X2009000200009>
- Vega R, Vázquez-Domínguez E, Mejía A, Cuarón AD (2007) Unexpected high levels of genetic variability and the population structure of an island endemic rodent (*Oryzomys couesi cozumelae*). *Biological Conservation* 137(2): 210–222. <https://doi.org/10.1016/j.biocon.2007.02.007>
- Vilà C, Amorim IR, Leonard JA, Posada D, Castroviejo J, Petrucci-Fonseca F, Crandall KA, Ellegren H, Wayne RK (1999) Mitochondrial DNA phylogeography and population history of the grey wolf *Canis lupus*. *Molecular Ecology* 8(12): 2089–2103. <https://doi.org/10.1046/j.1365-294x.1999.00825.x>
- Yakubu A, Salako AE, Abdullah AR (2011) Varimax rotated principal component factor analysis of the zoometrical traits of Uda sheep. *Archivos de Zootecnia* 60(231): 813–816. <https://doi.org/10.4321/S0004-05922011000300069>