



Foam nest construction and first report of agonistic behaviour in *Pleurodema tucumanum* (Anura, Leptodactylidae)

Melina J. Rodríguez Muñoz^{1,2}, Tomás A. Martínez^{1,2}, Juan Carlos Acosta¹,
Graciela M. Blanco¹

- 1 Gabinete DIBIOVA (Diversidad y Biología de Vertebrados del Árido). Departamento de Biología, FCEF, Universidad Nacional de San Juan, Avenida Ignacio de la Roza 590, Rivadavia J5400DCS, San Juan, Argentina
- 2 Consejo Nacional de Investigaciones Científicas y Técnicas, Godoy Cruz 2290, C1425FQB, Ciudad Autónoma de Buenos Aires Argentina

Corresponding author: Melina J. Rodríguez Muñoz (melina.rodriguez26@gmail.com)

Academic editor: A. M. Leal-Zanchet | Received 21 May 2018 | Accepted 27 December 2018 | Published 11 April 2019

Citation: Rodríguez Muñoz MJ, Martínez TA, Acosta JC, Blanco GM (2019) Foam nest construction and first report of agonistic behaviour in *Pleurodema tucumanum* (Anura: Leptodactylidae). Neotropical Biology and Conservation, 14(1): 117–128. <https://doi.org/10.3897/neotropical.14.e34841>

Abstract

Reproductive strategies are the combination of physiological, morphological, and behavioural traits interacting to increase species reproductive success within a set of environmental conditions. While the reproductive strategies of Leiuperinae are known, few studies have been conducted regarding the reproductive behaviour that underlies them. The aim of this study was to document the structural characteristics of nesting microsites, to describe the process of foam nest construction, and to explore the presence of male agonistic and chorus behaviour in *Pleurodema tucumanum*. Nests were found close to the edge of a temporary pond and the mean temperature of the foam nests was always close to the mean temperature of the pond water. Our observations corroborate that the foam nest construction phases for *P. tucumanum* are similar to those described for other Leiuperinae, including dispersion and the beating of the foam, realized by male hind limbs. We also recorded the first scientific observations of male agonistic encounter in *Pleurodema tucumanum*.

Keywords

Amphibians, Dry Chaco, male-male interaction, oviposition site, reproductive mode

Introduction

Reproductive modes can substantially vary among anuran species (Haddad and Prado, 2005) and are therefore considered of taxonomic value (Duellman and Trueb, 1986). Foam nest reproductive modes in anurans are a convergent behaviour that evolved independently in various frog lineages (Duellman and Trueb, 1986; Seymour and Loveridge, 1994; Faivovich *et al.* 2012). The ancestral oviposition mode for the Neotropical anuran clade Leiuperinae, composed by the genera *Edalorhina*, *Engystomops*, *Physalaemus*, and *Pleurodema*, is the floating foam nest (Grant *et al.* 2006; Pyron and Wiens, 2011; Faivovich *et al.* 2012).

The genus *Pleurodema* is notable for its particularly large variety of oviposition modes (Duellman and Veloso, 1977; Faivovich *et al.* 2012; Valetti *et al.* 2014). Faivovich *et al.* (2012) reported four egg-clutch structures proposed by different authors. However, recently Ferraro *et al.* (2016) rejects the egg-clutch structure proposed by Martori *et al.* (1994). Thus, three reproductive modes are known for *Pleurodema*: (1) eggs laid in subspherical gelatinous masses in groups of 1–6 eggs (Barrio, 1977; Valetti *et al.* 2009), (2) eggs laid in gelatinous strings (Fernández, 1927; Weigandt *et al.* 2004), and (3) eggs laid in a floating foam nest (Hödl, 1992; Cardoso and Arzabe 1993).

In arid environments, under temporarily harsh, unpredictable and variable conditions, species should possess special mechanisms for tolerating or avoiding stressful situations. During the aquatic phase, larvae and nests are often at risk of drying in ephemeral ponds (Spieler and Linsenmair, 1997). The foam nest structure protects eggs and tadpoles from dehydration, and is a beneficial reproductive strategy for species that inhabit arid environments (Prado *et al.* 2002; Zina, 2006). Also, choice of oviposition site plays an important role in the reproductive success of temporary pond amphibians (Resetarits and Wilbur, 1989). Various studies have suggested that female anurans select oviposition sites based on factors such as water depth (Caldwell, 1986), absence of predators (Howard, 1978; Resetarits and Wilbur, 1989), or water temperature (Howard, 1978). The suitability of a breeding site directly affects hatching success, larval performance, and, thereby, parental fitness (Resetarits and Wilbur, 1989). Thus, oviposition site-selection can have a strong impact on the spatial distribution of a species and thereby affect population dynamics (Hassel, 1987; Pearman and Wilbur, 1990) and the structure of communities (Morris, 2003). However, the mechanisms for oviposition site-selection in spatially and temporally variable environments are still poorly understood (Pearman and Wilbur, 1990).

Most *Pleurodema* species inhabit arid environments prone to strong seasonal fluctuations in temperature and precipitation, and are explosive breeders (Cei, 1980; Hödl, 1992; Cardoso and Arzabe, 1993). This temporal reproductive pattern is usually associated with marked spatial clumping around suitable oviposition sites (Wells, 1977). The explosive breeders form dense aggregations in which the males scramble for females. In this kind of breeders, struggles between males are common (Arak, 1983). For many anurans, male-male interactions consist of exclusively ter-

ritorial behaviours (Martins *et al.* 1998; Halloy and Espinosa, 2000), ranging from simple calls (warning, territorial, and/or encounter) to physical combat (Halliday and Tejedo, 1995). The scope of agonistic encounters varies and does not always result in physical confrontation, allowing the protagonists to avoid the risks and costs of bodily engagement (Kluge, 1981; Martins *et al.* 1998). The choice of behaviour and performance variation depends on the cost-benefit patterns of the behaviour and the communicated message (Enquist, 1985).

Pleurodema tucumanum PARKER 1927 is a small anuran, measuring between 40–45 mm snout-vent length (Cei, 1980), that inhabits central and northern Argentina (Vaira *et al.* 2012). In the central province of San Juan it is found exclusively in the Dry Chaco ecoregion of the Valle Fértil Department (Morrone, 2014). In this particular environment *P. tucumanum* often lives in sandy habitats, exhibiting burrowing behaviour, and breeding in temporary ponds formed by rains and/or flooding (Acosta comm. pers.). The reproduction of this species matches with the rainy season, in summer months (from November to April) (unpublished data). Ferraro *et al.* (2016) confirmed and described the final structure of the *P. tucumanum* foam nests. The process of construction of foam nests in Leiuperinae involves alternating and repeating phases of rest and activity. During the rest phases, the amplexant pair floated immobile, and in activity phases males beat the foam using their hindlimbs. There are several studies that describes this process in some *Pleurodema* species and other Leiuperinae (e.g. Hödl, 1990; 1992; Halloy and Fiaño, 2000; Valetti *et al.* 2014). However, there is no information about the process of construction in *P. tucumanum*. The objectives of this work were to 1) describe the structural characteristics of nesting microsites, 2) report the presence of agonistic and chorus behaviour among males of *P. tucumanum* and 3) document the construction process of these foam nests.

The study area is located in the region La Majadita (30.68°S, 67.50°W), Valle Fértil department, San Juan province, Argentina. This site is situated in the Arid Chaco ecoregion (Morrone, 2014). The climate is warm in the region with the mean annual temperature being 19.5 °C. The maximum mean temperature is 26 °C, and the minimum mean temperature is 11 °C. For this region, mean annual rainfall is recorded to be 358 mm, with means ranging between 152 and 656 mm (data provided by the Hydraulics Department of San Juan province, Argentina), with rains occurring predominantly in summer (December-March) (Cabrera, 1994).

The expedition was made on December 2013, following a small rain (3 mm; data provided by Servicio Meteorológico Nacional Argentino). During the trip, sampling was carried out through nocturnal walks along the river, using the visual encounter technique. At 2 am, we could observe a single temporary pond where were a lot of individuals of *P. tucumanum*. We sexed the individuals and counted them to obtain the abundance of individuals in this reproductive pond. We also measured the length and width of the pond, and maximum and minimum water depth using a tape (nearest 0.1 cm), and we made five different measures of water temperature to obtain an average of the temperature of the pond, using a thermocouple (TES TP-

K01; 1.62 mm diameter). In addition, we counted the number of amplexant pairs and foam nests located in this pond. In order to describe the foam nest temperatures and compare these temperatures with those of the amplexant pair and the air, we measured foam nest temperatures introducing a thermocouple (TES TP-K01; 1.62 mm diameter) 1 cm into the center of each foam nest. Air temperature was measured 1 cm above the foam using the same thermocouple, and body temperatures of the amplexant pair were measured using an infrared thermometer (GM550 Digital Laser Infrared Thermometer Temperature Measurement Electronic Point Gun) at a distance of 10 cm. In order to describe the process of construction of the foam nest, we recorded the whole process of construction using a digital camera (Nikon Coolpix S210V1.0). The time lapses for construction were obtained from the recorded video and were measured in minutes. While we were recording the foam nest construction, we also observed the behaviour of the remaining individuals in the reproductive pond. In addition to the nests recorded in December, during a subsequent sampling carried out in December 2015, we found new foam nests in other small bodies of water. For all observed foam nests (both found in December 2013 and in December 2015) we measured the following variables: depth of water under the nest, distance from the foam nest to the nearest shore, and maximum diameter of the foam nests, using a tape measure (nearest 0.1 cm).

Fifteen males and five females were observed in the pond. The dimensions of the pond were 435 cm by 86 cm, with a maximum depth of 7 cm and a minimum of 3.5 cm. The mean temperature of the pond water ($N=5$) was $20.7\text{ }^{\circ}\text{C}$ ($\pm 0.07\text{ }^{\circ}\text{C}$). From the total number of individuals, four of them were paired in amplexus. The remaining males were observed singing in chorus. In this reproductive pond, we observed three foam nests, one already constructed, and the others were seen since the beginning of the process of construction till the end (Table 1). We also observed the presence of agonistic male behaviour: we saw a combat between three males that involved frontal charging (Figure 1e) and included one climbing on and submerging another (Figure 1f), then, the third male attacked an amplexus pair. The aggression was repeated several times until the individual seemed to achieve his objective, interrupting the formation of the nest.

The foam nest construction process involved alternating phases of rest and activity (*sensu* Hödl, 1990). During the rest phases, the amplexant pair floated immobile. The male was in a position of axillary amplexus (Figure 1a–c), with his feet retracted and his cloaca upon that of the female. The active phase can be divided into three stages. The first stage began when the male slipped his feet between the thighs of the female and the cloacae of both were juxtaposed. The female laid her eggs while the male pushed his tarsal tubercles against the upward tilted cloacal region of the female and then rapidly brought his legs back to the initial position. In the second stage, the male beat the foam using alternating leg kicks. The movement involved the rotation of the femur and tibia perpendicular to the body axis, with the feet almost parallel to the body. The third and final stage began once a substantial amount of foam has been produced. Following the last kick, the male distributed the




Figure 1. Oviposition and reproductive behaviour of *Pleurodema tucumanum* in a temporary pond near La Majadita, Valle Fértil department, San Juan province, Argentina. **a–b** Amplexus and construction of foam nest and finalized nest corresponding to nest 1 (Table 1). **c–d** Amplexus and foam nest corresponding to nest 2 (Table 1). **e–f** Agonistic encounter between males. **g** Axillary amplexus.

foam by spreading his legs outward over the mass. The male then returned to the rest position. This behaviour was repeated until the foam nest was completed and egg deposition has finished.

In a subsequent sampling another 15 foam nest were recorded in other small bodies of water. The mean diameter of foam nests ($N=18$) was $104 \text{ mm} \pm 32.5 \text{ mm}$ (mean \pm SE), mean distance to the shore was $806.7 \text{ mm} \pm 63.4 \text{ mm}$ (mean \pm SE), and the mean water depth beneath the nests was $118 \text{ mm} \pm 63.3 \text{ mm}$ (mean \pm SE).

The mean foam nest temperature was similar to the mean water temperature, but always lower than the air temperature. Among temperate zone amphibians, aquatic egg masses are known to maintain temperatures slightly above the water surrounding them (Salthe and Mecham, 1974). All of the foam nests were located close to

the edges of the pond, in superficial water. Egg deposition and larval development in small temporary ponds markedly reduces vulnerability of eggs and tadpoles to aquatic predators (Duellman and Trueb, 1986; Hödl, 1986; Heyer *et al.* 1990). However, the use of those habitats as oviposition sites increases the vulnerability of developmental stages to desiccation (Zina, 2006). Eggs and larvae within foam nests placed in microsites that provide thermally favorable microclimate should develop somewhat faster than eggs and larvae within nests placed in cooler microsites (Dobkin and Gettinger, 1985), increasing the probability of tadpole survival.

Recorded construction times were very different between the two nests. The amplexant pair that dedicated more time to the foam nest construction produced a larger nest with more consistent foam (Figure 1a–b) than those that spent less time (Figure 1c–d). Such difference in foam nest size could be explained because of foam needs mechanical processing for the incorporation of air bubbles (Dalgetty and Kennedy, 2010). Thus nest size may depend on the time spent on construction. All males found in this study were singing in chorus, despite the fact that Martori *et al.* (1994) did not record the chorus behaviour in *P. tucumanum* in Córdoba Province. This behaviour constitutes a vocalized agonistic interaction with other males of the species with the objective of territorial establishment and female attraction (Wells, 1977; Penna *et al.* 2008; Velásquez *et al.* 2014). The agonistic behaviour found in this study has been observed in several pipids, leptodactylids, dendrobatids, hylids and ranids (e.g. Brattstrom and Yarnell, 1968; Rabb and Rabb, 1965; Wells, 1977; Halliday and Tejedo, 1995; Abrunhosa and Wogel, 2004), but there are no previous records of aggressive/agonistic combat behaviour in *Pleurodema* species.

Foam nest construction processes employed by *Pleurodema tucumanum* were analogous to those described for *P. diplolister* (Hödl, 1990) and *P. guayapae* (Valetti *et al.* 2014) and other Leiuperinae (e.g. Heyer and Rand, 1977; Dalgetty and Kennedy, 2010; Schaefer and Kehr, 2010). Male of *P. tucumanum* were observed utilizing their hind limbs to beat the foam, similar behaviour was found in *P. diplolister* (Hödl, 1992), *P. borellii* (Halloy and Fiaño, 2000), and *P. guayapae* (Valetti *et al.* 2014) and other Leiuperinae (e.g. Heyer and Rand, 1977; Dalgetty and Kennedy, 2010; Schaefer and Kehr, 2010). Similarly to our observations in *P. tucumanum*, in which during the rest phases the amplexant pair floated immobile and the male was in a position of axillary amplexus, axillary amplexus was observed in several Leiuperinae species (Barrio, 1964; Duellman and Veloso, 1977; Giaretta and Facure, 2009; Valetti *et al.* 2009; 2014; Velasco *et al.* 2017). The active phase can be divided into three stages that have been previously described for *Physalaemus ephippifer* (HÖDL, 1990), *Pleurodema diplolister* (Hödl, 1992), and *P. guayapae* (Valetti *et al.* 2014).

The data presented in this study verify the existence of *P. tucumanum* foam nest construction and confirms observations made by Ferraro *et al.* (2016). Surveys conducted days after oviposition led to determine that the ovoid plate nest proposed by



Figure 2. Observations on *Pleurodema tucumanum* in a temporary pond near La Majadita, Valle Fértil department, San Juan province, Argentina: **a** clutches found with low grade foam dilution **b** nest found with high grade of dilution, similar to the ovoid plate found by Martori *et al.* (1994) **c** clutch with larvae in Stage 22 of Gosner (1960).

Table 1. Data from variables measured in clutches from a temporary pond, near La Majadita, Valle Fértil department, San Juan province (30.68°S, 67.50°W), Argentina: foam nest temperature (Tfn), male temperature (Tm), female temperature (Tf), air temperature (Ta), time of construction and nest area. (*) No data obtained because the foam nest was already finished when found.

Nest	Tfn (°C)	Tm (°C)	Tf (°C)	Ta (°C)	Time of construction (min)
1	20.8	20.9	22.9	22.05	35
2	20.8	23.4	23.4	23.5	17
3*	20.4	–	–	22	–

Martori *et al.*, (1994) is the result of the dilution and collapse of the foam nest after several hours (Figure 2). Similar results were found by Ferraro *et al.* (2016) in the field and laboratory.

Pleurodema tucumanum is able to live in deserts and salty habitats (Ceï, 1980). The foam nests contain substances that delay larval development (Pisano, 1963), while protecting the eggs and early-stage embryo from dehydration (Prado *et al.* 2002; Zina, 2006). The *Pleurodema* genus appears to be a promising clade for evolutionary studies of physiological responses to extreme environmental conditions (Faivovich *et al.* 2012).

Our observations highlight the importance of studying reproductive behaviour systems. In addition, this study offers the first report of this type of reproductive behaviour strategies in Chacoan species, though further studies are needed to provide more detail for *P. tucumanum* reproductive behaviour such as the influence of both biotic and abiotic variables on oviposition site selection, timing of embryonic and larval development, the reproductive cycle, mating selection, among others.

Acknowledgments

We thank Rodrigo Nieva, Ana Paula Galdeano and Lia Piaggio Kokot for their help in fieldwork. We thank Jennifer Richardson and Magdalena Córdoba for their helpful comments on English language and we thanks Esteban Lavilla, Daiana Ferraro and the anonymous reviewers for comments and suggestions on earlier versions. We also thanks Secretaría de Ambiente y Desarrollo Sustentable Gobierno de San Juan, CICTCA-UNSJ and CONICET.

References

- ARAK, A. 1983. Male-male competition and mate choice in anuran amphibians. *In*: P. Bateson (eds.), *Mate choice*. Cambridge, United Kingdom, Cambridge University Press, p. 181–210.
- ABRUNHOSA, P.A.; WOGEL, H. 2004. Breeding behavior of the leaf-frog *Phyllomedusa burmeisteri* (Anura: Hylidae). *Amphibia-Reptilia*, **25**:125–135. <https://doi.org/10.1163/1568538041231157>
- BARRIO, A. 1977. Aportes para la elucidación del “status” taxonómico de *Pleurodema bibroni Tschudi* y *Pleurodema kriegi* (Müller) (Amphibia, Anura, Leptodactylidae). *Physis*, **37**: 311–331.
- BARRIO, A. 1964. Especies crípticas del género *Pleurodema* que conviven en una misma área, identificadas por el canto nupcial (Anura, Leptodactylidae). *Physis*, **24**:471–489.
- BRATTSTROM, B.H.; YARNELL, R.M. 1968. Aggressive behavior in two species of leptodactylid frogs. *Herpetologica*, **24**:222–228.
- CABRERA, A.L. 1994. Enciclopedia Argentina de Agricultura y jardinería, Tomo II, Fascículo 1: Regiones Fitogeográficas Argentinas. Buenos Aires, ACME, 85 p.
- CALDWELL, J.P. 1986. Selection of egg deposition sites: a seasonal shift in the southern leopard frog, *Rana sphenocephala*. *Copeia*, **1986**: 249–253. <https://doi.org/10.2307/1444923>
- CARDOSO, A.J.; ARZABE, C. 1993. Corte e desenvolvimento larvário de *Pleurodema diplostris* (Anura: Leptodactylidae). *Revista Brasileira de Biologia*, **53**: 561–570.
- CEI, J.M. 1980. *Amphibians of Argentina*. Monografía 2, Firenze, Monitore Zoologico Italiano, 609 p.
- DALGETTY, L.; KENNEDY, M.W. 2010. Building a home from foam túngara frog foam nest architecture and three-phase construction process. *Biology letters*, **6**: 293–296. <https://doi.org/10.1098/rsbl.2009.0934>
- DOBKIN, D.S.; GETTINGER, R.D. 1985. Thermal aspects of anuran foam nests. *Journal of Herpetology*, 271–275. <https://doi.org/10.2307/1564181>
- DUELLEMAN, W.E.; TRUEB, L. 1986. *Biology of amphibians*. New York, John Hopkins University Press, 670 p.
- DUELLEMAN, W.E.; VELOSO A.M. 1977. Phylogeny of *Pleurodema* (Anura: Leptodactylidae): a biographic model. *Occasional papers of the Museum of Natural History, University of Kansas*, **64**:1–46.

- ENQUIST, M. 1985. Communication during aggressive interactions with particular reference to variation in choice of behavior. *Animal Behaviour*, **33**:1152–1161. [https://doi.org/10.1016/S0003-3472\(85\)80175-5](https://doi.org/10.1016/S0003-3472(85)80175-5)
- FAIVOVICH, J.; FERRARO, D.P.; BASSO, N.G.; HADDAD, C.F.; RODRIGUES, M.T.; WHEELER, W.C.; LAVILLA, E.O. 2012. A phylogenetic analysis of *Pleurodema* (Anura: Leptodactylidae: Leiuperinae) based on mitochondrial and nuclear gene sequences, with comments on the evolution of anuran foam nests. *Cladistics*, **28**:460–482. <https://doi.org/10.1111/j.1096-0031.2012.00406.x>
- FERNÁNDEZ, K. 1927. Sobre la biología y reproducción de batracios argentinos. Segunda Parte. *Boletín de la Academia Nacional en Ciencias en Córdoba*, **29**: 271–328
- FERRARO, D.P.; PEREYRA, M.E.; BALDO D.; FAIVOVICH, J. 2016. The clutch structure of *Pleurodema tucumanum* (Anura: Leptodactylidae). *Salamandra*, **52**:48–52.
- GIARETTA, A.A.; FACURE, K.G. 2009. Habitat, egg laying behaviour, eggs and tadpoles of four sympatric species of *Pseudopaludicola* (Anura, Leiuperidae). *Journal of Natural History*, **43**: 995–1009. <https://doi.org/10.1080/00222930802702456>
- GOSNER, K.L. 1960. A simplified table for staging anuran embryos and larvae with notes on identification. *Herpetologica*, **16**:183–190.
- GRANT, T.; FROST, D.R.; CALDWELL, J.P.; GAGLIARDO, R.; HADDAD, C.F.B; KOK, P.J.R; MEANS, B.D.; NOONAN, B.P.; SCHARGEL, W.E.; WHEELER, W.C. 2006. Phylogenetic systematics of dart poison frogs and their relatives (Amphibia: Athesphatana: Dendrobatidae). *Bulletin of the American Museum of Natural History*, **299**:1–262. [https://doi.org/10.1206/0003-0090\(2006\)299\[1:PSODFA\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2006)299[1:PSODFA]2.0.CO;2)
- HADDAD, C.F.; PRADO, C.P. 2005. Reproductive modes in frogs and their unexpected diversity in the Atlantic Forest of Brazil. *BioScience*, **55**:207–217. [https://doi.org/10.1641/0006-3568\(2005\)055\[0207:RMIFAT\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2005)055[0207:RMIFAT]2.0.CO;2)
- HALLIDAY, T.; TEJEDO, M. 1995. Intrasexual selection and alternative mating behaviour. *Amphibian Biology*, **2**:419–468.
- HALLOY, M.; ESPINOSA, R.E. 2000. Territorial encounters and threat displays in the neotropical frog *Phyllomedusa sauvagii* (Anura: Hylidae). *Herpetological Natural History*, **7**:175–180.
- HALLOY, M.; FIAÑO, M. 2000. Oviposition site selection in *Pleurodema borellii* (Anura: Leptodactylidae) may be influenced by tadpole presence. *Copeia*, **2000**:606–609. [https://doi.org/10.1643/0045-8511\(2000\)000\[0606:OSSIPB\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2000)000[0606:OSSIPB]2.0.CO;2)
- HASSEL, M.P.1987. Detecting regulation in patchily distributed animal populations. *Journal of Animal Ecology* **56**:705–731. <https://doi.org/10.2307/5078>
- HEYER, W.R.; RAND, A.S. 1977. Foam nest construction in the leptodactylid frogs *Lepidodactylus pentadactylus* and *Physalaemus pustulosus* (Amphibia, Anura, Leptodactylidae). *Journal of Herpetology*, **11**: 225–228. <https://doi.org/10.2307/1563148>
- HEYER, W.R.; RAND, A.S.; DA CRUZ, C.A.G.; PEIXOTO, O.L.; NELSON, C.E. 1990. Frogs of Boracéia. *Arquivos de Zoologia (São Paulo)*, **31**: 231–410. <https://doi.org/10.2307/1446606>
- HÖDL, W. 1986. Foam nest construction in South American leptodactylid frogs. *Studies in herpetology*, **565–570**. <https://doi.org/10.2307/1563490>

- HÖDL, W. 1990. An analysis of foam nest construction in the Neotropical frog *Physalaemus ephippifer* (Leptodactylidae). *Copeia*, **1990**:547–554. <https://doi.org/10.2307/1446358>
- HÖDL, W. 1992. Reproductive behavior in the neotropical foam-nesting frog *Pleurodema diplolistris* (Leptodactylidae). *Amphibia-Reptilia*, **13**:263–274. <https://doi.org/10.1163/156853892X00472>
- HOWARD, R.D. 1978. The influence of male-defended oviposition sites on early embryo mortality in bullfrogs. *Ecology*, **59**: 789–798. <https://doi.org/10.2307/1938783>
- KLUGE, A.G. 1981. The life history, social organization, and parental behavior of *Hyla rosenbergi* Boulenger, a nest-building gladiator frog. *Miscellaneous publications, Museum of Zoology, University of Michigan*, **160**:11–70.
- MARTINS, M.; POMBAL, J.P.; HADDAD, C.F. 1998. Escalated aggressive behaviour and facultative parental care in the nest building gladiator frog, *Hyla faber*. *Amphibia-Reptilia*, **19**:65–73. <https://doi.org/10.1163/156853898X00331>
- MARTORI, R.; AÚN, L.; VIGNOLO, P.E. 1994. Aporte al conocimiento de la biología y distribución de *Pleurodema tucumanum* (Anura, Leptodactylidae). *Boletín de la Asociación Herpetológica Argentina*, **10**:18–19.
- MORRIS, D.W. 2003. Toward an ecological synthesis: a case for habitat selection. *Oecologia*, **136**: 1–13. <https://doi.org/10.1007/s00442-003-1241-4>
- MORRONE, J.J. 2014. Biogeographical regionalization of the Neotropical region. *Zootaxa*, **3782**:1–110. <https://doi.org/10.11646/zootaxa.3782.1.1>
- PEARMAN, P.B.; WILBUR, H.M. 1990. Changes in population dynamics resulting from oviposition in a subdivided habitat. *The American Naturalist*, **135**: 708–723. <https://doi.org/10.1086/285070>
- PENNA, M.; VELÁSQUEZ, N.; SOLÍS, R. 2008. Correspondence between evoked vocal responses and auditory thresholds in *Pleurodema thaul* (Amphibia; Leptodactylidae). *Journal of Comparative Physiology A*, **194**:361–371. <https://doi.org/10.1007/s00359-007-0312-0>
- PISANO, A. 1963. La espuma de los nidos de *Leptodactylus bufonius* y su acción inhibidora sobre el desarrollo de las larvas. *Archivos de Bioquímica, Química y Farmacia*, **10**:65–77.
- PRADO, C.P.A.; UETANABARO M.; HADDAD C.F. 2002. Description of a new reproductive mode in *Leptodactylus* (Anura, Leptodactylidae), with a review of the reproductive specialization towards terrestriality in the genus. *Copeia*, **2002**:1128–1133. [https://doi.org/10.1643/0045-8511\(2002\)002\[1128:DOANRM\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2002)002[1128:DOANRM]2.0.CO;2)
- PYRON, R.A.; WIENS, J.J. 2011. A large-scale phylogeny of Amphibia including over 2,800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics and Evolution*, **61**:543–583. <https://doi.org/10.1016/j.ympev.2011.06.012>
- RABB, G.; RABB, M.S. 1965. Effects of isolation on reproductive behavior in the pipid frog *Xenopus laevis*. *American Zoology*, **5**:685.
- RESEARITS, W.J.; WILBUR, H.M. 1989. Choice of oviposition site by *Hyla chrysoscelis*: role of predators and competitors. *Ecology*, **70**: 220–228. <https://doi.org/10.2307/1938428>
- SALTHER, S.N.; MECHAM, J.S. 1974. Reproductive and courtship patterns. *Physiology of the Amphibia*, **2**: 309–521. <https://doi.org/10.1016/B978-0-12-455402-3.50010-3>

- SCHAEFER, E.F.; KEHR, A.I. 2010. Comportamiento reproductivo de *Physalaemus albonotatus* (Steindachner, 1864) (Anura, Leiuperidae) en el nordeste argentino. *Revista Española de Herpetología*, **24**:27–39.
- SEYMOUR, R.S.; LOVERIDGE, J.P. 1994. Embryonic and larval respiration in the arboreal foam nests of the African frog *Chiromantis xerampelina*. *Journal of Experimental Biology*, **197**:31–46.
- SPIELER, M.; LINSENMAIR, K.E. 1997. Choice of optimal oviposition sites by *Hoplobatrachus occipitalis* (Anura: Ranidae) in an unpredictable and patchy environment. *Oecologia*, **109**: 184–199. <https://doi.org/10.1007/s004420050073>
- VAIRA, M.; AKMENTINS, M.; ATTADAMO, M.; BALDO, D.; BARRASSO, D.; BARRIONUEVO, S.; BASSO, N.; BLOTTO, B.L.; CAIRO, S.; CAJADE, R.; CÉSPEDEZ, J.; CORBALÁN, V.; CHILOTE, P.; DURÉ, M.; FALCIONE, C.; FERRARO, D.; GUTIÉRREZ, R.; INGARAMO, M.; JUNGES, C.; LAJMANOVICH, R.; LESCOANO, J.N.; MARANGONI, F.; MARTINAZZO, L.; MARTI, R.; MORENO, L.; NATALE, G.; PÉREZ IGLESIAS, J.; PELTZER, P.; QUIROGA, L.; ROSSET, S.; SANABRIA, E.; SANCHEZ, L.; SCHAEFER, E.F.; ÚBEDA, C.A.; ZARACHO, V.H. 2012. Categorización del estado de conservación de los anfibios de la República Argentina. *Cuadernos de Herpetología*, **26**:131–159.
- VALETTI, J.; GRENAT, P.R.; BARAQUET, M.; MARTINO, A.L. 2014. Reproductive biology of *Pleurodema guayapae* (Anura: Leptodactylidae: Leiuperinae). *International Journal of Tropical Biology and Conservation*, **62**:173–182. <https://doi.org/10.15517/rbt.v62i1.7763>
- VALETTI, J.A.; SALAS, N.E.; MARTINO, A.L. 2009. A new polyploid species of *Pleurodema* (Anura: Leiuperidae) from Sierra de Comechingones, Córdoba, Argentina and redescription of *Pleurodema kriegi* (Müller, 1926). *Zootaxa*, **2073**:1–21. <https://doi.org/10.5281/zenodo.187133>
- VELASCO, M.A.; ÚBEDA, C.A.; WILLIAMS, J.D.; KACOLIRIS, F.P. 2017. Reproductive biology of the critically endangered valcheta frog, *Pleurodema somuncurensis* (Anura: Leptodactylidae), from Patagonia, Argentina. *South American Journal of Herpetology*, **12**: 205–211. <https://doi.org/10.2994/SAJH-D-16-00049.1>
- VELÁSQUEZ, N.A.; OPAZO, D.; DÍAZ, J.; PENNA, M. 2014. Divergence of acoustic signals in a widely distributed frog: relevance of inter-male interactions. *PloS one*, **9**: e87732. <https://doi.org/10.1371/journal.pone.0087732>
- WEIGANDT, M.; ÚBEDA, C.A.; DIAZ, M. 2004. The larva of *Pleurodema bufoninum* Bell, 1843, with comments on its biology and on the egg strings (Anura, Leptodactylidae). *Amphibia-Reptilia*, **25**:429–437.
- WELLS, K.D. 1977. The social behaviour of anuran amphibians. *Animal Behaviour*, **25**:666–693. [https://doi.org/10.1016/0003-3472\(77\)90118-X](https://doi.org/10.1016/0003-3472(77)90118-X)
- ZINA, J. 2006. Communal nests in *Physalaemus pustulosus* (Amphibia: Leptodactylidae): experimental evidence for female oviposition preferences and protection against desiccation. *Amphibia-Reptilia*, **27**:148–150. <https://doi.org/10.1163/156853806776052092>

Resumo

Estratégias reprodutivas são combinações de características fisiológicas, morfológicas e comportamentais que interagem para aumentar o sucesso reprodutivo em certas condições ambientais. Embora as estratégias reprodutivas do Leiuperinae sejam conhecidas, poucos estudos foram realizados sobre os comportamentos reprodutivos que as sustentam. O objetivo deste trabalho foi documentar as características estruturais dos microsítios de oviposição, descrever o processo de construção do ninho de espuma e registrar a presença de comportamento agonístico e de coro em machos de *Pleurodema tucumanum*. Os ninhos foram encontrados próximos às margens de uma poça temporária e a temperatura média dos ninhos de espuma esteve sempre próxima da temperatura média da água. Nossas observações corroboram que as fases de construção do ninho de espuma para *P. tucumanum* são similares àquelas descritas para outros Leiuperinae, incluindo a dispersão e o batimento da espuma realizada pelos membros posteriores do macho. Também registramos pela primeira vez a observação de encontros agonísticos entre machos de *Pleurodema tucumanum*.

Palavras-chave

Anfíbios, Chaco Seco, Interações entre machos, modo reprodutivo, sítio de oviposição.