



FAUNISTIC NOTE

Small flies with high conservation value: first reliable record of *Hyperoscelis veternosa* (Diptera, Canthyloscelidae) from Romania supported by DNA barcode data

Lujza Keresztes¹, Andrei Bogdan Terec^{1,2}, Boróka-Zsuzsánna Jancsó^{1,2}, Avar
Lehel Dénes^{1,3}, Anna Dénes^{1,2}

1 Centre 3B, Laboratory of Advance Hydrobiology and Biomonitoring, Babeş-Bolyai University, Clinicilor 5–7, Cluj-Napoca 400006, Romania

2 Doctoral School of Integrative Biology, Babeş-Bolyai University, Republicii 44, Cluj-Napoca 400015, Romania

3 Institute of Interdisciplinary Research in Bio-Nano-Sciences, Babeş-Bolyai University, Treboniu Laurian 42, Cluj-Napoca 400271, Romania

Corresponding author: Andrei Bogdan Terec (andrei.terec@ubbcluj.ro)

Received 19 April 2023 | Accepted 31 May 2023 | Published 30 June 2023

Citation: Keresztes L, Terec AB, Jancsó B-Z, Dénes AL, Dénes A (2023) Small flies with high conservation value: first reliable record of *Hyperoscelis veternosa* (Diptera, Canthyloscelidae) from Romania supported by DNA barcode data. Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" 66(1): 165–171. <https://doi.org/10.3897/travaux.66.e105107>

Abstract

Here we present the first reliable record of the rare *Hyperoscelis veternosa* Mamaev and Krivosheina, 1969 from the headwaters area of the Someşul River, Apuseni Natural Park, Romania. Species level identification was made by detailed examination of the single female genitalia, supported by DNA barcode data. The new finding confirms the presence of the species in Romania, whereas a very old, single and doubtful record from the Bucharest area dates back to 1900 needs further confirmation. The presence of the second species of genus *Hyperoscelis* from Romania is of high conservation value, and confirm the persistence of a relict biodiversity of an old-growth forests-spot in the Carpathians.

Keywords

Canthyloscelidae, conservation, DNA barcode data, new record, relict species, the Carpathians, virgin forests.

The family Canthyloscelidae is a small group of Scatopsoidea flies, which has only 12 species worldwide with scattered or “relict-like” distribution in the Holarctic Region (6 species), southern South America (2 spp.) and New Zealand (4 spp.) (Hutson 1977; Martinovsky 1988; Haenni 1997; Oosterbroek 2006; Kjærandsen and Jaschhof 2019). According to the most recent revisions by Amorim (2000) and Ševčík et al. (2016), the family includes the subfamilies Synneurinae (with two genera, *Synneuron* Lundström, 1910 (with 3 spp.) and *Exiliscelis* Hutson, 1977 – with 1 sp.) and Canthyloscelinae (with two genera, *Hyperoscelis* Hardy and Nagatomi, 1960 (with 2 spp.) and *Canthyloscelis* Edwards, 1922 (with 6 spp.)). In Europe only three species occur so far, *Hyperoscelis eximia* (Boheman, 1858), *Hyperoscelis veternosa* Mamaev and Krivosheina, 1969, and *Synneuron annulipes* Lundström, 1910 with rather sporadic data (Martinovsky 1972; Haenni and Dufour 1983; Klasa 1991; Mikołajczyk 1993; Schacht 1993).

Members of the family are considered remnants of an ancient fauna restricted to some old growing primeval forests, with rarely collected adults and larvae detected from wet, rooting woods and stumps near waters (Roháček and Ševčík 2009; Marshall 2012).

The first record of Hyperoscelidae from Romania dates back to the beginning of the 20th century, when Mik (1900) recorded the species *Corynoscelis eximia* (Boheman, 1858; valid name: *H. eximia*), based on a single male specimen collected by Kiefer from the surroundings of Bucharest, Comana forest. The specimen was later revised and identified by Hutson (1977) as *H. veternosa* (Ceianu, 1998). The single known locality of the species *S. annulipes* from Romania is Mehadia (Banat region) and were cited by Zilahi-Sebess (1960) without any more exact collecting data. The third species, *H. eximia* were recorded by Ceianu (1998) from the Eastern Carpathians close to the Ukrainian border (Câmpulung Moldovenesc and Sadău) based on one male and one female individuals. Apart from these few records, no further data on this relict family in Romania are known so far.

Material examined. 1 female, Bihor county, Ic Ponor on the road to Padiș, Apuseni Mts., Romania, spruce fir forest in marshy area, 46.631716° N/22.805208° E, 1035 m a.s.l.; 7 May, 2021, Keresztes Lujza leg. The specimen (Fig. 1) was collected by sweeping the fallen roots and stumps from a marshy area near the spring sector of the river Someșul Cald. The specimen, with individual code DICN_SC2a_01, is deposited in the Diptera Collection of the Faculty of Biology and Geology (acronym DCFBG), University Babeș-Bolyai, Cluj-Napoca, Romania. Female terminalia were left overnight in 10% KOH and thereafter one hour in undiluted glacial acetic acid to neutralize and wash out the soap that was created from the soft tissues. Then, the male terminalia were transferred to a larger amount of glycerol in order to wash out the acid, and then to a drop of glycerol on a slide with rounded excavation. The slide was carefully transferred to the compound microscope in order to take the photos. Finally, the genital parts were placed in micro vials containing 50–50% ethanol and glycerol and were pinned under the dry specimen. Photos of the wing were taken




Figure 1. *Hyperoscelis veternosa*, female specimen DICN_SC2a_01. **A** Specimen, lateral view; **B** Right wing of female; **C** Abdomen terminal segments, dorsal view; **D** Abdomen terminal segments, ventral view.

with a stereomicroscope (Olympus SZ51) with a Canon EOS 650D digital camera attached. Photos were taken using a Canon EOS 750D digital camera, attached to the microscope (Olympus CX23), with an LM Digital SLR Adapter (MicroTechLab, Austria). Layer photos were combined using Zerene Stacker software.

Stacking results consist of 10–15 single exposures with the stereomicroscope and 20–50 exposures with the compound microscope.

One leg from the female specimen was sent to the Canadian Centre for DNA barcoding (Guelph, Ontario, Canada), for DNA extraction and bi-directional Sanger sequencing as part of the Romanian Macrozoobenthos Barcoding Initiative (ROMAC) in Barcode of Life Data System (BOLD). Publicly available *Hyperoscelis* sequences were downloaded from the BOLD database and a Neighbour-Joining tree was generated using 10,000 bootstrap replicates in Mega X (Kumar et al. 2018).

The single female collected by us is the second record of *H. veternosa* in Romania, after the first record of Mik (1900), under the name *Corynoscelis eximia* Boheman (see Hutson 1977 for discussion). Our data is the first reliable record of the species in Romania, and the first record of the species from the Apuseni National Park, Apuseni Mts. The habitat where the female specimen was collected is typical to *H. veternosa*, and is characterized by a large marshy area near the main course of the Someşul

Cald River, above 1000 m a.s.l. (Fig. 2). The area belongs to the Apuseni National Park, with restricted human activity in an old-growth spruce and fir forest with large amounts of fallen trees and dead wood accumulated as a result of undisturbed, natural forest succession.

The specimen collected by us was DNA-barcoded, and as a result, a 658 bp length sequence (the standard DNA region of the mitochondrial cytochrome oxidase gene *mtCOI*) was obtained and assigned to the Barcode Index Number (BIN) in BOLD: ADJ6424. Our DNA sequence was compared with the only two other additional sequences available on public genetic databases. The DNA sequence mined from GenBank (NCBI) (sample ID KT316865) and published by Ševčík et al. (2016) has 99,67% genetic similarity with our genetic dataset, while the second public DNA data mined from BOLD (sample ID NORSC2945-18) has 100% similarity with our data. The *H. veternosa* is the sister species of *H. eximia*, based on standard DNA data in BOLD with BIN number: ADS5898, having 11.89% distance to its nearest neighbour (Fig. 3).



Figure 2. Collecting locality of the specimen represents the reliable record of *Hyperoscelis veternosa* in Romania: Ic Ponor, the Apuseni Mountains, marshy spruce-fir forest along the main course of the Someşul Cald River, at 1035 m a.s.l. (46.631716° N/22.805208° E, 1035 m a.s.l.), leg. Keresztes L.




Figure 3. NJ-tree (Kimura-2-distance) with all barcodes available for the European Canthyloscelidae fauna, with their associated BIN numbers. The single specimen collected by us is marked in bold.

Our new data are of great conservation value, as the species has rather sporadic data across Europe and is closely related with primeval forest-type associations, unique ecosystems that are disappearing from most parts of our continent. The presence of the species in the perimeter of the Apuseni National Park protected area suggest the presence of a still highly diverse virgin forests in the Carpathians. The spruce-fir and old-growth beech forests around the headwaters of the Someșul Cald River area needs special conservation efforts and good forest management, as the ongoing global climate change with catastrophic meteorological events affects seriously the area (as it was, for example, in 2017 or in 2022, resulted in massive windfalls of timbers) in combination with the anthropogenic pressure (pasturing, tourism), which will decimate, for sure, these unique ecosystems in the forthcoming decades. The newly discovered *H. veternosa* should be a good indicator species of a still pristine forest ecosystem in this area, therefore we suggest including it, together with the sister species, *H. eximia* in the national red-lists of endangered species with high conservation value.

Acknowledgements

The present work received financial support from a national grant offered by the Executive Unit for Financing Higher Education, Research, Development and Innovation, UEFSCDI, of the Romanian Ministry of Education and Research, PN-III-P2-2.1-PED-2019-0214; nr. 476PED/2020.

References

- Amorim DS (2000) A new phylogeny and phylogenetic classification for the Canthyloscelidae (Diptera: Psychodomorpha). *Canadian Journal of Zoology* 78(6): 1067–1077. DOI:10.1139/z00-010.
- Boheman CH (1858) Bidrag till Lapplands dipter-fauna. Öfversikt Kungliga Vetenskapsakademien Forhandlingar. 15: 55–57. [in German]
- Ceianu I (1998) Genul *Hyperoscelis* Hardy and Negrotomi, 1960 (Diptera: Nematocera: Canthyloscelidae) în România. *Buletin de Informare a Societății lepidopterologice Române* 9(3–4): 283–290. [in Romanian]
- Edwards FW (1922) XXXIII. — Preliminary note on a new genus of Scatopsid flies from New Zealand. *The Annals and magazine of natural history; zoology, botany, and geology* 9: 267–269. DOI:10.1080/00222932208632673.
- Hutson AM (1977) A revision of the families Syneuridae and Canthyloscelidae (Diptera). *Bulletin of the British Museum (Natural History) Entomology* 35(3): 67–100. DOI:10.5962/bhl.part.4753.
- Haenni JP (1997) Family Canthyloscelidae. In Papp L, Darvas B (Eds): *Contributions to a Manual of Palaearctic Diptera. Nematocera and Lower Brachycera. Vol. 2.* Science Herald, Budapest 273–279.
- Haenni JP, Dufour C (1983) Première capture en Suisse d'un représentant de la famille raliqne des Canthiloscelidae (Diptera, Nematocera). *Bulletin de la Société Entomologique Suisse* 56: 187–189. [in French]
- Hardy DE, Nagatomi A (1960) An unusual new Nematocera from Japan (Diptera) and a new family name. *Pacific Insects* 2: 263–269.
- Kjærandsen J, Jaschof M (2019) New records and first DNA barcodes of the family Canthyloscelidae (Diptera) in Fennoscandia. *Norwegian Journal of Entomology* 66: 81–93.
- Klasa A (1991) *Hyperoscelis eximia* (Boheman) (Diptera, Canthyloscelidae). A representative of the new to the Polish fauna family of Nematoceros Flies. *Annals of the Upper Silesian Museum - Entomology* 2: 217–220.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. DOI:10.1093/molbev/msy096

- Mamaev BM, Krivosheina NP (1969) New data on the morphology and ecology of the Hyperoscelidae (Diptera, Nematocera) (in Russian). *Entomologicheskoe Obozrenje* 48(4): 933–942.
- Marshall SA (2012) *Flies: The natural history and diversity of Diptera*. Firefly Books Ltd, p. 136.
- Martinovsky J (1972) *Hyperoscelis eximia* (Boheman) in der Tschechoslowakei und die systematische übersicht der Familie Hyperoscelidae (Diptera). *Acta entomologica bohemoslovaca* 69: 351–354. [in German]
- Martinovsky J (1988) Czechoslovak species of the family Canthyloscelidae (Diptera). *Zpravy čes. Spol. Ent.* 24: 29–32.
- Mik J (1900) Dipterologische Miscellen. (2. Serie)- Wiener Entomologische Zeitung - 19: 71–76. [in German]
- Mikołajczyk W (1993) *Hyperoscelis eximia* (Bohem.) its new localities of Poland. *Polskie Pismo Entomologiczne* 62: 289–290.
- Oosterbroek P (2006) *The European Families of the Diptera*. KNNV Publishing, Utrecht, 205 pp. DOI: 10.1163/9789004278066.
- Roháček J, Ševčík J (2009) Opetiidae, Platypezidae. In: Roháček J, Ševčík J (Eds) *Diptera of the Poľana Protected Landscape Area – Biosphere Reserve (Central Slovakia)*. SNC SR, Administration of the PLA – BR Poľana, Zvolen, 150–155.
- Schacht W (1993) Fundort Schöngesing Erstnachweis einer Fliegenfamilie für Deutschland (Diptera, Canthyloscelidae). *Entomofauna* 14(6): 81–84. [in German]
- Ševčík J, Kasprák D, Mantič M, Fitzgerald S, Ševčíková T, Tóthová A, Jaschhof M (2016) Molecular phylogeny of the megadiverse insect infraorder Bibionomorpha sensu lato (Diptera). *PeerJ* 4:e2563. DOI: 10.7717/peerj.2563.
- Zilahi-Sebess G (1960) *Fonalascsapúak I. - Nematocera I. (Magyarország állatvilága - Fauna Hungariae* 55. XIV. kötet, Diptera I., 2. füzet). [in Hungarian]