



FAUNISTIC NOTE

# Trigonalynoidea (Hymenoptera: Apocrita) – a new superfamily of wasps recorded in Romania

Alexandru-Mihai Pintilioaie<sup>1</sup>

<sup>1</sup> SC Biodiversity Research and Consulting SRL, Iași, Romania

Corresponding author: *Alexandru-Mihai Pintilioaie* (alexandrupintilioaie@gmail.com)

Received 7 March 2022 | Accepted 19 May 2022 | Published 30 June 2022

**Citation:** Pintilioaie A-M (2022) Trigonalynoidea (Hymenoptera: Apocrita) – a new superfamily of wasps recorded in Romania. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 65(1): 67–70. <https://doi.org/10.3897/travaux.64.e83252>

## Abstract

The superfamily Trigonalynoidea, along with the species *Pseudogonalos hahnii* (Spinola, 1840), is recorded for the first time in Romania. An up-to-date distribution in Europe and a brief description of its biology is presented, together with a picture of the specimen collected in Romania.

## Keywords

distribution, Europe, hyperparasitism, Insecta, trigonalynids.

Trigonalynoidea is a small superfamily of parasitic wasps, comprising 16 genera and around 120 species, all belonging to a single family – Trigonalynidae (Chen et al. 2020). Various authors spell the name of the family either as Trigonalidae or Trigonalynidae, but the latter is used here (for details see the paper of Engel and Lelej 2020).

The distribution of Trigonalynidae is a cosmopolitan one (aside from alpine and arctic zones), with a maximum number of species occurring in the tropics. Europe has only one species, namely *Pseudogonalos hahnii* (Spinola, 1840) (Fig. 1), which is also present in Asia (China, Kazakhstan, Mongolia and Siberia). In Europe, *P. hahnii* is known from Belgium, Czech Republic, Estonia, Finland, France, Germany, Greece, Italy, Latvia, Lithuania, the Netherlands, Poland, Russia, Slovakia, Switzerland, Ukraine and the United Kingdom (Väänänen et al. 2018). Although the species is known from most of the European countries, it appears to be collected only in small numbers, being relatively rare (Broad 2016; Väänänen et al. 2018). This scarcity of



**Figure 1.** *Pseudogonalos hahnii* (Spinola, 1840), habitus of the specimen collected in Romania.

records can be partially explained by the very short life span of the adults, which seems to be at most eight days (Carmean 1991).

*Pseudogonalos hahnii* was found to be a eurytopic species, being collected in both xerothermous and moist habitats (Schnee 2011).

The trigonalynoids have a very interesting biology. The female of *Pseudogonalos hahnii* lays thousands of tiny eggs on various plants (*Artemisia vulgaris* L., *Epilobium angustifolium* L., *Phragmites australis* (Cav.) Trin. ex Steud., *Pinus* sp., Poaceae, *Prenanthes purpurea* L., *Pteridium aquilinum* (L.) Kuhn, *Rubus* sp., *Urtica dioica* L., *Vaccinium myrtillus* L., etc), which needs to be consumed by a secondary host (sawfly or lepidoptera larvae) in order to hatch (Väänänen et al. 2018). It was shown that the ovarioles of *Pseudogonalos hahnii* can contain more than 10.000 eggs and it can lay more than 1000 eggs per day in laboratory conditions (Bischoff 1936). After hatching in the secondary host gut, trigonalyno larva searches for its primary host, which is represented by Ichneumonidae larvae (Insecta: Hymenoptera), in the case of *Pseudogonalos hahnii* (Väänänen et al. 2018). The known species of both primary and secondary hosts of *Pseudogonalos hahnii* are summarized in Väänänen et al. (2018).

**Material examined:** Romania: 1 specimen (♀); Bacău county, Comănești; 46.4286°N/26.4399°E; 472 m alt.; 15 June 2020; Pintilioaie Alexandru-Mihai leg. The specimen was collected from herbaceous vegetation using a sweeping net and is preserved in the personal collection of the author, housed in Iași, Romania.

## Acknowledgements

The author is thankful to Steve Compton and Theo Peeters for providing relevant papers on the subject; to Alexandra Florina Popa for providing useful comments regarding the presence of the species in Romania; to Christian Schmid-Egger and Arjan Stroo for helping with the identification of the specimen; to Cosmin Mancu for helping with the picture of the specimen, and also to the anonymous reviewers who improved the first version of this paper.

## References

- Bischoff H (1936) Beiträge zur Lebensgeschichte der *Pseudogonalos hahni* (Spin.). Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1–3: 51–63 [in German].
- Broad G (2016) Checklist of British and Irish Hymenoptera - Trigonalynoidea. Biodiversity Data Journal 4: e7935. doi: 10.3897/BDJ.4.e7935
- Carmean D (1991) Biology of the Trigonalynoidea (Hymenoptera), with notes on the vespine parasitoid *Bareogonalos canadensis*. New Zealand Journal of Zoology 18(2): 209–214. <https://doi.org/10.1080/03014223.1991.10757968>

- Chen H-Y, Hong C-D, Achterberg C van, Pang H (2020) New species and new records of Trigonalysidae (Hymenoptera) from Tibet, China. *ZooKeys* 918: 83–98. <https://doi.org/10.3897/zookeys.918.49729>
- Engel MS, Lelej AS (2020) On the spelling of family-group names based on the genus *Trigonalys* Westwood (Hymenoptera: Trigonalidae versus Trigonalysidae). *Entomologist's Monthly Magazine* 156: 58–60. <https://doi.org/10.31184/M00138908.1561.3989>
- Schnee H (2011) Beitrag zur Kenntnis der Biologie von *Pseudogonalos hahnii* (Spinola) (Hymenoptera, Trigonalidae und Ichneumonidae). *Entomologische Nachrichten und Berichte* 55: 27–32. [In German].
- Väänänen S, Paukkunen J, Soon V, Budrys E (2018). Occurrence and biology of *Pseudogonalos hahnii* (Spinola, 1840) (Hymenoptera: Trigonalidae) in Fennoscandia and the Baltic states. *Entomologica Fennica*, 29(2): 86–96. <https://doi.org/10.33338/ef.71220>