

Range dynamics of *Walterinnesia morgani* (Serpentes, Elapidae) during climatic oscillations in Iran

Ali Gholamifard¹, Mehmet Kürşat Şahin²

¹ Department of Biology, Faculty of Sciences, Lorestan University, 6815144316 Khorramabad, Iran

² Department of Biology, Faculty of Science, Hacettepe University, Ankara, Türkiye

<https://zoobank.org/69B225B8-6965-4682-BB64-3A5332803B37>

Corresponding author: Ali Gholamifard (gholamifard.a@lu.ac.ir)

Academic editor: Yurii Kornilev ♦ Received 16 June 2023 ♦ Accepted 20 November 2023 ♦ Published 18 December 2023

Abstract

Reptiles have a crucial part in maintaining global biodiversity and the functioning of dynamic ecosystems, owing to their ecological roles and functions. Nevertheless, these organisms are susceptible to human-induced disruptions and the deterioration of their habitats, leading to their categorization as the third most endangered group of vertebrates on a global scale. Understanding the spatial distribution of reptiles is crucial due to their often specific habitat needs and limited vagility. Morgan's black cobra (*Walterinnesia morgani*) is a secretive venomous snake species that has thus far received little attention in Iranian scientific literature. The aim of the present study was to determine the existing distribution pattern of the cobra and to speculate on how climatic changes might affect it. Maximum entropy modeling was used to examine a dataset consisting of 16 occurrence records gleaned from field observations and the literature. The niche of the species was predicted using current and future climate change forecasts and bioclimatic and topographical characteristics. The models predicted a future reduction in the wide distribution region of *W. morgani* in southern and western Iran. It was discovered that climatic factors like temperature range, precipitation dynamics, and river proximity all played a key role in shaping the pattern of distribution. The predicted suitable areas for *W. morgani* were dependent on water sources; however, future scenarios showed a decline in suitable habitats. This study underscores the importance of conservation efforts in light of the potential implications of climate change on this species. To further understand the range shifts and adaptive strategies of the species, further study of its ecology and dispersal dynamics is required.

Key Words

ecological niche, MaxEnt, Morgan's black cobra, precipitation, topography

Introduction

Reptiles, although they account for one-third of global terrestrial vertebrate diversity, have lagged behind other groups, such as birds and mammals, in terms of ecological studies (Biber et al. 2023). Due to the fact that reptiles are typically characterized by specific habitat needs and limited vagility (Guedes et al. 2018), understanding their spatial distribution is crucial. In this regard, revealing species distribution patterns improves our knowledge of the relationship between landscape and species. Climatic oscillations, like topographic and vegetation cover patterns, have contributed to the establishment of the distribution

of species in different biomes (Bobrowski et al. 2018; Şahin et al. 2022a).

Although reptile species are widely distributed throughout the world, they are at risk of extinction due to habitat loss and degradation, pollution, invasive species, diseases, and climate change (Gibbons et al. 2000). Anthropogenic impacts—not only habitat use but also misinformation about their venomosity—have disproportionately threatened snake species (Saha et al. 2018). Therefore, evaluating the conservation status of these quite secretive organisms and determining their distribution patterns are important.

The herpetofauna of Iran comprises 81 species of snakes belonging to 34 genera and 7 Families (Rajabizadeh 2018).

Family Elapidae, including Elapinae (3 genera, 3 species) and Hydrophiinae (1 genus, 10 species), is more diverse than Viperidae, including Crotalinae (1 genus, 1 species) and Viperinae (7 genera, 10 species) (Safaei-Mahroo et al. 2015; Rajabizadeh 2018). Despite increasing efforts to determine herpetofaunal distributions in Iran (Hosseini Yousefkhani 2019; Vaissi 2021a, b, 2022), snake species remain understudied (Hosseinzadeh et al. 2017; Moradi et al. 2021).

The genus *Walterinnesia* Lataste, 1887 has two species known commonly as desert black snakes or black desert cobras including the type species *Walterinnesia aegyptia* Lataste, 1887 and *Walterinnesia morgani* (Mocquard, 1905) (Fig. 1). Both species occur only in the Middle East. The type locality of *W. morgani* is Khuzestan Province, southwestern Iran (Uetz et al. 2023). Populations of the species in Iran were previously considered as *W. aegyptia* by Farzanpay (1989), Latifi (1985, 1991, 2000), and Leviton et al. (1992). Nilson and Rastegar-Pouyani (2007), based on the lower scale row counts around the neck and a banded juvenile pattern (Fig. 2), recognized the eastern populations of the genus (from Turkey and Saudi Arabia to Iran) as *W. morgani*; however, Rajabizadeh (2018) considered the Iranian populations of the genus as *Walterinnesia aegyptia morgani*.




Figure 1. Occurrence records of *W. morgani* in Iran.




Figure 2. Recent historical (1981–2010) habitat suitability of *W. morgani* in Iran (warmer colors refer to high suitability level).

Walterinnesia morgani, Morgan's black cobra or black desert cobra, is a venomous snake species that inhabits the Arabian Peninsula (Kuwait and Saudi Arabia), the extreme south of Turkey, Syria, and the majority of Iraq, as well as western and southern Iran (Ilam, Kermanshah, Khuzestan, Bushehr, Fars, and Hormozgan Provinces) (Joger 1984; Gasperetti 1988; Latifi 2000; Uğurtaş et al. 2001; Sindaco et al. 2006; Safaei-Mahroo et al. 2015). In a specific concept, the Zagros Mountains display a wide biodiversity pattern, as well as the Central Iranian Plateau and the northern Persian Gulf, in one of the well-known biodiversity hotspots (Irano-Anatolian Biodiversity Hotspot) in the Palearctic Realm (Gholamifard 2011; Mittermeier et al. 2011). *Walterinnesia morgani* is one of the representative charismatic species of this region, where its shiny black color draws attention and, as most of the sightings of this species lead to its death, is very important from the aspect of conservation.

Black desert cobras display nocturnal activity patterns (Uğurtaş et al. 2001). During the hottest parts of the day, they seek refuge in burrows or beneath rocks. The seasonal activity of desert cobras may be influenced by local climate conditions (Baran et al. 2006). In some regions, for instance, they may be less active during the hottest period of summer and more active during the cooler months (Baran et al. 2021). Nonetheless, they are capable of being active even in extreme temperatures and are acclimated to harsh desert environments (Baran et al. 2021).

In spite of the known distribution of Morgan's black cobra, *W. morgani*, (IUCN status is Least Concern (LC)) in the Central and Southern Zagros Mountains, and its medical importance (Abid et al. 2020), the national literature on the species is scarce. The main reason for this is probably the difficulty of studying this mysterious and nocturnally active venomous snake. Therefore, the aim of this study was to reveal the current potential distribution pattern and speculate on the effects of climate oscillations on the potential distribution pattern of the black desert cobra in Iran.

Materials and methods

Study area and input data

This study was conducted within the borders of the Islamic Republic of Iran (25.08–39.77°N, 44.04–63.3°E; Fig. 1). A total of 16 occurrence records of *W. morgani* were obtained from the literature (Farzanpay 1989; Latifi 2000; Nilson and Rastegar-Pouyani 2007; Fathinia et al. 2010; Gholamifard and Rastegar-Pouyani 2012; Gholamifard et al. 2012) and our field observations (Suppl. material 1). In cases in which the locality information was not georeferenced, an online geographic system application (i.e., Google Earth Pro) was used to ascertain the most precise location possible. All records were georeferenced using the WGS84 coordinate system and checked with (QGIS Development Team 2023).

Ecological niche modelling (ENM)

Bioclimatic variables and one topographic layer (elevation) were downloaded from the CHELSA database (<https://chelsa-climate.org/>) at a spatial resolution of 30 arc-second raster grids (Karger et al. 2017; Brun et al. 2022). Additionally, one other topographic variable (distance to river) was obtained from the study of (Gavashelishvili et al. 2018) (Suppl. material 2). Each layer was clipped for the study area in QGIS. Pearson correlations between variables were calculated in R v4.3 (R Core Team 2023) and highly correlated variables were eliminated ($r \geq |0.8|$). Six climatic and two topographic variables were retained and used in model construction based on the Pearson correlation coefficient: mean diurnal air temperature range (Bio_2), annual range of air temperature (Bio_7), daily mean air temperatures of the driest quarter (Bio_9), precipitation amount of the driest month (Bio_14), precipitation seasonality (Bio_15), mean monthly precipitation amount of the driest quarter (Bio_17), distance to river (d_river), elevation (elev). The contributions of the variables are given in Table 1.

All used variables were employed to predict the species niche under recent (1970–2000) and future (2071–2100) climate change projections (GFDL-ESM4, IPSL-CM6A-LR, MPI-ESM1-2-HR, MRI-ESM2-0, and UKESM1-0-LL) with the lowest and the highest limits of the shared socioeconomic pathways (SSPs) from the Coupled Model Intercomparison Project Phase 6 (CMIP6) as these models have the best performance for Eurasia (Eyring et al. 2016; Sun et al. 2022) (Suppl. material 3). Using the R package *spThin*, to reduce the effects of spatial autocorrelation (Boria et al. 2014) we utilized occurrence data that were separated by more than 2 km (Aiello-Lammens et al. 2015). One of the main issues with ENM is that it may lead to model overfitting if a large number of predictor variables are used in conjunction with a small sample size (Fielding and Bell 1997). Therefore, the Akaike Information Criterion corrected for small sample sizes (AICc) was applied to the potential distribution pattern (Hurvich and Tsai 1989).

Maximum entropy modeling was utilized since it is a robust method that can be applied to presence and pseudo-absence data. Based on presence and pseudo-absence data, this algorithm can predict the presence of a species with a probability between 0 and 1 (Phillips et al. 2009). A total of 2000 background points for *W. morgani* were sampled at random throughout the survey area. The potential habitat suitability was modeled by implementing MaxEnt 3.4.1 in R using the ‘kuenm’ package (Phillips et al. 2017; Cobos et al. 2019). To create the models for this cobra species, 80% of the occurrence data was used for generating the candidate models, and the remaining 20% for independent presence as test data.

In order to optimize the model complexity for the cobra species, 31 combinations of MaxEnt’s 5 feature classes [hinge (h), threshold (t), product (p), quadratic (q), and linear (l)] along with 17 regulation multiplier values (0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 8, 10) were evaluated. This combination of options enabled us to find the best-fitting model representing our data by generating diverse candidate models (Muscarella et al. 2014; Cobos et al. 2019). Following that, the Akaike Information Criterion corrected for small sample sizes (AICc) (with the lowest values) as well as the AUC values (with the highest values) were both used to determine the best models (Hurvich and Tsai 1989). Significance tests were performed using partial ROC (Peterson et al. 2008) and predictive power with a 5% omission rate (Anderson et al. 2003). Lastly, all model inputs were transformed into binary predictions by using the minimum training presence as the threshold to distinguish suitable areas from unsuitable ones (Pearson et al. 2007; Şahin et al. 2022a).

Results

The regions anticipated to be suitable for *W. morgani* exhibited a significant area under the curve (AUC) value of 0.955 ± 0.014 . Furthermore, the recent historical and future bioclimatic predictions (2071–2100) under the SSP126 and SSP585 scenarios had the lowest values for Akaike Information Criterion corrected (AICc) and the delta AICc at 0.169 and 0, respectively. (Figs 2, 3A–J; Table 2).

Table 1. Percentage contribution of the environmental layers used in species distribution modeling of *Walterinnesia morgani*.

Bio 2 mean diurnal air temperature range	Bio 7 annual range of air temperature	Bio 9 daily mean air temperatures of the driest quarter	Bio 14 precipitation amount of the driest month	Bio 15 precipitation seasonality	Bio 17 mean monthly precipitation amount of the driest quarter	d_river distance to river	elev elevation
10	4.3	27.7	0.4	14.7	1.1	31.9	10

Table 2. Summary for selecting the best model for species distribution maps of *W. morgani* via ‘kuenm’ package.

All candidate models	Statistically significant models	MaxEnt features	AICc	wAICc	Delta AICc	AUC	Mean AUC ratio
1054	1054	linear, quadratic, product	418.125	0.169	0.00	0.955 ± 0.014	1.866

AICc: A corrected AIC score, used for a small sample size by increasing the cost for each parameter.

wAICc: The model weight is the relative likelihood for each model, divided by the total relative likelihood for all models that were considered.

Delta AICc: The difference between the model with the lowest score (the “best” model) and the AICc score for each model.

AUC: Area under the curve is a measure of the accuracy of the model.

Mean AUC ratio ≥ 1.00 , $p < 0.05$ means predictions are significantly better than a random model.




Figure 3. Future habitat suitability (2071–2100) of *W. morgani* in Iran based on **A.** GFDL-ESM4 under optimistic scenario (ssp 126); **B.** GFDL-ESM4 under pessimistic scenario (ssp 585); **C.** IPSL-CM6A-LR under optimistic scenario (ssp 126); **D.** IPSL-CM6A-LR under pessimistic scenario (ssp 585); **E.** MPI-ESM1-2- HR under optimistic scenario (ssp 126); **F.** MPI-ESM1-2- HR under pessimistic scenario (ssp 585); **G.** MRI-ESM2-0 under optimistic scenario (ssp 126); **H.** MRI-ESM2-0 under pessimistic scenario (ssp 585); **I.** UKESM1- 0-LL under optimistic scenario (ssp 126); **J.** UKESM1- 0-LL under pessimistic scenario (ssp 585) (warmer colors refer to high suitability level).

The fact that the AUC data were very close to 1 showed that the potential distribution area revealed by the locality data obtained from the distribution area of the desert cobra displayed a much better performance than a random prediction. For *W. morgani*, 1054 statistically significant models that had different regularization multipliers and feature classes were evaluated. Linear, quadratic and product (LQP) features along with a regularization multiplier of 0.8, and with the lowest delta AICc, was the best model. The most important variables were *d_river* (31.9%), *Bio_9* (27.7%) and *Bio_15* (14.7%), respectively.

According to the analysis, the potential distribution area of *W. morgani* covers a water source-dependent pattern in western Iran. Additionally, as can be seen in future climatic conditions, the potential distribution range would be narrower than under recent historical conditions. Habitat loss is expected in each future climatic scenario, but its level varies depending on the ssp level (Table 3). For instance, scenarios in ssp 126 level forecast the species range change percentile to be between -9.814 and -18.028%, however, the ssp 585 scenarios may result in a much great-

er loss of between 49.337% and 71.262% of the species' suitable habitats in the distribution range (Suppl. material 4: fig. S1A–J; Table 3). As a result of all these scenarios, it is expected that the possible distribution of *W. morgani* will contract in southwestern Iran in the future.

Discussion

Numerous biotic and abiotic factors have significant impacts on the distribution of species (Pearson et al. 2007). To understand the responses of *W. morgani* to climatic oscillations during recent history and under future possible scenarios, we employed ENM as a useful analytical instrument to assess the range of the species. Climate changes affect all aspects of biodiversity, from organisms (organismal diversity) to biomes (ecological diversity) (Gaston and Spicer 2004), and pose a significant threat to the integrity of ecosystems (Bellard et al. 2012). Additionally, it is speculated that vertebrate species will face serious problems of adaptation related to alterations in their climatic niches in the near future (Quintero and Wiens 2013). It should be noted that the distribution of reptiles can be significantly affected by direct and/or indirect anthropogenic activities, such as human population size and human-driven climatic changes (Bickford et al. 2010).

Based on these results, most of the suitable predicted areas were slightly wider than the present potential distribution of *W. morgani*. This might be the possible effect of “*d_river*” as one of the highest contributors to shape the distribution pattern via the water requirement of the species. Additionally, daily and annual temperature cycles have also contributed to the potential distribution pattern as well as the seasonal factors, especially the driest seasonal precipitation dynamics. This overall bioclimatic and topographic pattern is observed in many herpetile species in and nearby the study area (Naumov et

Table 3. Species range change (SRC) of *W. morgani* in recently suitable habitats (gain/loss) by 2081–2100 under optimistic (ssp126) and pessimistic (ssp585) scenarios in different climate-change projections.

	Percent loss	Percent gain	Species range change
GFDL-ESM4 ssp 126	18.859	1.279	-17.58
GFDL-ESM4 ssp 585	49.337	4.739	-44.598
IPSL-CM6A-LR ssp 126	19.613	1.636	-17.977
IPSL-CM6A-LR ssp 585	71.262	0.518	-70.744
MPI-ESM1-2- HR ssp 126	13.15	3.336	-9.814
MPI-ESM1-2- HR ssp 585	65.135	0.274	-64.861
MRI-ESM2-0 ssp 126	19.854	5.286	-14.568
MRI-ESM2-0 ssp 585	58.527	10.632	-47.894
UKESM1- 0-LL ssp 126	30.247	12.219	-18.028
UKESM1- 0-LL ssp 585	77.18	7.599	-69.581

al. 2020; Kurnaz and Eroğlu 2021; Şahin et al. 2022a, b; Vaissi 2022; Kurnaz 2023).

On the other hand, our results show that there will be a future decrease in suitable habitats for *W. morgani*, but the level of the decreasing trend varies depending on the SSP levels and different future scenario sets (Suppl. material 4: fig. S1A–J; Table 3). This trend is compatible with many ecological niche modeling studies that have been applied to herptiles in the study area and nearby regions (Kurnaz and Şahin 2021; Bozkurt 2022). Additionally, this contraction pattern has also been predicted for future distribution trends in many lizard species (Vaissi 2022). Nonetheless, the genus *Lacerta* and two leopard geckos exhibit the opposite trend in the Anatolian Peninsula and Iran, respectively (Hosseinian Yousefkhani and Nabizadeh 2022; Gül et al. 2023).

When compared to other large vertebrate groups, the ability of reptiles to migrate is quite restricted (Hickling et al. 2006). Moreover, our knowledge about the ecology and dispersal dynamics of *W. morgani* has received little or no attention. Based on Figs 2, 3, the current research assumed an infinite capacity for dispersal across species and made predictions about its range changes through 2100. Despite contractions and expansions, the potential distribution range of the species under each ssp126 scenario remains less changed (Fig. 3A, C, E, G, I; Table 3); however, under ssp585 scenarios, the species will be contract in southwestern Iran (Fig. 3B, D, F, H, J; Table 3). In the pessimistic scenario (ssp585) for the IPSL-CM6A-LR climate projection, in particular, it is expected that there would be 71.26% habitat loss with only 0.51% habitat gain and as a result, there would be a 70.74% negative range change (Suppl. material 4: fig. S1D; Table 3). Additional factors, such as geographic barriers, may influence the migration rate of a species (Morena-Rueda et al. 2012). In our case, distance to the river and elevation parameters, as topographic variables, make significant contributions to the species distribution. Even though the Zagros Mountains can be assessed as a remarkable geographic barrier, the main limitation factor is the “distance to the rivers” for *W. morgani*. Because it seems that the cobra tends not to occur very distant from water resources. Moreover, even though the potential distribution map displays suitable habitats in central or eastern regions of Iran, it should be noted that this could be the consequence of bioclimatic and topographic similarities in distinct areas that the species is biogeographically incapable of inhabiting. A similar pattern to ours was observed in several studies which evaluate narrower regions (Heidari 2021; Bozkurt 2022; Şahin 2022b).

On the other hand, the model algorithm that was used in this study did not take into account some parameters, such as parasitism, disease, habitat loss, and fragmentation that might affect the realistic distribution of the species (Todd et al. 2010). In the meantime, a recent study that focused on snake bite risk in Iran showed that the northern and western parts of Iran would have greater risk than the rest of the country (Yousefi et al. 2023). However, our results pointed out that southwestern Iran could

be a potentially suitable region for Morgan’s black cobra. No common potential distribution model was reported for *W. morgani*, neither country-based nor across-range, by including up-to-date topographic variables. Thus, our results suggest that *W. morgani* should be monitored in the near future, with more occurrence records collected in order to better understand its distribution and to increase awareness of human-cobra conflict as a result of anthropogenic activities. To conclude, it is desirable to predict the potential distribution of this species in Iran from the aspect of medical treatment of possible bites of the species and to prepare its antivenom.

Author contributions

A. Gholamifard and M.K. Şahin conceived the study. A. Gholamifard collected the field data and documentation. M.K. Şahin carried out the sampling design, statistical analysis and performed modeling. A. Gholamifard and M.K. Şahin contributed equally to writing the manuscript.

Acknowledgments

We thank Dr. Erica DeMilio for the proof reading and the respected anonymous reviewers and Editor Dr. Yuri Kornilev for improving the manuscript. We are grateful to Professor Aaron M. Bauer for editing the final version of this manuscript.

References

- Abid I, Jemel I, Alonazi M, Ben Bacha A (2020) A new Group II Phospholipase A2 from *Walterinnesia aegyptia* venom with antimicrobial, antifungal, and cytotoxic potential. *Processes* 8(12): e1560. <https://doi.org/10.3390/pr8121560>
- Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP (2015) SpThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography* 38(5): 541–545. <https://doi.org/10.1111/ecog.01132>
- Anderson RP, Lew D, Peterson AT (2003) Evaluating predictive models of species’ distributions: criteria for selecting optimal models. *Ecological Modelling* 162(3): 211–232. [https://doi.org/10.1016/S0304-3800\(02\)00349-6](https://doi.org/10.1016/S0304-3800(02)00349-6)
- Baran I, Kumlutaş Y, Avcı A, Ilgaz Ç (2006) Second record of *Walterinnesia aegyptia* Lataste, 1887 from Southern Anatolia. *Herpetozoa* 9(2): 87–90.
- Baran I, Avcı A, Kumlutaş Y, Olgun K, Ilgaz Ç (2021) Türkiye Amfibi ve Sürüngenleri. Palme Publishing, Ankara, 223 pp. [in Turkish]
- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F (2012) Impacts of climate change on the future of biodiversity. *Ecology Letters* 15(4): 365–377. <https://doi.org/10.1111/j.1461-0248.2011.01736.x>
- Biber MF, Voskamp A, Hof C (2023) Potential effects of future climate change on global reptile distributions and diversity. *Global Ecology and Biogeography* 15(3): 354–358. <https://doi.org/10.1111/2022.05.07.490295>

- Bickford D, Howard SD, Ng DJJ, Sheridan JA (2010) Impacts of climate change on the amphibians and reptiles of Southeast Asia. *Biodiversity and Conservation* 19: 1043–1062. <https://doi.org/10.1007/s10531-010-9782-4>
- Bobrowski M, Bechtel B, Böhner J, Oldeland J, Weidinger J, Schickhoff U (2018) Application of thermal and phenological land surface parameters for improving ecological niche models of *Betula utilis* in the Himalayan region. *Remote Sensing* 10(6): e814. <https://doi.org/10.3390/rs10060814>
- Boria RA, Olson LE, Goodman SM, Anderson RP (2014) Spatial filtering to reduce sampling bias can improve the performance of ecological niche models. *Ecological Modelling* 275: 73–77. <https://doi.org/10.1016/j.ecolmodel.2013.12.012>
- Bozkurt E (2022) Ecological niche modelling of the genus *Anatololacerta* under past, historical and future bioclimatic conditions. *Folia Biologica* 70: 45–53. https://doi.org/10.3409/fb_70-2.06
- Brun P, Zimmermann NE, Hari C, Pellissier L, Karger DN (2022) Global climate-related predictors at kilometer resolution for the past and future. *Earth System Science Data* 14(12): 5573–5603. <https://doi.org/10.5194/essd-14-5573-2022>
- Cobos ME, Peterson AT, Barve N, Osorio-Olvera L (2019) kuenm: An R package for detailed development of ecological niche models using Maxent. *PeerJ* 7: e6281. <https://doi.org/10.7717/peerj.6281>
- Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ, Taylor KE (2016) Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development* 9(5): 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>
- Farzanpay R (1989) Ophiology. University Publication Center, Tehran, 248 pp. [In Persian]
- Fathinia B, Rastegar-Pouyani N, Darvishnia H, Rajabizadeh M (2010) The snake fauna of Ilam Province, southwestern Iran. *Iranian Journal of Animal Biosystematics* 6(1): 9–23. <https://doi.org/10.22067/ijab.v6i1.9153>
- Fielding AH, Bell JF (1997) A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environmental Conservation* 24(1): 38–49. <https://doi.org/10.1017/S0376892997000088>
- Gasperetti J (1988) Snakes of Arabia. *Fauna of Saudi Arabia* 9: 340–343.
- Gaston KJ, Spicer JJ (2004) *Biodiversity: An introduction* (2nd edn.). Blackwell Publishing, UK, 191 pp.
- Gavashelishvili A, Yuriy A, Yarovenko YA, Babayev EA, Mikeladze G, Gurielidze Z, Dekanoidze D, Kerdikoshvili N, Ninua L, Paposhvili N (2018) Modeling the distribution and abundance of Eastern Tur (*Capra cylindricornis*) in the Caucasus. *Journal of Mammalogy* 99(4): 885–897. <https://doi.org/10.1093/jmammal/gyy056>
- Gholamifard A (2011) Endemism in the reptile fauna of Iran. *Iranian Journal of Animal Biosystematics* 7(1): 13–29. <https://doi.org/10.22067/ijab.v7i1.25218>
- Gholamifard A, Rastegar-Pouyani N (2012) Systematics and distribution of *Walterinnesia morgani* (Mocquard, 1905) (Elapidae) in Iran. 5th Asian Herpetological Conference, June 2, Chengdu, China.
- Gholamifard A, Rastegar-Pouyani N, Esmaili HR (2012) Annotated checklist of reptiles of Fars Province, Southern Iran. *Iranian Journal of Animal Biosystematics* 8(2): 155–167. <https://doi.org/10.22067/ijab.v8i2.25584>
- Gibbons WJ, Scott DE, Ryan TJ, Buhlmann KA, Tuberville TD, Metts BS, Greene JL, Mills T, Leiden Y, Poppy S (2000) The global decline of reptiles, Déjà Vu Amphibians: reptile species are declining on a global scale. six significant threats to reptile populations are habitat loss and degradation, introduced invasive species, environmental pollution, disease, unsustainable use, and global climate change. *BioScience* 50(8): 653–666. [https://doi.org/10.1641/0006-3568\(2000\)050\[0653:TGDORD\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2000)050[0653:TGDORD]2.0.CO;2)
- Guedes TB, Sawaya RJ, Zizka A, Laffan S, Faurby S, Pyron RA, Bernils RS, Jansen M, Passos P, Prudente ALC (2018) Patterns, biases and prospects in the distribution and diversity of Neotropical snakes. *Global Ecology and Biogeography* 27(1): 14–21. <https://doi.org/10.1111/geb.12679>
- Gül S, Kuşlutaş Y, İlğaz Ç, Candan K (2023) Climatic envelopes of the genus *Lacerta* Linnaeus, 1758 in Türkiye: an application of ecological niche modeling. *Environmental Science and Pollution Research* 30: 56382–56397. <https://doi.org/10.1007/s11356-023-26351-4>
- Heidari N (2021) Ecological niche differentiation between *Lacerta media* and *Lacerta strigata* (Sauria: Lacertidae) in Iran. *Biologia* 76: 901–905. <https://doi.org/10.2478/s11756-020-00601-z>
- Hickling R, Roy DB, Hill JK, Fox R, Thomas CD (2006) The distributions of a wide range of taxonomic groups are expanding polewards. *Global Change Biology* 12(3): 450–455. <https://doi.org/10.1111/j.1365-2486.2006.01116.x>
- Hosseiniyan Yousefkhani SS (2019) Review to the last ecological niche modelling studies on Iranian herpetofauna and their importance for species conservation. *Journal of Biological Studies* 1(4): 165–167.
- Hosseiniyan Yousefkhani SS, Nabizadeh H (2022) Response to the future climate change effect of two Leopard geckos' distribution (Sauria: Eublepharidae) in Iran. *Journal of Asia-Pacific Biodiversity* 15(3): 354–358. <https://doi.org/10.1016/j.japb.2022.04.004>
- Hosseinzadeh MS, Ghezellou P, Kazemi SM (2017) Predicting the potential distribution of the endemic snake *Spalerosophis microlepis* (Serpentes: Colubridae), in the Zagros Mountains, Western Iran. *Salamandra* 53(2): 294–298.
- Hurvich CM, Tsai CL (1989) Regression and time series model selection in small samples. *Biometrika* 76(2): 297–307. <https://doi.org/10.1093/biomet/76.2.297>
- Joger U (1984) The venomous snakes of the Near and Middle East. Wiesbaden, L. Reichert, 115 pp.
- Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler M (2017) Climatologies at high resolution for the earth's land surface areas. *Scientific Data* 4(1): 1–20. <https://doi.org/10.1038/sdata.2017.122>
- Kurnaz M, Eroğlu Aİ (2021) Habitat preferences of *Testudo graeca*, Linnaeus 1758 (Testudines; Testudinidae) with a new locality record from eastern Black Sea region of Turkey. *Scientific Reports in Life Sciences* 2(2): 30–38. <https://doi.org/10.22034/srls.2021.527238.1015>
- Kurnaz M, Şahin MK (2021) A contribution to the biogeography and taxonomy of two Anatolian mountain brooknewts, *Neurergus barani* and *N. strauchii* (Amphibia: Salamandridae) using ecological niche modeling. *Turkish Journal of Zoology* 45(1): 54–64. <https://doi.org/10.3906/zoo-2007-37>
- Kurnaz M (2023) Geographic distribution of two *Montivipera* taxa using ecological niche modeling. *Journal of Bioscience* 48: 1–7. <https://doi.org/10.1007/s12038-023-00333-7>
- Latifi M (1985) Snakes of Iran. Department of the Environment, Tehran, 221 pp. [In Persian]
- Latifi M (1991) Snakes of Iran. Society for the Study of Amphibians and Reptiles, Oxford, Ohio, 159 pp. <https://doi.org/10.2307/1446120>

- Latifi M (2000) Snakes of Iran (3rd edn.). Department of the Environment, Tehran, 478 pp. [In Persian]
- Leviton AE, Anderson SC, Adler K, Minton SA (1992) Hand-Book to Middle East Amphibians and Reptiles. Society for the Study of Amphibians and Reptiles, Oxford, Ohio, 252 pp.
- Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C (2011) Global Biodiversity Conservation: The Critical Role of Hotspots. Biodiversity hotspots. Springer, 3–22. https://doi.org/10.1007/978-3-642-20992-5_1
- Mocquard MF (1905) Diagnoses de quelques espèces nouvelles de reptiles. Bulletin du Muséum d'Histoire Naturelle 11(2): 76–78.
- Moradi SH, Rastegar-Pouyani E, Hosseini Yousefkhani SS, Zargan J (2021) Evaluation ecological niche between *Platyceps rhodorachis* and *P. karelini* (Serpentes: Colubridae) in Iran. Iranian Journal of Animal Biosystematics 17(2): 147–155. <https://doi.org/10.22067/ijab.2022.69956.1014>
- Morena-Rueda G, Pleguezuelos JM, Pizarro M, Montori A (2012) Northward shifts of the distributions of Spanish reptiles in association with climate change. Conservation Biology 26(2): 278–283. <https://doi.org/10.1111/j.1523-1739.2011.01793.x>
- Muscarella R, Galante PJ, Soley Guardia M, Boria RA, Kass JM, Uriarte M, Anderson RP (2014) ENM Eval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models. Methods in Ecology and Evolution 5(11): 1198–1205. <https://doi.org/10.1111/2041-210X.12261>
- Naumov BY, Popgeorgiev GS, Kornilev YV, Plachiyski DG, Stojanov AJ, Tzankov ND (2020) Distribution and ecology of the Alpine newt *Ichthyosaura alpestris* (Laurenti, 1768)(Amphibia: Salamandridae) in Bulgaria. Acta Zoologica Bulgarica 72: 83–102.
- Nilson G, Rastegar-Pouyani N (2007) *Walterinnesia aegyptia* Lataste, 1887 (Ophidia: Elapidae) and the status of *Naja morgani* Mocquard 1905. Russian Journal of Herpetology 14(1): 7–14.
- Pearson RG, Raxworthy CJ, Nakamura M, Peterson AT (2007) Predicting species distributions from small numbers of occurrence records: A test case using cryptic geckos in Madagascar. Journal of Biogeography 34(1): 102–117. <https://doi.org/10.1111/j.1365-2699.2006.01594.x>
- Peterson AT, Papeş M, Soberón J (2008) Rethinking receiver operating characteristic analysis applications in ecological niche modeling. Ecological Modelling 213(1): 63–72. <https://doi.org/10.1016/j.ecolmodel.2007.11.008>
- Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. Ecological Applications 19(1): 181–197. <https://doi.org/10.1890/07-2153.1>
- Phillips SJ, Dudík M, Schapire RE (2017) Maxent software for modeling species niches and distributions (version 3.4.1). Biodiversity Informatics. http://biodiversityinformatics.amnh.org/open_source/maxent/
- QGIS Development Team (2023) QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>
- Quintero I, Wiens JJ (2013) Rates of projected climate change dramatically exceed past rates of climatic niche evolution among vertebrate species. Ecology Letters 16(8): 1095–1103. <https://doi.org/10.1111/ele.12144>
- R Core Team (2023) R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna. <https://www.R-project.org>
- Rajabizadeh M (2018) Snakes of Iran. Iranshenasi Publishing, Tehran, 496 pp. [In Persian]
- Safaei-Mahroo B, Ghaffari H, Fahimi H, Broomand S, Yazdani M, Najafi-Majid E, Hosseini Yousefkhani SS, Rezazadeh E, Hosseinzadeh MS, Nasrabadi R, Rajabizadeh M, Mashayekhi M, Mote-sharei M, Naderi A, Kazemi SM (2015) The herpetofauna of Iran: checklist of taxonomy, distribution and conservation status. Asian Herpetological Research 6(4): 257–290. <https://doi.org/10.16373/j.cnki.ahr.140062>
- Saha A, McRae L, Dodd Jr CK, Gadsden H, Hare KM, Lukoschek V, Böhm M (2018) Tracking global population trends: population time-series data and a living planet index for reptiles. Journal of Herpetology 52(3): 259–268. <https://doi.org/10.1670/17-076>
- Sun C, Zhu L, Liu Y, Wei T, Guo Z (2022) CMIP6 model simulation of concurrent continental warming holes in Eurasia and North America since 1990 and their relation to the Indo-Pacific SST warming. Global and Planetary Change 213: e103824. <https://doi.org/10.1016/j.gloplacha.2022.103824>
- Şahin MK, Candan K, Karakasi D, Lymberakis P, Poulakakis N, Kumlutaş Y, Yildirim E, Ilgaz Ç (2022a) Ecological niche differentiation in the Anatolian rock lizards (Genus: *Anatololacerta*) (Reptilia: Lacertidae) of the Anatolian Peninsula and Aegean Islands. Acta Herpetologica 17(2): 165–175. https://doi.org/10.36253/a_h-13089
- Şahin MK, Kumlutaş Y, Yanchukov A, Çetintaş O, Candan K, Ilgaz Ç, Ayaş Z (2022b) The Quaternary range dynamics of the dwarf lizard, *Parvilacerta parva* (Boulenger, 1887) (Squamata, Lacertidae) in the Anatolian Peninsula. Journal of Wildlife and Biodiversity 6(1): 79–86. <https://doi.org/10.22120/jwb.2021.540551.1259>
- Sindaco R, Serra G, Menegon M (2006) New data on the Syrian Herpetofauna, with a newly-recorded species of snake. Zoology in the Middle East 37(1): 29–38. <https://doi.org/10.1080/09397140.2006.10638146>
- Todd BD, Willson JD, Gibbons JW (2010) The global status of reptiles and causes of their decline. In: Sparling DW, Linder G, Bishop CA (Eds) Ecotoxicology of Amphibians and Reptiles (2nd Edn.). SETAC Press, Pensacola, 47–67. <https://doi.org/10.1201/EBK1420064162-c3>
- Uetz P, Freed P, Aguilar R, Reyes F, Hošek J [Eds] (2023) The Reptile Database. <http://www.reptile-database.org> [accessed (9 May 2023)]
- Uğurtaş IH, Papenfuss TJ, Orlov NL (2001) New record of *Walterinnesia aegyptia* Lataste, 1887 (Ophidia: Elapidae: Bungarinae) in Turkey. Russian Journal of Herpetology 8(3): 239–245.
- Vaissi S (2021a) Historic range dynamics in Kaiser's Mountain Newt (*Neurergus kaiseri*): insights from phylogeographic analyses and species distribution modeling. Ecology and Evolution 11(12): 7622–7633. <https://doi.org/10.1002/ece3.7595>
- Vaissi S (2021b) Potential changes in the distributions of Near Eastern Fire Salamander (*Salamandra infraimmaculata*) in response to historical, recent and future climate change in the Near and Middle East: implication for conservation and management. Global Ecology and Conservation 29: e01730. <https://doi.org/10.1016/j.gecco.2021.e01730>
- Vaissi S (2022) Response of Iranian lizards to future climate change by poleward expansion, southern contraction, and elevation shifts. Scientific Reports 12(1): 1–21. <https://doi.org/10.1038/s41598-022-06330-4>
- Yousefi M, Hosseini Yousefkhani SS, Grünig M, Kafash A, Rajabizadeh M, Rastegar-Pouyani E (2023) Identifying high snakebite risk area under climate change for community education and antivenom distribution. Scientific Reports 13(1): e8191. <https://doi.org/10.1038/s41598-023-35314-1>

Supplementary material 1

Species occurrence records of *W. morgani* in Iran from literature and our field trips

Authors: Ali Gholamifard, Mehmet Kürşat Şahin

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e107947.suppl1>

Supplementary material 2

Variables used for modeling the distribution of black desert cobra (*Walterinnesia morgani*)

Authors: Ali Gholamifard, Mehmet Kürşat Şahin

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e107947.suppl2>

Supplementary material 3

Climate forcing data and source for the CHELSA CMIP6 ISIMIP3 data

Authors: Ali Gholamifard, Mehmet Kürşat Şahin

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e107947.suppl3>

Supplementary material 4

Species range change (SRC) of *W. morgani* in recently suitable habitats (gain/loss) by 2081–2100 under optimistic (ssp126) and pessimistic (ssp585) scenarios

Authors: Ali Gholamifard, Mehmet Kürşat Şahin

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/herpetozoa.36.e107947.suppl4>