



# Rediscovery of *Serangium montazerii* Fürsch in Georgia and updated list of the Coccinellidae of Georgia

Alain Migeon<sup>1</sup>, Tea Arabuli<sup>2,3</sup>

<sup>1</sup> CBGP, INRAE, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France

<sup>2</sup> Institute of Entomology, Agricultural University of Georgia, University Campus at Digomi David Aghmashenebeli Alley, 13<sup>th</sup> km, Tbilisi, Georgia

<sup>3</sup> Institute of Zoology, Ilia State University, Kakutsa Cholokashvili Ave 3/5, Tbilisi, Georgia

<http://zoobank.org/0E5BCF84-85E6-48D6-88DD-2CD7DE63956F>

Corresponding author: Alain Migeon ([alain.migeon@inrae.fr](mailto:alain.migeon@inrae.fr))

Academic editor: Levan Mumladze ♦ Received: 20 November 2020 ♦ Accepted: 8 July 2021 ♦ Published: 12 January 2022

## Abstract

This article provides an updated list of the Coccinellidae of Georgia, recording 84 species. *Serangium montazerii* Fürsch nec *S. parcesetosum* Sicard is also rehabilitated as a species actually present in the country.

## Key words

beetles, Caucasus, checklist, faunistic entomology

## Introduction

Merkviladze and Kvavadze (2002) provided a list that names 84 species of Coccinellidae found in Georgia. Among the species listed, *Rhyzobius lophanthae* (Blaisdell, 1892), *Novius cardinalis* (Mulsant, 1850), *Harmonia conformis* (Boisduval, 1835), and *Harmonia axyridis* (Pallas, 1773) are exotic species introduced for biological control. In the close vicinity of Georgia, Orlova-Bienkowskaja and Bieńkowski (2017) also recorded *Cryptolaemus montrouzieri* Mulsant, 1853, and *Serangium montazerii* Fürsch, 1995, which are not listed in the Georgian fauna, despite the introduction of *Serangium parcesetosum* Sicard, 1929 in the country as predator of *Dialeurodes citri* (Ashmead, 1885) during the seventies (Timofeyeva and Hoang Duc Nhuan 1978).

Here we propose an updated list of the Coccinellidae found in Georgia, taking into account the actual phylogeny and nomenclature. We also report the rediscovery of *Serangium montazerii* Fürsch, 1995 nec *Serangium parcesetosum* Sicard, 1929 in an orange grove near Batumi, and

make address nomenclatural confusion about the species in Georgia.

## Material and methods

### Site description

The sampled site is located at Chakvis Tskali, Kobuleti District, Adjara (41.71843°N, 41.73841°E) in a familial orange grove and garden. Further orange trees located 6 km eastward along the Chakvistskali River were also sampled. Observations were conducted on 25-VI-2019.

The climate of the region is characteristic of the Batumi area and the Black Sea Georgian coast, with high precipitations (avg. 2515 mm/year), mild winters (avg. 7 °C in January) and mild summers (avg. 23 °C in July). The region has warm temperate, almost subtropical, climate conditions (Bohn et al. 2000–2003). The vegetation is represented by hygrophilous thermophytic mixed deciduous broadleaved

forest (Bohn et al. 2000–2003). The presence of many citrus orchards, with their specific pests, have made Batumi a focus of biocontrol attempts, with release of exotic Coccinellidae during the Soviet period (Timofeyeva and Hoang Duc Nhuan 1978).

### Sampling method

Beating trays were used to collect ladybird beetles by beating and shaking the branches of orange trees. The ladybird beetles were then collected with an aspirator and subsequently identified to species level in the laboratory using available literature (Gourreau 1974, Iablokoff-Knזורian 1978, Poorani 1998, Bieńkowski 2018, 2020).

### Systematics

The list of the species recorded from Georgia is established following Kovář (2007), Seago et al. (2011), Escalona and Slipinski (2012), Che et al. (2017, 2021) and Zhang et al. (2018) for supra-generic levels; Tomaszewska and Szawaryn (2016): genus *Chnootriba*; Pang et al. (2020): genus *Novius*; Szawaryn and Tomaszewska (2020): genus *Rhyzobius*; Nattier et al. (2021) and Tomaszewska et al. (2021): genera *Anisocalvia* and *Hippodamia*.

## Results and discussion

### *Serangium montazerii* on *Citrus* sp.

This non-native species intentionally introduced in Georgia (Adjara) was initially identified as *Serangium parcesetosum* Sicard, 1929 (Timofeyeva and Hoang Duc Nhuan, 1978). These authors also reassigned it to its original genus after the transfer to the genus *Catana* by Chapin (1940). They were followed by latter authors and by Escalona and Slipinski (2012).

However, Fürsch (1995) described a new species, *Serangium montazerii* Fürsch, 1995 from Iranian specimens (Mazandran Province). Subsequently Duverger (1998) as-

signed all the introduced populations of *Serangium* in Western Europe, which were resulting from the first one introduced by Timofeyeva and Hoang Duc Nhuan (1978), to this latter species. However, after examination of specimens from Corsica, Coutanceau (2006), and Coutanceau and Malausa (2014) assigned the French introduced populations to *S. parcesetosum*. Their works, published in French in an amateurs' journal, remained largely unknown. Escalona and Slipinski (2012) in their revision of the Microweiseinae subfamily did not examine any members of the two taxa. The specimens introduced in France (continental and Corsica, Malausa et al. (1988)), and in Turkey (Yigit and Canhilal 2005)) came from this Georgian source (Timofeyeva and Hoang Duc Nhuan 1978). Likewise, neither Fürsch (1995), nor Duverger (1998), nor Coutanceau (2006) examined or compared the two taxa. However, Poorani (1998) noted that the two species can be distinguished by the shape of the right paramere and by the inter-ocular distance. Bieńkowski (2020) also underlined the same diagnostic characters and revised illustrations of the genitalia of both taxa. Moreover, the population introduced in Georgia in the 1970's (Timofeyeva and Hoang Duc Nhuan 1978) came from North India (Raniket, Uttarakhand) where *S. montazerii* occurs, while *S. parcesetosum* is distributed into Central and Southern India (Poorani 1998).

We found one specimen of *S. montazerii* on orange trees (Figure 1). It was female and we could not perform a determination using genitalia. However, the shape of the head with an inter-ocular distance equal to twice the size of the eyes (Figure 1B) allows us to assign this specimen to *S. montazerii*. The location is close to Batumi, and corresponds to the area of first historical introduction of the species in Georgia, the country of first use of the species for biological control of *D. citri* (Timofeyeva and Hoang Duc Nhuan 1978). According to Yigit and Canhilal (2005), the species was still present in Georgia in 1990. It has probably been present in the country since 1975, but was not listed by Merkviladze and Kvavadze (2002). The species is also present northward in the vicinity of Sochi (Orlova-Bienkowskaja and Bieńkowski 2017; Bieńkowski



**Figure 1.** *Serangium montazerii* Fürsch, 1995, collected at Chakvis Tskali, 25-VI-2019 on an orange tree. **A:** dorsal view; **B** front view of the head, showing the inter-ocular distance close to 2× eye diameter. Pictures from CBGP – Continental Arthropod Collection (2021).

2018, 2020; Bieńkowski and Orlova-Bienkowskaja 2020) in Russia.

## Other species found on *Citrus* sp.

Along with *S. montazerii*, we found many *Harmonia axyridis* (Pallas, 1773) both *succinea* and *axyridis* forms, *Novius cardinalis* (Mulsant, 1850), *Rhyzobius lophanthae* (Blaisdell, 1892). All of these species are introduced. Among the native species, we found *Hyperaspis campestris* Herbst, 1793, *Scymnus haemorrhoidalis* Herbst, 1797, *Scymnus subvillosus* (Goeze, 1777), *Scymnus interruptus* (Goeze, 1777), *Scymnus rubromaculatus* (Goeze, 1777) and *Nephus quadrimaculatus* (Herbst, 1783).

In addition, we report one observation of *Hippodamia undecimnotata* (Schneider, 1792) in Chalaubani (Kakheti, Gurjaani, 41.6291°N, 45.7946°E) 21-VI-2019 on Carduaceae, a new district and a new province not recorded in Merkviladze and Kvavadze (2002).

## List of the species of Georgia

When a taxon's nomenclature has been changed since the publication of Merkviladze and Kvavadze (2002), the original indication is reported. Many species were not listed by Kovář (2007) in the Catalogue of Palearctic Coleoptera. We add some indications for these species. The list is in alphabetical order. The supra generic classification follows Che et al. (2021).

### Family Coccinellidae Latreille, 1807

#### Subfamily Coccinellinae Latreille, 1807

##### Tribe Chilochorini Mulsant, 1846

- Six species are recorded from Georgia.

*Chilocorus bipustulatus* (Linnaeus, 1758)

*Chilocorus renipustulatus* (Scriba, 1790)

*Exochomus octosignatus* (Gebler, 1830)

*Brumus octosignatus* Gebler, 1830

- This species is not recorded in Georgia by Kovář (2007).

*Exochomus quadripustulatus* (Linnaeus, 1758)

*Exochomus undulatus* Weise, 1878

*Parexochomus nigromaculatus* (Goeze, 1777)

*Exochomus nigromaculatus* Goeze, 1777

##### Tribe Coccidulini Mulsant, 1846

- Three species can be considered as present in Georgia.

*Coccidula rufa* (Herbst, 1783)

*Coccidula scutellata* (Herbst, 1783)

*Rhyzobius lophanthae* (Blaisdell, 1892)

*Lindorus lophanthae* (Blaisdell, 1892)

- *R. lophanthae* is an introduced species established in a large number of countries in Europe and around the World (Roy and Migeon, 2010, Rondoni et al., 2020).

##### Tribe Coccinellini Latreille, 1807

- Thirty-six species are recorder from Georgia.

*Adalia bipunctata* (Linnaeus, 1758)

*Adalia decempunctata* (Linnaeus, 1758)

*Anatis ocellata* (Linnaeus, 1758)

- This species is not recorded in Georgia by Kovář (2007).

*Anisocalvia quatuordecimguttata* (Linnaeus, 1758)

*Calvia quatuordecimguttata* Linnaeus, 1758

*Calvia rosti* Weise, 1871

- Following Kovář (2007) *C. quatuordecimguttata* and *C. rosti* are synonyms.

*Anisocalvia quindecimguttata* (Fabricius, 1777)

- This species is not recorded in Georgia by Kovář (2007).

*Anisosticta novemdecimpunctata* (Linnaeus, 1758)

- This species is not recorded in Georgia by Kovář (2007).

*Aphidecta obliterated* (Linnaeus, 1758)

*Bulaea lichatschovii* (Hummel, 1827)

*Calvia decemguttata* (Linnaeus, 1767)

*Coccinella hieroglyphica* Linnaeus, 1758

- This species is not recorded in Georgia by Kovář (2007).

*Coccinella magnifica* Redtenbacher, 1847

*Coccinella divaricata* Olivier, 1808

- Following Kovář (2007) *C. magnifica* and *C. divaricata* are synonyms.

*Coccinella quinquepunctata* Linnaeus, 1758

*Coccinella septempunctata* Linnaeus, 1758

*Coccinula quatuordecimpustulata* (Linnaeus, 1758)

*Coccinula sinuatomarginata* (Faldermann, 1837)

*Halysia sedecimguttata* (Linnaeus, 1758)

*Harmonia axyridis* (Pallas, 1773)

- This species is not recorded in Georgia by Kovář (2007). *Harmonia axyridis* has been introduced in 1927 (Iablokoff-Khnzorian 1982), but wasn't observed until 2002 (Merkviladze and Kvavadze 2002) in Eastern Georgia (Lagodekhy Reserve). Genetic studies would be necessary to assess the origin of the actual population(s), perhaps resulting of spreading from the admixed Western-Europe population (Lombaert et al. 2011).

*Harmonia conformis* (Boisduval, 1835)

- This species is not recorded in Georgia by Kovář (2007). *Harmonia conformis* is an introduced species (Iablokoff-Khnzorian 1982, 1983), also established in France (Coutanceau 2009) and Egypt (Iablokoff-Khnzorian 1982) against the psyllid *Acizzia uncatoides*. Its establishment has been confirmed in 2002 (Merkviladze and Kvavadze 2002).

*Harmonia quadripunctata* (Pontippidan, 1763)

*Hippodamia apicalis* (Weise, 1879)

*Semiadalia apicalis* (Weise, 1879)

*Hippodamia notata* (Laicharting, 1781)

*Semiadalia notata* Laicharting, 1781

- This species is not recorded in Georgia by Kovář (2007).

*Hippodamia schneideri* (Weise, 1878)

*Semiadalia schneideri* (Weise, 1878)

*Hippodamia tredecimpunctata* (Linnaeus, 1758)

- This species is not recorded in Georgia by Kovář (2007).

*Hippodamia undecimnotata* (Schneider, 1792)

*Semiadalia undecimnotata* Schneider, 1792

*Hippodamia variegata* (Goeze, 1777)

*Myrrha octodecimguttata* (Linnaeus, 1758)

*Oenopia conglobata* (Linnaeus, 1758)

*Synharmonia conglobata* Linnaeus, 1758

*Oenopia impustulata* (Linnaeus, 1758)

*Synharmonia impustulata* Linnaeus, 1758



*Oenopia lyncea* (Olivier, 1808)

*Synharmonia lyncea* Olivier, 1808

- This species is not recorded in Georgia by Kovář (2007).

*Oenopia oncina* (Olivier, 1808)

*Propylea quatuordecimpunctata* (Linnaeus, 1758)

*Psyllobora vigintiduopunctata* (Linnaeus, 1758)

*Thea vigintiduopunctata* Linnaeus, 1758

*Sospita oblongoguttata* (Linnaeus, 1758)

*Neomysia oblongoguttata* Linnaeus, 1758

- This species is not recorded in Georgia by Kovář (2007).

*Sospita vigintiguttata* (Linnaeus, 1758)

- This species is not recorded in Georgia by Kovář (2007).

*Tytthaspis sedecimpunctata* (Linnaeus, 1758)

- This species is not recorded in Georgia by Kovář (2007).

*Vibidia duodecimguttata* (Poda, 1761)

### Tribe Diomini Gordon, 1999

- One species is recorded from Georgia.

*Diomus rubidus* (Motschulsky, 1837)

*Scymnus (Diomus) rubidus* Motschulsky, 1837

- This species is not recorded in Georgia by Kovář (2007). However, it is also present in Daghestan (Bieńkowski 2018) from where it was described by Motchulsky (1837).

### Tribe Epilachnini Mulsant, 1846

- Three species are recorded from Georgia.

*Chnootriba argus* (Geoffroy, 1785)

*Epilachna argus* (Geoffroy, 1785)

- This species is not recorded in Georgia by Kovář (2007).

*Chnootriba elaterii* (Rossi, 1794)

*Epilachna chrysomelina* Fabricius, 1775

- This species is not recorded in Georgia by Kovář (2007).

*Subcoccinella vigintiquatuorpunctata* (Linnaeus, 1758)

### Tribe Hyperaspini Mulsant, 1846

- Five species are recorded from Georgia.

*Hyperaspis campestris* (Herbst, 1793)

*Hyperaspis reppensis* (Herbst, 1793)

*Hyperaspis femorata* (Motchulsky, 1783)

- This species is not recorded in Georgia by Kovář (2007) but in Azerbaijan and Armenia.

*Hyperaspis transversoguttata* Weise, 1878

- This species is not recorded in Georgia by Kovář (2007) but in Armenia.

*Hyperaspis erythrocephala* (Fabricius, 1787)

*Oxynychus erythrocephala* (Fabricius, 1787)

### Tribe Noviini Mulsant, 1846

- One species is recorded from Georgia.

*Novius cardinalis* (Mulsant, 1850)

*Rodolia cardinalis* (Mulsant, 1850)

- This species is not recorded in Georgia by Kovář (2007). *Novius cardinalis* is an introduced species now present in many countries and considered as the most

effective among the ladybird beetles released (Rondoni et al. 2020), consuming the scale insect *Icerya purchasi*.

### Tribe Platynaspini Mulsant, 1846

- One species is recorded from Georgia.

*Platynaspis luteorubra* (Goeze, 1777)

### Tribe Scymnini Mulsant, 1846

- Twenty species can be considered as present in Georgia.

*Clitostethus arcuatus* (Rossi, 1794)

*Nephus (Bipunctatus) bipunctatus* (Kugelann, 1794)

*Scymnus (Nephus) bipunctatus* Kugelann, 1794

- This species is not recorded in Georgia by Kovář (2007).

*Nephus (Nephus) quadrimaculatus* (Herbst, 1783)

*Scymnus (Nephus) quadrimaculatus* Herbst, 1783

- This species is not recorded in Georgia by Kovář (2007).

*Nephus (Nephus) redtenbacheri* Mulsant, 1846

*Scymnus (Nephus) redtenbacheri* Mulsant, 1846

- This species is not recorded in Georgia by Kovář (2007).

*Scymniscus biflammulatus* (Motchulsky, 1837)

- This species is not recorded by Merkviladze and Kva-vadze (2002). However, it has been described from Georgia, with no more indication of location. Its status needs to be investigated.

*Scymniscus biguttatus* (Mulsant, 1850)

*Scymnus (Scymnus) biguttatus* Mulsant, 1850

- This species is not recorded in Georgia by Kovář (2007). It was first reported in the original description of *Scymnus bipustulatus* Motchulsky, 1837 from Chirvan, now in Azerbaijan. Schneider and Leder (1878) report it under the name of *S. bipustulatus* Motchulsky, from Borjomi (Samtskhe-Javakheti) and Savenko (1953) under the name of *S. (Sidis) biguttatus* Mulsant from Tbilisi.

*Scymnus (Neopullus) haemorrhoidalis* Herbst, 1797

*Scymnus (Neopullus) limbatus* Stephens, 1832

*Scymnus (Pullus) testaceus* (Motschulsky, 1837)

- This species is not recorded in Georgia by Kovář (2007). *Scymnus testaceus* has been described from The Caucasus Baths, with no more indication.

*Scymnus (Pullus) argutus* Mulsant, 1850

*Scymnus (Pullus) auritus* Westman in Thunberg, 1795

*Scymnus (Pullus) ferrugatus* (Moll, 1785)

- This species is not recorded in Georgia by Kovář (2007).

*Scymnus (Pullus) fraxini* Mulsant, 1850

*Scymnus (Pullus) subvillosus* (Goeze, 1777)

*Scymnus (Scymnus) apetzi* Mulsant, 1846

*Scymnus (Scymnus) frontalis* (Fabricius, 1787)

*Scymnus (Scymnus) interruptus* (Goeze, 1777)

*Scymnus (Scymnus) magnomaculatus* Fürsch, 1958

*Scymnus (Scymnus) quadriguttatus* Capra, 1921

- This species is not recorded in Georgia by Kovář (2007).

*Scymnus (Scymnus) marginalis* (Rossi, 1794)

*Scymnus (Scymnus) nigrinus* Kugelann, 1794

- This species is not recorded in Georgia by Kovář (2007).

*Scymnus (Scymnus) rubromaculatus* (Goeze, 1777)

- This species is not recorded in Georgia by Kovář (2007).

#### Tribe Stethorini Dobzhansky, 1924

- Two species recorded from Georgia.

*Stethorus gilvifrons* (Mulsant, 1850)

*Scymnus (Scymnus) gilvifrons* Mulsant, 1850

- This species is not recorded in Georgia by Kovář (2007).

*Stethorus pusillus* (Herbst, 1797)

*Stethorus punctillum* Weise, 1891

#### Tribe Sticholotidini Mulsant, 1846

- Two species are recorded from Georgia.

*Pharoscygnus armenus* Iablokoff-Khnzorian, 1970

- This species is not recorded in Georgia by Kovář (2007).

*Pharoscygnus smirnovi* Dobzhanskiy, 1927

#### Tribe Tetrabrachini Kapur, 1948

- Three species are recorded from Georgia.

*Tetrabrachys caucasicus* (Weise, 1878)

*Lithophilus caucasicus* Weise, 1878

*Tetrabrachys connatus* (Creutzer in Panzer, 1796)

*Lithophilus connatus* Panzer, 1796

- This species is only recorded by Kovář (2007) in central Europe, ranging from Austria to Ukraine and from Montenegro to Bulgaria. The Georgian records were reported by Scheinder and Leder (1878) in Lailasch (Racha-Leckhumi and Kveno-Svaneti), by Savenko (1953) in Kutaisi (Imereti). Iablokoff-Khnzorian (1983) only cites the Caucasus region. Specific investigations are necessary to confirm/infirm its presence in Georgia.

*Tetrabrachys weisei* Reitter, 1880

*Lithophilus weisei* Reitter, 1880

#### Subfamily Microweiseinae Leng, 1920

##### Tribe Serangiini Pope, 1962

- Only one (introduced) species is recorded from Georgia.

*Serangium montazerii*, Fürsch, 1995

## Conclusion

The Georgian fauna of Coccinellidae is containing 84 species. Two species (*S. biflammulatus* and *S. montazerii*) have been added to the Merkviladze and Kvavadze (2002) list, but one species is dubious for the country (*T. conatus*); a further two species are now considered as synonyms. Thirty species were not listed in the Catalogue of Palearctic Coleoptera (Kovář 2007).

## Potential species to be sought

Among the biocontrol agents introduced into Georgia, *Cryptolaemus montrouzieri* Mulsant, 1853 is an important predator of mealybugs. The species is reported from Sochi (Orlova-Bienkowskaja and Bienkowski 2017) in high numbers (Orlova-Bienkowskaja et al. 2018) and could be still present in Georgia. Eight other species, released between 1935 and 1990, have not been reported since their introduction in Georgia, nor along the Russian Black Sea Caucasus coast (Bienkowski and Orlova-Bienkowskaja 2020).

## Acknowledgements

We thank the family Kontselidze for their welcome and for allowing us to take samples in their garden. We also thank Oldřich Nedvěd and an anonymous reviewer for their important and valuable comments which contributed to the improvement of the manuscript. This work was partially supported (AM) by the European Union's Horizon2020 research and innovation program [grant 773 902-SuperPests].

## დასკვნა

საქართველოს ჭიამაიების (*Coccinellidae*) ფაუნა 84 სახეობით არის წარმოდგენილი. მერვევილაძე, ყვავაძის (2002) სახეობების სიას დაემატა ორი სახეობა (*S. biflammulatus* and *S. montazerii*), მაგრამ ჩველისტში შეტანილი ერთი სახეობის (*T. conatus*) არსებობა ქვეყნის ტერიტორიაზე საეჭვოა; ორი სახეობა გადაყვანილია სინონიმში. ნაშრომში წარმოდგენილი 30 სახეობა არ არის რეგისტრირებული პალეარქტიკის ხეშეშფრთიანთა კატალოგში (Kovář 2007).

## References

- Bienkowski AO (2018) Key for identification of the ladybirds (Coleoptera: Coccinellidae) of European Russia and the Russian Caucasus (native and alien species). *Zootaxa* 4472: 233–260. <https://doi.org/10.11646/zootaxa.4472.2.2>
- Bienkowski AO (2020) Identification of the ladybirds (Coleoptera Coccinellidae) of the European part of Russia and Northern Caucasus [In Russian]. A.N. Severtsov Institute of Ecology and Evolution of the Russian Academy of Sciences, Moscow, 140 pp.
- Bienkowski AO, Orlova-Bienkowskaja MJ (2020) History of the Biodiversity of Ladybirds (Coccinellidae) at the Black Sea Coast of the Russian Caucasus in the Last 120 Years – Does the Landscape Transformation and Establishment of *Harmonia axyridis* Have an Impact? *Insects* 11: 824. <https://doi.org/10.3390/insects11110824>
- Bohn U, Gollub G, Hettwer C, Neuhäuslová Z, Raus T, Schlüter H, Weber H (2000–2003) Karte der natürlichen Vegetation Europas [Map of the Natural Vegetation of Europe]. Maßstab / Scale 1:2 500 000. Landwirtschaftsverlag, Münster.
- Centre de Biologie pour la Gestion des Populations (2021) CBGP – Continental Arthropod Collection (V2 edn). Portail Data INRAE. <https://doi.org/10.15454/D6XAKL>

- Chapin EA (1940) New genera and species of lady-beetles related to *Serangium* Blackburn (Coleoptera: Coccinellidae). *Journal of the Washington Academy of Sciences* 30(6): 263–272.
- Che L, Zhang P, Deng S, Escalona H.E, Wang X, Li Y, Pang H, Vandenberg N, Ślipiński A, Tomaszewska W, Liang D (2021) New insights into the phylogeny and evolution of lady beetles (Coleoptera: Coccinellidae) by extensive sampling of genes and species. *Molecular Phylogenetics and Evolution*, 156: e107045. <https://doi.org/10.1016/j.ympev.2020.107045>
- Che LH, Zhang SQ, Li Y, Liang D, Pang H, Ślipiński A, Zhang P (2017) Genome-wide survey of nuclear protein-coding markers for beetle phylogenetics and their application in resolving both deep and shallow-level divergences. *Molecular Ecology Resources* 17(6): 1342–1358. <https://doi.org/10.1111/1755-0998.12664>
- Coutanceau J-P (2006) *Serangium parcesetosum* Sicard 1929 en Corse: réhabilitation et nouvelles localités (Col. Coccinellidae). *Le Coléoptériste* 9(3): 180.
- Coutanceau J-P (2009) *Harmonia conformis* (Boisduval, 1835) : une Coccinelle australienne introduite et acclimatée en France (Col. Coccinellidae). *Bulletin de la Société entomologique de France* 114: 110.
- Coutanceau J-P, Malausa J-C (2014) Coléoptères Coccinellidae introduits en France métropolitaine comme agents de lutte biologique. *Harmonia* 13: 9–21.
- Duverger C (1998) *Serangium montazerii* Fursch 1995, (Coleoptère Coccinellidae) introduit en France et Corse sous le nom de *Serangium parcesetosum* Sicard 1929. *Bulletin de la Société linnéenne de Bordeaux* 26(2): 97–98.
- Escalona HE, Ślipiński A (2012) Generic revision and phylogeny of Microweiseinae (Coleoptera: Coccinellidae). *Systematic Entomology* 37(1): 125–171. <https://doi.org/10.1111/j.1365-3113.2011.00601.x>
- Fürsch H (1995) A new *Serangium*-species from Iran. *Nachrichtenblatt der Bayerischen Entomologen* 44(1/2): 20–21.
- Gourreau J-M (1974) Systématique de la tribu des Scymini (Coccinellidae). *Annales de Zoologie, Ecologie animale Hors-série*, 1–221.
- Iablokoff-Khznorlan SM (1982) Les coccinelles Coléoptères-Coccinellidae: Tribu Coccinellini des régions Paléarctique et Orientale. Société Nouvelle des Éditions Boubée, Paris.
- Iablokoff-Khznorlan SM (1983) A review of the ladybirds in the fauna of the USSR (Coleoptera, Coccinellidae) [in Russian]. *Sbornik, Institut Zoologii, Akademiya Nauk Armianskoi SSR* 19: 94–161.
- Kováč I (2007) Family Coccinellidae Latreille, 1807. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup Vol. IV, 568–631.
- Lombaert E, Guillemaud T, Thomas CE, Lawson Handley LJ, Li J, Wang S, Pang H, Goryacheva I, Zakharov IA, Jousset E, Poland RL, Migeon A, Van Lenteren J, De Clercq P, Berkvens N, Jones W, Estoup A (2011) Inferring the origin of populations introduced from a genetically structured native range by approximate Bayesian computation: case study of the invasive ladybird *Harmonia axyridis*. *Molecular Ecology* 20: 4654–4670. <https://doi.org/10.1111/j.1365-294X.2011.05322.x>
- Malausa JC, Franco E, Brun P (1988) Acclimatation sur la côte d'azur et en corse de *Serangium parcesetosum* [Col.: Coccinellidae] prédateur de l'aleurode des citruses, *Dialeurodes citri* [Hom.: Aleyrodidae]. *Entomophaga* 33(4): 517–519. <https://doi.org/10.1007/bf02373187>
- Merkviladze MS, Kvavadze ES (2002) List of ladybirds (Coleoptera, Coccinellidae) of Georgia. *Proceedings of the Institute of Zoology* 21: 149–155.
- Nattier R, Michel-Salzat A, Almeida LM, Chifflet-Belle P, Magro A, Salazar K, Kergoat GJ (2021) Phylogeny and divergence dating of the ladybird beetle tribe Coccinellini Latreille (Coleoptera: Coccinellidae: Coccinellinae). *Systematic Entomology* 46: 632–648. <https://doi.org/10.1111/syen.12480>
- Orlova-Bienkowskaja MJ, Bieńkowski AO (2017) Alien Coccinellidae (Ladybirds) in Sochi National Park and its vicinity, Russia. *Nature Conservation Research* 2(4): 96–101. <https://doi.org/10.24189/ncr.2017.044>
- Orlova-Bienkowskaja MJ, Spiridonov SE, Butorina NN, Bieńkowski AO (2018) Coinvasion by the ladybird *Harmonia axyridis* (Coleoptera: Coccinellidae) and its parasites, *Hesperomyces virescens* (Ascomycota: Laboulbeniales) and *Parasitylenchus bifurcatus* (Nematoda: Tylenchida, Allantonematidae), in the Caucasus. *PLOS ONE* 13(11): e0202841. <https://doi.org/10.1371/journal.pone.0202841>
- Pang H, Tang X-F, Booth RG, Vandenberg N, Forrester J, Mchugh J, Ślipiński A (2020) Revision of the Australian Coccinellidae (Coleoptera). Genus *Novius* Mulsant of tribe Noviini. *Annales Zoologici* 70: 1–24. <https://doi.org/10.3161/00034541ANZ2020.70.1.001>
- Poorani J (1998) A new species of *Serangium* Blackburn (Coleoptera: Coccinellidae), with a key to species, from India. *Journal of Biological Control* 12: 55–61.
- Rondoni G, Borges I, Collatz J, Conti E, Costamagna AC, Dumont F, Evans EW, Grez AA, Howe AG, Lucas E, Maisonhaute J-É, Onofre Soares A, Zaviezo T, Cock MJW (2021) Exotic ladybirds for biological control of herbivorous insects – a review. *Entomologia Experimentalis et Applicata* 169: 6–27. <https://doi.org/10.1111/eea.12963>
- Roy H, Migeon A (2010) Ladybeetles (Coccinellidae). Chapter 8.4. *Bio-Risk* 4: 293–313. <https://doi.org/10.3897/biorisk.4.49>
- Savenko R (1953) Faune of the coccinellids (Coleoptera Coccinellidae) of Georgia [in Russian]. *Trudy Instituta Zoologii Akademii Nauk Gruzinskoy SSR* 11: 127–140
- Seago AE, Giorgi JA, Li J, Ślipiński A (2011) Phylogeny, classification and evolution of ladybird beetles (Coleoptera: Coccinellidae) based on simultaneous analysis of molecular and morphological data. *Molecular Phylogenetics and Evolution* 60: 137–151. <https://doi.org/10.1016/j.ympev.2011.03.015>
- Schneider O, Leder H (1878) Beiträge zur Kenntniss des kaukasischen Käferfauna. 346–353.
- Szawaryn K, Tomaszewska W (2020) New and known extinct species of *Rhyzobius* Stephens, 1829 shed light on the phylogeny and biogeography of the genus and the tribe Coccidulini (Coleoptera: Coccinellidae). *Journal of Systematic Palaeontology* 18: 1445–1461. <https://doi.org/10.1080/14772019.2020.1769751>
- Timofeeva TV, Nhuan HD (1978) Morphological and biological characteristics of Indian lady-bird *Serangium parcesetosum* Sicard (Coleoptera, Coccinellidae), predator of the citrus whitefly in Adzharia. *Entomologicheskoe Obozrenie* 57(2): 302–308.
- Tomaszewska W, Escalona HE, Hartley D, Li J, Wang X, Li H-S, Pang H, Ślipiński A, Zwick A (2021) Phylogeny of true ladybird beetles (Coccinellidae: Coccinellini) reveals pervasive convergent evolution and a rapid Cenozoic radiation. *Systematic Entomology* 46: 611–631. <https://doi.org/10.1111/syen.12479>
- Tomaszewska W, Szawaryn K (2016) Epilachnini (Coleoptera: Coccinellidae) A revision of the world genera. *Journal of Insect Science* 16: 1–91. <https://doi.org/10.1093/jisesa/iew082>
- Yigit A, Canhilal R (2005) Establishment and dispersal of *Serangium parcesetosum* Sicard (Coleoptera, Coccinellidae), a predatory beetle of citrus whitefly, *Dialeurodes citri* Ashm. (Homoptera, Aleyrodidae) in the East Mediterranean region of Turkey. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 112(3): 268–275.
- Zhang SQ, Che LH, Li Y, Liang D, Pang H, Ślipiński A, Zhang P (2018) Evolutionary history of Coleoptera revealed by extensive sampling of genes and species. *Nature Communications* 9(1): 205. <https://doi.org/10.1038/s41467-017-02644-4>