



New data on some microhymenopteran families from Lagodekhi Protected Area, with new records for Georgia (Sakartvelo) and the Caucasus

George Japoshvili^{1,2}

1 *Institute of Entomology, Agricultural University of Georgia, Digomi David Aghmashenebeli Alley, #240, Tbilisi – 0119, Georgia*

2 *Invertebrate Research Center, Tetrtsklebi, Telavi Municipality – 2200, Telavi, Georgia*

<http://zoobank.org/19F7E4A8-F25E-41C0-BFC0-2AFC31E8BE61>

Corresponding author: George Japoshvili (giorgij70@yahoo.com)

Academic editor: Levan Mumladze ♦ **Received:** 3 February 2022 ♦ **Accepted:** 5 April 2022 ♦ **Published:** 20 June 2022

Abstract

We present data on the distribution of seven wasp families, Ceraphronidae, Diapriidae, Ismaridae, Megaspilidae, Platygasteridae, Proctotrupidae and Scelionidae, collected in Georgia (Sakartvelo). Thirty-three genera are recorded for the first time from Georgia and 25 of them for the Caucasus. From these, 16 species level taxa are newly recorded in Georgia, and 14 identified species are new records for the Caucasus. This brings the number of known insect species in the Lagodekhi Protected Area (LPA) to 1682.

Key words

Ceraphronidae, Diapriidae, Megaspilidae, Platygasteridae, Proctotrupidae, Scelionidae, parasitic wasps, South Caucasus

Introduction

Except for some families belonging to the Chalcidoidea superfamily, microhymenopterans in Georgia (Sakartvelo) are scarcely studied. Notwithstanding this neglect, the group contains some of the most important biological control agents of phytophagous insects, in particular selected species of Scelionidae (Japoshvili et al. 2022). Moreover, parasitoid wasps are a major component of many terrestrial ecosystems, and may constitute up to 20% of all insect species (LaSalle and Gauld 1991; Godfray 1994; Memmott et al. 1994). Here, I present hymenopteran survey results conducted in Lagodekhi Protected Area (LPA). The LPA was established in 1912, and is located in the extreme north-eastern part of Georgia along the southern slope of the Greater Caucasus Mountains. Elevations in this region range from 590 to 3500 m. The LPA is among the most diverse and best-studied protected areas in Georgia, with 1666 insect

species records to date (Japoshvili et al. 2022). More than 10 publications have been published during last 10 years detailing hundreds of new records for fauna within the LPA and Georgia (Japoshvili et al. 2016, 2017, 2019; Japoshvili and Kostjukov 2016, 2017; Kostjukov et al. 2016a,b; 2017; Kostjukov and Japoshvili 2016a,b; Gul Aslan et al. 2017; Japoshvili 2017, 2018; Dobosz et al. 2018; Dvorak et al. 2018; Otto and Japoshvili 2018; Riedel et al. 2018; Japoshvili and Khachikov 2021). Here we provide additional new faunistic records of hymenopterans in the LPA, documenting the first records of these taxa in Georgia and/or the Caucasus.

Material and methods

We collected samples in Lagodekhi Protected Area using malaise traps during the entire growing season, from April to

November of 2014. Malaise traps were placed in seven different areas with the following vertical zonal characteristics: H1: low-altitude forest (450–750 m), 666 m, N41.852483, E46.287767; H2: mid-altitude forest (750–1250 m), 847 m, N41.85585, E46.292733; H3: high-altitude forest (1250–1800 m), 1351 m, N41.871467, E46.311533; H4: subalpine forest (1800–2000 m), 1841 m, N41.882733, E46.32185; H5: subalpine forest (1800–2000 m), 1902 m, N41.885583, E46.324117; H6: Subalpine meadows and shrublands (2000–2500 m), 2230 m, N41.89805, E46.333883; H7: alpine zone (above 2500 m), 2559 m, N41.906183, E46.3334.

Malaise traps were obtained from B & N Entomological services (<http://www.entomology.org.uk/>). Traps were emptied every 10 (± 2) days and stored specimens in 96% alcohol. Subsequently, samples were CPD dried and card mounted (Noyes, 2019). After sorting, all microhymenoptera not belonging to the Chalcidoidea superfamily and/or Cynipoidea were sent to Dr Lubomir Masner, who screened, sorted and prepared the specimens for further determination. Identifications were done by L. Masner using published keys (Medvedev 1978; Townes and Townes 1981; Kozlov and Kononova 1983).

All voucher and type specimens were deposited in the entomological collection of the Agricultural University of Georgia, Tbilisi, Georgia. New records for Georgia are indicated with one asterisk, two asterisks next to the taxon name indicates the taxon, new to the Caucasus.

Results and discussion

Annotated list of species

Order: Hymenoptera

Family: Diapriidae Haliday, 1833

- Only two species, *Psilus punctiventris* (Kozlov) and *Spilomicrus cursor* Kieffer, were known from Georgia (Aphazeti and Achara (Batumi)), according to Medvedev (1978).

Genus: *Entomacis* Förster, 1856**

Entomacis perplexa (Haliday, 1857)

- GEORGIA • 1 Female; H3; 3.V.2014. 1 Female; H4-5; 25.VI-5.VII.2014. 1 Female; H6; 23.V-13.VI.2014.

Entomacis platyptera (Haliday, 1857)

- GEORGIA • 1 Female; H1; 25.V-4.VI.2014.

Family: Ismaridae Thomson, 1858**

- This family was historically considered as a subfamily of Diapriidae. However, based on recent integrative taxonomic investigation, the Ismarinae subfamily was elevated to family rank including the single genus *Ismarus* Haliday (Sharkey et al. 2012). There are no previous records of this family occurring in the Caucasus region.

Genus: *Ismarus* Haliday, 1835

Ismarus dorsiger (Haliday, 1831)

- GEORGIA • 1 Female; H5; 5-15.VII.2014. 1 Female; H6; 26.VII-6.VIII.2014.

Ismarus haliday Foerster, 1850

- GEORGIA • 1 Female, 1 Male; H6; 23.V-13.VI.2014.

Family: Megaspilidae Ashmead, 1893

- Only one genus and one species, *Megaspilus dux* (Curtis), was previously known from Georgia, according to Alekseev and Radchenko (2001).

Genus: *Trichosteresis* Förster, 1856**

Trichosteresis glabra (Boheman, 1832)

- GEORGIA • 2 Females; H5; 23.V-13.VI.2014. 2 Females; H5; 5-15.VII.2014. 1 Male; H5; 26.IX-6.X.2014.

Family: Platygastriidae Haliday, 1833

- Prior to our study, 5 species representing 5 genera were recorded from Georgia, including *Allotropa conventus* (Maneval) and *Synopeas figitifformis* Thomson (Medvedev 1978), *Acerotella boter* (Walker), *Amblyaspis aliena* (Nees), *Leptaces laodices* (Walker) (Timokhov 2019).

Genus: *Iphitrachelus* Haliday, 1836**

Iphitrachelus gracilis Masner, 1957

- GEORGIA • 1 Male; H1; 4-14.VI. 2014.

Iphitrachelus lar Haliday, 1835

- GEORGIA • 1 Female; H1; 25.V-4.VI.2014. 1 Female; H6; 5-15.VIII.2014.

Genus: *Metaclisis* Förster, 1856**

Metaclisis areolata (Haliday, 1835)

- GEORGIA • 1 Female; H1; 17-23.IV.2014.

Genus: *Platystasius* Nixon, 1937**

Platystasius transversus (Thomson, 1859)

- GEORGIA • 1 Female; H1; 5-15.VI.2014. 1 Female; H2; 25.V-4.VI.2014. 1 Female; H2; 25.VI-5.VII.2014. 1 Female; H6; 5-15.VII.2014.

Family: Proctotrupidae Latreille, 1802

- According to Lelej (2012), the following 19 species have been recorded from Georgia prior to our studies: *Brachyserphus parvulus* (Nees), *Codrus picicornis* (Förster), *Cryptoserphus aculeator* (Haliday), *C. dilates* Townes, *C. flavipes* (Provancher), *C. medius* Townes, *Disogmus areolator* Haliday, *D. basalis* (Thomson), *Exallonyx ater* (Gravenhorst), *E. brevicornis* (Haliday), *E. crenicornis* (Nees), *E. minor* Townes, *E. wasmani* Kieffer, *Mischoserphus obesus* Townes, *Phaenoserphus chittii* (Morley), *P. nigripes* (Ashmead), *P. viator* (Haliday) and *Proctotrupes gravidator* (Linnaeus). *M. arcuator* (Stelfox) was recorded by Izadihzhadeh et al. (2016) from Georgia.

Genus: *Brachyserphus* Hellen, 1941

Brachyserphus parvulus (Nees, 1834)

- GEORGIA • 4 Females, 1 Male; H5; 5-15.VII.2014.

Genus: *Disogmus* Förster, 1856

Disogmus areolator (Haliday, 1839)

- GEORGIA • 3 Females, 4 Males; H1; 25.V-4.VI.2014. 1 Female, 1 Male; H2; 15-25.V.2014. 1 Female; H4-5; 25.VI-5.VII.2014. 2 Females; H5; 5-15.VII.2014. 2 Males; H6; 23.V-13.VI.2014.

Genus: *Exallonyx* Kieffer, 1904

Exallonyx brevicornis (Haliday, 1839)

- GEORGIA • 2 Females; H4-5; 25.VI-5.VII.2014. 1 Female; H6; 5-15.VII.2014. 1 Female; H5; 23.V-13.VI.2014.

Mischoserphus arcuator (Stelfox, 1950)

- GEORGIA • 1 Female; H1; 25.V-4.VI.2014. 3 Females; H4-5; 25.VI-5.VII.2014.

Genus: *Phaenoserphus* Kieffer, 1908

Phaenoserphus calcar (Haliday, 1839)**

- GEORGIA • 1 Male; H4; 5-15.VII.2014.

Genus: *Tretoserphus* Townes, 1981**

Tretoserphus laricis (Haliday, 1839)

- GEORGIA • 4 Females; H3; 3.V.2014.

Family: Scelionidae Haliday, 1839

- Nineteen species belonging to the Scelionidae family have been recorded prior to our studies from Georgia: *Duta tenuicornis* (Dodd), *Proteleas tridentatus* Kozlov, *Telenomus chloropus* (Thomson), *T. hofmanni* Mayr, *T. oophagus* Nikolskaja, *Trimorus elongatus* (Kieffer), *T. pedes* (Kieffer), *Trissolcus tumidus* (Mayr) (Lelej 2012), *Encyrtoscelio apterus* (Szelenyi), *Eremioscelio cydnoides* Priesner, *Psilanteria bicolor* (Kieffer), *Teleas quinquespinosus* Szabo, *T. sibiricus* Kieffer and *T. sulcatus* (Kozlov) (Timokhov 2019b). Recently, Japoshvili et al. (2021) reported following species from Georgia: *T. belenus* (Walker), *T. colemani* (Crawford), *T. cultratus* (Mayr), *T. scutellaris* (Thomson), *T. semistratus* (Nees).

Genus: *Idris* Förster, 1856**

Idris pedestris (Kieffer, 1908)

- GEORGIA • 1 Female; H1; 26.VII-6.VIII.2014. 1 Female; H1; 17.IX-6.X.2014. 1 Female; H2; 20.VII-5.VIII.2014. 1 Female; H2; 23.VIII.-4.IX.2014.

Genus: *Scelio* Latreille, 1805*

Scelio rugulosus Latreille, 1805**

- GEORGIA • 1 Female; H1; 25.V-4.VI. 1 Female; H5; 5-15.VIII.

Genus: *Teleas* Latreille, 1809

Teleas lamellatus Szabo, 1956*

- GEORGIA • 1 Female; H5; 5-15.VIII.2014.

Genus: *Trissolcus* Ashmead, 1904

Trissolcus basalis (Wollaston, 1858)*

- GEORGIA • 1 Female; H2; 5-15.VII.2014. 1 Female; H5; 23.V-13.VI.2014.

Trissolcus cultratus (Mayr, 1879)

- GEORGIA • 2 Females; H1; 26.VII-6.VIII.2014. 1 Female; H2; 25.V-1.VI.2014. 1 Female; H3; 12-31.V.2014. 13 Females; H6; 23.V-13.VI.2014. 1 Female; H6; 5-15.VII.2014.

Genus: *Xenomerus* Walker, 1836**

Xenomerus ergenna Walker, 1836

- GEORGIA • 10 Females; H6; 25.VI-5.VII.2014. 1 Female; H5; 15-25.VI.2014.

Additional remarks

In addition, a number of taxa were recorded in the LPA which have unknown species identities. These include *Aphanogmus* sp. (Ceraphronidae) (1 Female; H1; 5-15.VI.2014) (from this genus, the only species, *A. steinitzi* Priesner, was recorded from Achara region of Georgia (Medvedev 1978)), *Acanosema* sp.** (1 Female; H1; 25.VI.-5.VII.2014), *Acropiasta* sp.** (1 Female; H2; 15-25.VI.2014), *Aneurhynchus* sp.** (1 Female; H3; 15-25.VI.2014), *Basalys* sp.** (1 Female; H3; 15-25.VI.2014), *Pantoclis* sp.** (1 Female; H4; 5-15.VII.2014), *Paramesius* sp.** (1 Male; H1; 5-15.VI.2014), *Synacra* sp.** (1 Female; H6; 5-15.VII.2014), *Trichopria* sp.** (1 Female; H6; 25.VII-3.VIII.2014), *Cinetus* sp.* (1 Female; H4; 5-15.VII.2014) (all from the family Diapriidae), *Atritomellus* sp.** (1 Female; H6; 23.V-3.VI.2014), *Conostigmus* sp.** (1 Female; H3; 3.V.2014), *Dendrocerus* sp.* (1 Female; H6; 5-15.VII.2014), *Lagynodes* sp.* (1 Female; H6; 15-25.VII.2014) (all from the family Megaspilidae), *Acerotella* sp. (1 Female; H2; 15-25.V.2014), *Allotropa* sp. (1 Female; H6; 15-25.VII.2014), *Amblyaspis* sp. (1 Female; H6; 15-25.VII.2014), *Isocybus* sp.** (1 Female; H6; 25.VI.-5.VII.2014), *Piestopleura* sp.** (1 Female; H6; 5-15.VII.2014), *Platygaster* sp.** (1 Female; H3; 3.V.2014), *Pyrgaspis* sp.** (1 Female; H5; 5-15.VIII.2014), *Tetrabaenus* sp.** (1 Female; H1; 26.VII.-6.VIII.2014), *Leptacis* sp.* (1 Male; H1; 5-15.VI.2014. 1 Female; H6; 5-15.VII.2014), *Synopeas* sp.* (1 Female; H6; 5-15.VIII.2014), *Cryptoserphus* sp. (1 Female; H4-5; 25.VI.-5.VII.2014) (all from the family Platygasteridae), *Anteris* sp.** (1 Female; H1; 5-15.VI.2014), *Gryon* sp.* (1 Female; H1; 5-15.VI.2014), *Sparasion* sp.* (1 Male; H5; 5-15.VIII.2014), *Telenomus* sp. (1 Male; H5; 5-15.VII.2014) and *Trimorus* sp. (1 Male; H6; 5-15.VII.2014) (all from the family Scelionidae).

In summary, new records to Georgia or the Caucasus were identified, and a number of unidentified taxa belonging to seven wasp families, Ceraphronidae, Diapriidae, Ismaridae, Megaspilidae, Platygasteridae, Proctotrupidae and Scelionidae, were collected in the LPA. Sixteen of these species were new country records for Georgia (14 new for the Caucasus) and thus the number of insect species recorded from LPA has reached 1682 species (Japoshvili et al. 2022). From the species identified to morpho species level, 33 genera, have been newly recorded from Georgia and 25 of them are new for the Caucasus. Undoubtedly, the current knowledge of hymenopterans of Georgia and the Caucasus in general is incomplete and further research will almost certainly reveal many new taxa.

Acknowledgements

We would like to express our gratitude to Dr Lubomir Masner, Research Associate (Agriculture and Agri-Food Canada, K.W. Neatby Building, Ottawa, Ontario K1A 0C6, Canada), for his kind help in species identifications. We also thank the Shota Rustaveli National Science Foundation for financial support (project ref. FR/221/7-110/13).

დასკვნა

ნაშრომში წარმოდგენილია სიფრიფანაფრთიანების კვლევის შედეგები ლაგოდეხის დაცული ტერიტორიებიდან. შესწავლილი იქნა შვიდი ოჯახი (Ceraphronidae, Diapriidae, Ismaridae, Megaspilidae, Platygastriidae, Proctotrupidae and Scelionidae) რომელთაგან ერთი ოჯახი წარმოადგენს პირველ შეტყობინებას კავკასიის რეგიონისთვის. 16 სახეობა ახალია საქართველოსთვის, მათგან 14 კი კავკასიისთვის. ასევე, სახეობებიდან, რომლებიც გაირკვა მორფო სახეობის დონეზე 33 გვარი ახალია საქართველოსათვის, მათგან 25 კი კავკასიისთვის. კვლევის შედეგების შედეგად ლაგოდეხის დაცული ტერიტორიებისთვის აღწერილ მწერთა სახეობების რაოდენობა გაიზარდა 1682 სახეობამდე.

References

- Alekseev V, Radchenko T (2001) Cerapronoid wasps (Hymenoptera, Cerapronoidea) of the fauna of the Ukraine. *Vestnik Zoologii* 35(3):3-16.
- Dobosz R, Japoshvili G, Krivokhatsky V, Wařala R (2018) Contributions to the knowledge of neuropterid insects (Neuropterida: Raphidioptera, Neuroptera) of Georgia (Sakartvelo). Part II. *Annals of the upper Silesian museum in bytom entomology* 26:1-21.
- Dvorak L, Japoshvili G (2018) The scorpionflies (Mecoptera) from Lagodekhi Reserve, with some distributional and seasonal activity data. *Zootaxa* 4407(1):137-140.
- Godfray H (1994) *Parasitoids: Behavioural and Evolutionary Ecology*. Princeton University Press, Princeton, 473 pp.
- Gul Aslan E, Mumladze L, Japoshvili G (2017) List of leaf beetles (Coleoptera: Chrysomelidae) from Lagodekhi reserve with new records for Transcaucasia and Georgia. *Zootaxa* 4277(1):86-98.
- Izadihzhadeh M, Talebi A, Kolyada V (2016) First record of the family Proctotrupidae (Hymenoptera: Proctotrupoidea) from Iran, with five new species records. *Turkish Journal of Zoology* 40:27-34.
- Japoshvili G (2017) New records of Encyrtids (Hymenoptera: Chalcidoidea: Encyrtidae) from Georgia, with description of seven new species. *Journal of Asia-Pacific Entomology* 20(3):866-877.
- Japoshvili G (2018) New records of *Centrodora* Forster (Hymenoptera: Aphelinidae) from the Caucasus. *Turkish Journal of Zoology* 42:118-120.
- Japoshvili G, Arabuli T, Salakaia M, Tskaruashvili Z, Kirkitadze G, Talamas E (2022) Surveys for *Halyomorpha halys* (Stål)(Hemiptera: Pentatomidae) and its biocontrol potential by parasitic wasps in the Republic of Georgia (Sakartvelo). *Phytoparasitica* 50(1):127-137.
- Japoshvili G, Gordiashvili N, Injia S, Tsiklauri Kh, Sulamanidze G (2022) Updated annotated checklist of insects from Lagodekhi protected areas, Sakartvelo (Georgia). *Munis Zoology and Entomology* 17(2): 793-841.
- Japoshvili G, Kostjukov V (2016) The First Record of *Pronotalia* Gradwell, 1957 (Hymenoptera, Eulophidae) Wasps from Georgia. *Entomological Review* 96(8):1089-1091.
- Japoshvili G, Kostjukov V (2017) Three New Minotetrastichus (Hymenoptera: Eulophidae) Species Records from Transcaucasia. *Journal of the Entomological Research Society* 19(2):95-100.
- Japoshvili G, Kostjukov V, Kosheleva O (2016a) First record of *Anaprostocetus* Graham, 1987 (Hymenoptera: Eulophidae) from Transcaucasia, Biological plant protection as the basis of ecosystem stabilization 9:125-128.
- Japoshvili G, Kostjukov V, Kosheleva O (2016b) New Record of *Tamarixia* (Hymenoptera: Eulophidae) from Transcaucasia with Some Taxonomic and Biogeographical Notes. *Journal of Entomological Research Society* 18(3):113-120.
- Japoshvili G, Kostjukov V, Kosheleva O, Podvarko A (2019) New Record of *Aprostocetus caudatus* Species Group (Hymenoptera, Eulophidae) from Georgia. *Journal of Entomological Research Society* 21(3):361-367.
- Japoshvili G, Kostjukov V, Kosheleva O, Yegorenkova E (2017) New Record of *Neotrichoporoides* (Hymenoptera: Eulophidae) from Georgia with some Taxonomic and Biogeographical Notes. *Journal of the Entomological Research Society* 19(2):67-71.
- Japoshvili G, Mumladze L (2019) New species and new records of *Aphelinus* Dalman (Hymenoptera: Chalcidoidea: Aphelinidae) from Lagodekhi Reserve (Sakartvelo--Georgia), with diversity and distribution along an elevational gradient. *Turkish Journal of Zoology* 43:192-202.
- Khachikov E, Iljina E, Gasanova N, Japoshvili G (2021) Preliminary faunistic study on the rove beetles (Coleoptera: Staphylinidae) of Republic of Dagestan, with some additional information from northern Caucasus. *Zootaxa* 4950(2):248-266.
- Kostjukov V, Japoshvili G (2016) First record of *Holcotetrastichus* Graham, 1987 (Hymenoptera: Eulophidae) from Transcaucasia. *Biological plant protection as the basis of ecosystem stabilization* 9: 148-151.
- Kostjukov V, Japoshvili G (2016) First record of Murcia-group of the genus *Tetrastichus* (Hymenoptera: Eulophidae) from Transcaucasia. *Biological plant protection as the basis of ecosystem stabilization* 9:162-166
- Kostjukov V, Japoshvili G (2018) New record of *Hyperteles* (Hymenoptera: Eulophidae) species from Transcaucasia. *Annals of Agrarian Science* 16(1):42-44.
- Kostjukov V, Kosheleva O, Japoshvili G (2016) First record of *Mischotetrastichus* Graham, 1987 (Hymenoptera, Eulophidae) from Transcaucasia. *Eurasian Entomological Journal* 15(5):464-465.
- Kozlov M, Kononova S (1983) *Telenominae of the USSR*. Nauka, Leningrad, 336 pp. (In Russian)
- Lelej AS (2012) *Annotated catalogue of the Insects of Russian Far East. Volume 1. Hymenoptera*. Dalnauka, Vladivostok, 635 pp.
- LaSalle J, Gauld I (1991) Parasitic Hymenoptera and the biodiversity crisis. *Redia* 74:315-334.
- Medvedev G (1978) *Keys to the Insects of the European part of USSR. Vol. 3 Part 2*. Nauka, Leningrad, 756 pp. (In Russian)
- Memmott J, Godfray H, Gauld I (1994) The structure of a tropical host-parasitoid community. *Journal of Animal Ecology* 63:521-540.
- Mikhailov K, Otto S, Japoshvili G (2017) A new species from the *Clubiona caerulea* group from the Caucasus (Araneae: Clubionidae). *Zoology in the Middle East* 63(4):362-368.
- Mumladze L, Ulrich W, Asanidze Z, Japoshvili G (2017) An inverse elevational species richness gradient of Caucasian vascular plants and Encyrtidae (Hymenoptera, Chalcidoidea). *Écoscience* 24(1-2):75-79.
- Noyes JS (2019) *Universal Chalcidoidea Database*. World Wide Web electronic publication. <https://www.nhm.ac.uk/our-science/data/chalcidoidea/database/>.
- Otto S, Japoshvili G (2018) The spiders (Arachnida: Araneae) of the Lagodekhi Reserve, Georgia: faunistic results of a transect study and an updated checklist. *Arachnology* 17(8):375-391.
- Riedel M, Diller E, Japoshvili G (2018) The Ichneumonid fauna (Hymenoptera: Ichneumonidae) of Lagodekhi Reserve, Sakartvelo (Georgia), with descriptions of four new species. *Linzer biologische Beiträge* 50(2):1447-1507.
- Sharkey MJ, Carpenter JM, Vilhelmsen L, Heraty J, Liljebld J, Dowling AP, Schulmeister S, Murray D, Deans AR, Ronquist F, Krogmann L,

- Wheeler WC (2012). Phylogenetic relationships among superfamilies of Hymenoptera. *Cladistics* 28(1):80-112.
- Talamas E, Buffington M, Hoelmer K (2017) Revision of Palearctic *Trisolcus* Ashmead (Hymenoptera: Scelionidae). *Journal of Hymenoptera Research* 56:3-185.
- Timokhov A (2019a) New records of platygastrid wasps (Hymenoptera: Platygastridae) from Russia and Georgia. *Proceedings of the Russian Entomological Society* 90:9-12.
- Timokhov A (2019b) New data and corrections to the fauna of scelionid wasps (Hymenoptera: Scelionidae) of Russia. *Proceedings of the Russian Entomological Society* 90:13-21.
- Townes H, Townes M (1981) A revision of the Serphidae (Hymenoptera). *Memoirs of the American Entomological Institute* 32:1-541