ÖGH Herpetozoa
Austrian Herpetological Society

Herpetozoa 37: 1–10 (2024)
DOI 10.3897/herpetozoa.37.e111749

Reproductive effort of *Plestiodon copei* (Squamata, Scincidae), a highland viviparous lizard from Central Mexico

Hermilo Sánchez-Sánchez[1], Adriana Jocelyn Morales-Gonzaga[2], Ken Oyama[3], J. Carlos Alvarado-Avilés[2]

1 *Centro de Investigación en Recursos Bióticos (CIRB), Universidad Autónoma del Estado de México (UAEMéx), Km. 14.5 Carretera Toluca-Ixtlahuaca, Toluca, Estado de México, CP 50200, Mexico*
2 *Facultad de Ciencias, Universidad Autónoma del Estado de México (UAEMéx), Km. 15.5 Carretera Toluca-Ixtlahuaca, Toluca, Estado de México, CP 50200, México*
3 *Escuela Nacional de Estudios Superiores (ENES) Unidad Morelia, Universidad Nacional Autónoma de México (UNAM), Antigua Carretera a Pátzcuaro No. 8701, Col. Ex-Hacienda de San José de la Huerta, Michoacán, Morelia, CP 58190, Mexico*

https://zoobank.org/373AA447-7987-4745-B7C8-29ABC892B395

Corresponding author: J. Carlos Alvarado-Avilés (jalvaradoa@uaemex.mx)

Academic editor: Johannes Foufopoulos ♦ **Received** 28 August 2023 ♦ **Accepted** 14 December 2023 ♦ **Published** 8 January 2024

Abstract

Female reproductive effort is defined as the proportion of total energy or resources devoted to reproduction. In reptiles, there is frequently high inter- and intra-population variation related to several factors, such as food availability, climatic conditions, age and size, all of which, in turn, also influence survival and future reproduction. The present study is the first reproductive effort analysis of a population of the high-mountain scincid lizard *Plestiodon copei* in central Mexico, focusing on relative litter mass (RLM), investment per capita (INV) and productivity (PROD). We also compared the reproductive efficacy of *P. copei* to those of other Mexican congeners. We collected 24 gravid females of *P. copei* over a 4-year period and recorded a total of 90 neonates born in captivity. We found significant variation in neonatal mass amongst individual females and across years. We recorded an average litter size of 3.75 and an average litter mass of 1.25 g, which were positively correlated with both the size and total mass of the reproductive females. The RLM, INV and PROD values (0.301, 0.428 g and 1.236 g × year^{-1}, respectively) for *P. copei* were greater than those for other Mexican species in the *P. brevirostris* group (0.290, 0.412 g and 1.135 g × year^{-1}), revealing that females of *P. copei* from Tenango invest a comparatively high amount of their resources in reproduction.

Key Words

life history, litter mass, litter size, reproductive investment, skink lizard, Trans-Mexican Volcanic Belt

Introduction

Reproductive investment is a central aspect of the theory of life history evolution, which states that an individual must allocate an optimal proportion of its available energy to reproduction which, in turn, will be reflected in statistics, such as in neonate/egg size and mass, litter/clutch size and mass and reproductive effort (Stearns 1992; Schwarzkopf 1993; Shine 2005). In ectothermic organisms, reproductive investment is strongly affected by both physiological and ecological constraints. In reptiles, the reproductive investment of females is strongly influenced by body size and pelvic opening (Congdon and Gibbons 1987; Sinervo and Licht 1991; Rodríguez-Romero et al. 2002, 2004, 2005; Ramírez-Bautista et al. 2016, 2017; Suárez et al. 2018). However, local environmental conditions also affect the reproductive investment of females and can lead to significant variations in the number and mass of eggs or

Copyright *Hermilo Sánchez-Sánchez et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1. A) Adult female and **B**) neonate of *Plestiodon copei*. Photo Credits **A:** H. Sánchez-Sánchez; **B:** JC. Alvarado-Avilés.

neonates (Stearns 1992; Roitberg et al. 2013; Muñoz 2015; Suárez et al. 2018). The metabolic rate, food intake and environmental temperature are also related to reproductive investment, whereby lizard species inhabiting temperate environments may accelerate their sexual maturation and reproduce at smaller sizes, leading to smaller clutches/litters (Warne and Charnov 2008; Meiri et al. 2012).

Female lizards can also exhibit remarkable inter- and intraspecific variations in their reproductive life history traits (Cruz-Elizalde and Ramírez-Bautista 1998; López-Alcaide et al. 2020) due to environmental factors associated with temperature and precipitation regimes, as well as evolutionary history factors, such as phylogenetic constraints and adaptations to adverse environmental conditions, such as cold temperatures in high-elevation forests (Vitt and Price 1982; De Marco 1989; Rohr 1997; Abell 1999; Shine 2005; Castro-Franco et al. 2011).

In the mountainous regions of Mexico, several studies have analysed the reproductive strategies utilised in different genera of lizards, such as *Barisia* (Guillette and Casas-Andreu 1987), *Phrynosoma* (Zamudio and Parra-Olea 2000; Suárez et al. 2018), *Sceloporus* (Feria-Ortiz et al. 2001; Rodríguez-Romero et al. 2002, 2004, 2005; Bastiaans et al. 2013; Ramírez-Bautista et al. 2016; Maciel-Mata 2017; López-Alcaide et al. 2020) and *Plestiodon* (Guillette 1983; Vitt and Cooper 1986; Ramírez-Bautista et al. 1996, 1998; Ramírez-Bautista and Arizmendi 2004; Feria-Ortiz et al. 2007; López et al. 2008; García 2009; Chávez 2012; Laguna 2014; Morales 2014; Muñoz 2015; Bañuelos et al. 2016), in which both the characteristics of females and their progeny were evaluated. Some studies on the reproduction of sympatric or phylogenetically related species of the genus *Plestiodon* indicate that inter- and intra-population variations in reproductive patterns are related to environmental heterogeneity and can lead to local adaptations (Mathies and Andrews 1995; Wapstra and O'Reilly 2001; Rodríguez-Romero et al. 2002, 2004, 2005; Shine 2005; Chávez 2012; Ramírez-Bautista et al. 2016; Maciel-Mata 2017).

To estimate the reproductive investment of female reptiles, several indices have been developed that take into consideration the characteristics of both females and their progeny. Relative clutch mass/relative litter mass (RCM/RLM = fraction of female body mass devoted to reproduction) is the most frequently used index for estimating reproductive effort in lizards for both oviparous and viviparous species (Tinkle and Hadley 1975; Vitt and Congdon 1978; Vitt and Price 1982; Shine and Schwarzkopf 1992; Suárez et al. 2018; Padilla-Pérez et al. 2022). Other indices that have been estimated for several lizard species include the productivity index (PROD = total mass of offspring produced in one year) and the per capita reproductive investment index (INV = investment per capita in the offspring) (Meiri et al. 2012; Rutschmann et al. 2016).

Plestiodon copei (Fig. 1) is an endemic and high-elevation specialist skink lizard distributed in temperate pine and pine-oak forests (i.e. elevation range ca. 2466–3966 m) in the Trans-Mexican Volcanic Belt (TMVB) (Alvarado-Avilés et al. 2020). *Plestiodon copei* is a species with asynchronous gonadal activity and an autumn reproductive period (Guillette 1983; Ramírez-Bautista et al. 1996). This species produces one litter per year and apparently displays parental care, but little is known about the reproductive investment of the females of the species. The present study aimed to characterise the reproductive strategies adopted by females of *P. copei* in a mountainous region of Central Mexico by estimating the reproductive effort under extreme environmental conditions that can lead to different adaptive responses relative to those of other species of the same genus.

Materials and methods

We collected 24 late-pregnant females of *P. copei* from March–early April 2014 to 2017 near Tenango del Valle (19°05'54.74"N, 99°38'26.43"W) at an elevation of ca. 3030 m in the TMVB of central Mexico. Gravid females of *P. copei* were identified *in situ* by their greater abdominal volume. All females were captured in a landscape dominated by pine forests (*Pinus teocote*, *P. montezumae* and *P. rudis*) and grasslands (*Muhlenbergia macroura*) and surrounded by agricultural fields (Alvarado-Avilés et al. 2020).

We transported the collected females to the Laboratory of Genetic and Molecular Evolution (Universidad Autónoma del Estado de México) and individually housed them in plastic boxes (400 × 220 × 180 mm) with natural substrate (e.g. soil, gravel, mulch) and refuge objects (e.g. rocks and bark) obtained from the capture site. All females were maintained according to the thermal conditions described by García (2009), were fed live

food every two days (crickets and mealworms) and were supplied with water *ad libitum*.

Reproductive effort estimation

We recorded the snout–vent length (SVL) and the total mass (TM = female body mass before birth) of each gravid female after capture. Females were checked at least once per day for 2 to 3 weeks and weighed daily until parturition. After birth, the following data were immediately recorded: absolute mass (AM = female body mass after birth), litter size (LS = number of neonates produced by a female), litter mass (LM = sum of the masses of all neonates produced by a female) and mean litter mass (MLM = average mass of the neonates of each litter).

The reproductive effort of *P. copei* females was estimated as follows: (a) relative litter mass; RLM = LM/AM; (b) reproductive investment per capita; INV = (TM−AM)/LS; and (c) estimated productivity; PROD = (MLM × LS) × year^{-1} (Tinkle and Hadley 1975; Vitt and Congdon 1978; Vitt and Price 1982; Rodríguez-Romero et al. 2002, 2004, 2005; Castro et al. 2011; Meiri et al. 2012; Bastiaans et al. 2013; Muñoz 2015; Rutschmann et al. 2016; Suárez et al. 2018). After all the measurements were made, both the females and neonates were released at the place where the female was captured. We also estimated the reproductive effort of other Mexican *Plestiodon* species by examining published data. Recently, taxonomic changes were considered for undescribed species (Pavón-Vázquez et al. 2018).

Data analysis

We checked all variables for normality using the Shapiro–Wilk test and for the presence of outliers with box plots in SPSS ver. 24 (IBM Corporation 2016). To estimate whether the variations in litter characteristics could be explained by female traits, we performed linear regression analysis on the LS, LM, RLM, INV and PROD variables, using the SVL, TM and AM of the females as independent variables. To evaluate whether female mass affects reproductive characteristics, we conducted two regression analyses using LS and MLM as dependent variables and the residuals of the regression of AM on SVL (a measure of robustness) as independent variables (Castro-Franco et al. 2011). We estimated the coefficient of variation (CV = standard deviation/mean × 100%), a standardised measure of the dispersion of a frequency distribution (Sokal and Rohlf 2012; Hutchings 2021) of SVL, LS, LM, MLM and the three reproductive effort indices; values whose CVs were ≤ 5% were considered invariant and those whose CVs were > 5% were considered variable, as described by Castro-Franco et al. (2011) and Suárez et al. (2018) in studies on reproductive effort in Mexican lizard species. Finally, after checking the homogeneity of the variances with Levene's tests and not assuming the same variances, we analysed whether the mass of neonates at birth differed between litters and the annual variation in mean litter mass using Welch's ANOVA and Games–Howell post hoc tests in SPSS ver. 24.

Results

Births

Twenty-four gravid females were captured over a 4-yr period. The SVL ranged from 51.8 mm to 69.0 mm and the mean size was 62.711 mm (SD = 4.141 mm, CV = 6.6%, n = 24). In captivity, the females gave birth to a total of 90 neonates (range = 2−5 neonates per litter, mean = 3.750, SD = 0.897, CV = 23.9%) from mid-April to the last week of June. LMs ranged from 0.659 g to 1.844 g (mean = 1.250 g, SD = 0.310, CV = 31.2%, n = 24), while MLMs ranged from 0.256 g to 0.405 g (mean = 0.330 g, SD = 0.033, CV = 10.0%, n = 24).

Reproductive effort

For *P. copei*, the estimated range for RLM was 0.129−0.530 (mean = 0.301, SD = 0.094, CV = 31.2%, n = 24), the range for INV was 0.190−0.694 g (mean = 0.428 g, SD = 0.103, CV = 24.1%, n = 24) and the range for PROD was 0.659−1.844 g × year^{-1} (mean = 1.236 g × year^{-1}, SD = 0.326, CV = 26.4%, n = 24).

Female characteristics and their effects on reproductive output

Correlation analyses revealed significant positive correlations between female SVL and LS ($r = 0.427$, $R^2 = 0.182$, $F_{1,23} = 4.903$, $p = 0.037$) and LM ($r = 0.458$, $R^2 = 0.210$, $F_{1,23} = 5.851$, $p = 0.024$) (Fig. 2A, B). Similarly, we found significant positive correlations between female TM and both litter traits (LS: $r = 0.616$, $R^2 = 0.379$, $F_{1,23} = 13.451$, $p = 0.001$; LM: $r = 0.527$, $R^2 = 0.278$, $F_{1,23} = 8.455$, $p = 0.008$) (Fig. 2C, D). Analyses between the SVL of females and the three estimated reproductive investment indices revealed a significant correlation only for PROD ($r = 0.433$, $R^2 = 0.188$, $F_{1,23} = 5.082$, $p = 0.034$), but not for INV ($r = 0.386$, $R^2 = 0.149$, $F_{1,23} = 3.885$, $p = 0.062$) or RLM ($r = 0.253$, $R^2 = 0.064$, $F_{1,23} = 1.502$, $p = 0.233$). Additionally, a negative correlation between female AM and RLM was observed ($r = -0.291$, $R^2 = 0.540$, $F_{1,23} = 9.041$, $p = 0.006$). We also observed no significant relationships between LS ($r = 0.206$, $R^2 = 0.042$, $F_{1,23} = 0.976$, $p = 0.334$) or LM ($r = 0.024$, $R^2 = 0.001$, $F_{1,23} = 0.013$, $p = 0.912$) and the TM length of females. Finally, when comparing the masses of the neonates, we observed significant variations amongst both litters (Fig. 3) ($F_{23,20.871} = 5.765$, $p < 0.001$) (Suppl. material 1), and years ($F_{3,44.102} = 6.530$, $p < 0.001$). The MLM varied amongst years and was greater in 2014 (mean = 0.3604 g) than in other years (2015 = 0.3136 g; 2016 = 0.3301 g and 2017 = 0.3266 g) (Fig. 4).



Figure 2. Relationships between individual *Plestiodon copei* female size (SVL) and (**A**) litter size, (**B**) litter mass. Additionally, relationships between individual female total mass and (**C**) litter size and (**D**) litter mass.



Figure 3. Within-litter variation in neonate mass (g) of *Plestiodon copei*. The horizontal line shows estimated mean litter mass (MLM) for the species.

Discussion

In several lizard species, reproductive investment has been shown to be associated with phenotypic and physiological traits, both of which are optimised by natural selection (Tinkle and Handley 1973; Vitt 1974; Rodríguez-Romero et al. 2002). The reproductive effort indices estimated in this study for *Plestiodon copei* showed that females invest a high amount of their resources in the development of their offspring and that the size of the female correlates with the size and aggregate weight of the offspring. In addition, Castro-Franco et al. (2011) discussed the pattern between the estimated values of RCM/RLM and their coefficients of variation (CVs) in several lizard species. They concluded that high RCM/RLM values and CV values above 20% could indicate a recurrent pattern for lizard species inhabiting extreme environments. In the case of *P. copei*, this



Figure 4. Annual variation in mean litter mass (g) of *Plestiodon copei*. Values are means ± SE. Letters denote differences between years, based on Games-Howell post-hoc test.

reproductive effort is associated with its habitat at high elevations within the TMVB. This condition has also been documented for other viviparous high-mountain lizard species within the TMVB, such as *Sceloporus bicanthalis* (Rodríguez-Romero et al. 2002), *Phrynosoma orbiculare* (Suárez et al. 2018) and Mexican skink lizards, such as *Plestiodon* sp. from Tlaxcala and *P. dugessi* (Muñoz 2015), all of which have high RLM values and > 20% variation. In the specific case of *P. copei*, the CVs of the three indices used to estimate reproductive effort were > 26%, also indicating high reproductive investment for the population studied.

The gravid females varied in both SVL and TM; these variables, in turn, showed significant correlations with litter size and litter mass. Ford and Siegel (1989) showed that several phenotypic traits can vary within reptile populations in response to different biotic and abiotic factors and that phenotypic plasticity can be expected in life-history traits, as might be the case in *P. copei*. The reproductive investment of *P. copei* females is strongly influenced by maternal size; specifically, in *P. copei*, litter size increases with increasing maternal body size. Similar results were reported by Guillette (1983) and Ramírez-Bautista et al. (1996) for viviparous or oviparous lizard species sympatric with *P. copei* within the TMVB (Rodríguez-Romero et al. 2002, 2004, 2005; Suárez et al. 2018).

The timing of birth in captivity for these study animals was consistent with the birth period reported by García (2009) for *P. copei*. Similarly, the birth months of *P. copei* were the same as those of other *Plestiodon* species inhabiting the TMVB (e.g. *P. dugesii*, *P. indubitus*, *P. lynxe* and *Plestiodon* sp. PT) (López et al. 2008; Chávez 2012; Lagunas 2014; Muñoz 2015). The date of parturition of lizards is typically associated with an abundance of food resources, which favours the growth and survival of neonates (Olsson and Shine 1998; Ramírez-Bautista et al. 1998, 2004; Chávez 2012; Lagunas 2014; Muñoz 2015), increasing their probability of reaching sexual maturity earlier (Chávez 2012).

The minimum gravid female sizes reported here indicate that Tenango females can reach sexual maturity at relatively small body sizes (SVL = 51.8 mm), as the SVL of the smallest pregnant female previously reported was 56.0 mm (Ramírez-Bautista et al. 1996); this finding indicates a relatively fast life history for the Tenango population. Ramírez-Bautista et al. (1996) also reported the growth rates of *P. copei* in the first year of life; by comparing the SVL of the smallest gravid female from Tenango, we can confirm that at least some *P. copei* females from Tenango become sexually mature in the year after birth. Variation in the body size of sexually mature females has been associated with the specific climatic conditions (temperature and precipitation) to which a population is exposed (Wapstra and O'Reilly 2001; Arribas and Galán 2005; Gutiérrez et al. 2010), as well as a combination of biotic (predation, foraging capacity, food availability and quality) and abiotic factors (climatic conditions) (Du et al. 2005, 2014; Roitberg et al. 2013; Hosseinian et al. 2014; Cruz-Elizalde and Ramírez-Bautista 2016; Maciel-Mata 2017; Wang et al. 2017; López-Alcaide et al. 2020). Based on the records collected for 59 years from meteorological stations near each population, the mean annual temperature in our study area was greater (Tenango = 13.4 °C) than those in other *P. copei* population sites reported by other authors (Cuajimalpa: 10.1 °C; Zoquiapan = 11.6 °C; Ajusco = 10.0 °C) (Suppl. material 2), suggesting that warmer temperatures in Tenango may favour *P. copei* females reaching sexual maturity at a smaller size (Kubisch et al. 2012; Cabezas-Cartes et al. 2018).

It has been shown that, particularly for ectothermic species living at high elevations, temperature is an important factor shaping life history strategies, as well as determining growth rate and age and size at sexual maturity (Atkinson and Sibly 1997; Angilletta et al. 2006; Amat and Meiri 2018; Padilla-Pérez and Angilleta Jr. 2022). For example, populations living in relatively warmer environments, such as the Tenango population compared to those of Cuajimalpa, Zoquiapan and Ajusco, are expected to have faster gonadal maturation. This would allow sexual maturity to be reached at a smaller size relative to populations living in colder regions, where sexual maturity is delayed and, in turn, reflected in the larger average body size of the gravid females (Wapstra et al. 2001; Morrison and Hero 2003; Angilletta Jr. et al. 2006; Meiri et al. 2013; Padilla-Pérez and Angilletta Jr. 2022).

The effects of temperature on determining the size of females of *P. copei* at sexual maturity are similar to those of other high-mountain lizards, such as *Mediodactylus heterocercus* and *Darevskia derjugini*. In all three lizard species, sexually mature females with smaller minimum body sizes were encountered under relatively warm temperature conditions, while sexually mature females with larger minimum sizes were found in comparatively cooler environments. (Altunışık and Eksilmez 2020; Altunışık et al. 2022). This variation in reproductive strategy amongst and within lizard species (i.e. the existence of females with early reproduction and small body size versus females with later reproduction at larger sizes) has already been recognised by earlier herpetologists within the fast–slow continuum hypothesis of life history theory (Adolph and Porter 1996; Shine 2005; Rojas-González et al. 2008; Pérez-Mendoza and Zúñiga-Vega 2014; Cruz-Elizalde and Ramírez-Bautista 2016; Boretto et al. 2018). The fast–slow continuum hypothesis proposes that organisms inhabiting colder environments and higher altitudes adopt a slow reproductive strategy, resulting

in delayed maturation, as well as comparatively smaller litters. Conversely, in warmer climates and at lower altitudes, a rapid reproductive strategy is observed, marked by early maturation and larger litters (Promilsow and Harvey 1990; Shine 2005; Li et al. 2014; Pérez-Mendoza and Zúñiga-Vega 2014). Comparing these conditions between the populations of *P. copei* in Tenango and San Lorenzo Acopilco, we observe that the Tenango population follows a faster strategy, while the San Lorenzo Acopilco population follows a slower strategy. Thus, in Tenango, which is a population living at relatively warm temperatures at lower elevations, natural selection favours sexually mature females with smaller body sizes and larger litters. Conversely, in San Lorenzo Acopilco, which has higher elevations and colder temperatures, natural selection favours larger gravid females, but smaller litters (García 2009). Moreover, the possibility that a fast life history strategy may lead to a decrease in adult female survival has also been documented (Tinkle et al. 1970; Stearns 1992; Rohr 1997; Shine 2005; Bestion et al. 2015; Boretto et al. 2017).

The mass of the offspring is another reproductive characteristic of *P. copei* that, if influenced by environmental temperature, suggests a rapid life history strategy for Tenango animals. According to the PROD index, the total mass of offspring produced in a year is greater in Tenango than in San Lorenzo Acopilco. This pattern is similar to that observed in another temperate-adapted skink species, *Eulamprus tympanum* (Rohr 1997); in populations living at higher environmental temperatures and lower elevations, the total offspring mass is significantly greater than that in populations living at colder temperatures and higher elevations. In conclusion, female lizards living in colder environments, as in the case of San Lorenzo Acopilco, evolve a slow strategy characterised by a combination of sexually mature females with larger body sizes and lower offspring masses (Adolph and Porter 1996; Rohr 1997; Badyaev and Ghalambor 2001; Pérez-Mendoza and Zúñiga-Vega 2014; López-Alcaide et al. 2020).

Our analysis of the masses of individual *P. copei* neonates revealed marked differences both within and between litters (see Fig. 3), which have been attributed to the availability of resources in the environment (Bleu et al. 2013; Wang et al. 2017) or predation pressures (Shine and Downes 1999; Bestion et al. 2014) to which the females are exposed during gestation. However, other factors could also explain this observed variation, such as sperm storage by females or multiple paternity events (Bateson et al. 2011; Chávez 2012; Muñoz 2015). Such factors have been suggested to be important in species of the genus *Plestiodon* (Ramírez-Bautista et al. 1996; Bateson et al. 2011) and may matter for *P. copei* as well, although this has not yet been demonstrated for this species.

Moreover, the differences in MLM observed between years suggest that there are other factors that could influence the reproductive investment of *P. copei* females in Tenango. Possible factors include differences in the SVL of females or in environmental conditions, such as temperature variation and food availability (Ballinger 1977; Ji et al. 2007; Warner et al. 2007; Bleu et al. 2013; Cruz-Elizalde and Ramírez-Bautista 2016; Guo et al. 2022), which could be responsible for the heavier litters produced in 2014 compared to those in the three subsequent years. First, we ruled out the possibility that the greater MLM in 2014 was due to female size variation because the female SVL did not significantly differ across the four years. Second, data from the nearest meteorological station showed that there was no significant variation in environmental temperature that might account for the heavier litters in 2014 than in 2015–2017.

However, the lower MLMs recorded since 2015 could be attributed to reduced food availability, as the collection site showed repeated signs of human-caused grass fires due to traditional agricultural and grazing activities that allowed fires to spread into the forest. This type of vegetation disturbance has been shown – due to the lethal surface temperatures caused by fire – to negatively impact the abundance of ground-dwelling beetles which, in turn, are an important food source for *P. copei* (Nunes et al. 2006; Elia et al. 2012). The quantity and quality of the diet of females strongly influence their reproductive output (Selman and Huston 1996; Warner et al. 2007; Lovern and Adams 2008). Low feeding during fire years reduced MLM not only during that year, but also in subsequent years, as food resources for *P. copei* females decreased (Ballinger 1997; Warner et al. 2015). Bleu et al. (2013) documented that food availability is correlated with reproductive investment in *Zootoca vivipara*. Furthermore, in *Vipera aspis*, the size of the neonate is influenced by the food intake of the female immediately before ovulation and litter size depends on both long-term reserves and recent food intake (Bonnet et al. 2001). Thus, if human-caused grass fires in Tenango indeed resulted in a significant decrease in prey items consumed by females prior to fertilisation and during pregnancy, this could explain why MLM decreased in 2015–2017; however, this hypothesis needs to be tested in future studies.

In conclusion, by comparing the RLM, INV and PROD indices used to evaluate reproductive effort, we observed that *P. copei* had slightly greater values than the mean estimated for species in the *P. brevirostris* group (Table 1). The high reproductive effort values obtained for *P. copei* might be a result of inhabiting areas with extreme environmental conditions (e.g. high altitude) within the TMVB. Elevation is considered an important factor influencing the evolution of life history traits, especially in *P. copei*, as it is the most important abiotic factor determining its geographic distribution within the TMVB (Alvarado-Avilés et al. 2020). For example, colder temperatures and limited food availability at higher altitudes have been shown to lead to greater reproductive effort in high-altitude species, such as *P. copei* (Rohr 1997; Badyaev and Ghalambor 2001). Overall, we observed that the RLM obtained for *P. copei* was similar to the estimated values for other species of the Scincidae family (Vitt and Price 1982; Qualls and Shine 1997). However, when we examined data from only the species *Plestiodon* from Mexico, we observed that the RLM estimated for *P. copei* was slightly greater than those reported for *P. dugesii* and *Plestiodon* sp. PT from Jalisco and Tlaxcala, which live

Table 1. Reproductive effort estimations for several Mexican species of *Plestiodon*.

Species	Location	RCM / RLM	INV (g)	PROD (g × year^{-1})	Reference
Plestiodon bilineatus (n = 1)	Valparaíso, Zacatecas	–	–	1.820	Bañuelos et al. (2016)
Plestiodon sp. PT* (n = 7)	Volcán La Malinche, Tlaxcala	0.231	0.432	0.850	Muñoz (2015)
Plestiodon sp. PT*(n = 17)	Chilchota, Puebla	0.334	0.325	1.168	Chávez (2012)
Plestiodon sp. OX* (n = 1)	Santa Inés del Monte, Oaxaca	–	0.515	1.240	López et al. (2008)
P. indubitus (n = 1)	Sur de Tres Marías, Morelos	–	–	0.640	López et al. (2008)
P. copei (n = 3)	San Lorenzo Acopilco, CDMX	–	–	1.020	García (2009)
P. copei (n = 24)	Tenango, Estado de México	0.301	0.428	1.236	This study
P. dicei (n = 1)	Pablillo, Nuevo León	–	–	1.212	Feria-Ortiz et al. (2007)
P. dicei (n = 1)	San Antonio de las Alazanas, Coahuila	–	–	1.164	Feria-Ortiz et al. (2007)
P. dugesii (n = 21)	Mazamitla, Jalisco	0.292	0.362	1.001	Muñoz (2015)
Averages		0.290	0.412	1.135	

* Undescribed *Plestiodon* species reported by Pavón-Vázquez et al. (2018): *Plestiodon* sp. PT from Puebla and Tlaxcala; *Plestiodon* sp. OX from Oaxaca.

at lower elevations (Muñoz 2015). Conversely, the RLM obtained for *P. copei* was slightly lower than that estimated for the population of *Plestiodon* sp. PT from Puebla (Chávez 2012). We also observed that the PROD index calculated for *P. copei* from Tenango was greater than that (0.830) estimated by Meiri et al. (2012) for this species.

The observations presented here suggest that the reproductive strategy of *P. copei* from Tenango may have been shaped by the prevailing extreme environmental conditions (Castro-Franco et al. 2011; Bestion et al. 2015; Muñoz 2015), although it will be necessary to evaluate this hypothesis in future studies using both larger sample sizes and additional study sites. Furthermore, it will also be important to evaluate the effects of global warming and habitat loss on the reproductive strategies and sex ratio of this specialist skink.

Acknowledgments

We thank to E. Carbajal and S. H. Cuadros for their assistance in the field. Collecting permits (SGPA/DGVS/08371/14) were provided by the Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT). We also thank the anonymous reviewers for comments and suggestions to the manuscript.

References

Abell AJ (1999) Variation in clutch size and offspring size relative to environmental conditions in the lizard *Sceloporus virgatus*. Journal of Herpetology 33: 173−180. https://doi.org/10.2307/1565712

Adolph SC, Porter WP (1996) Growth, seasonality, and lizard life histories: age and size at maturity. Oikos 77: 267−278. https://doi.org/10.2307/3546065

Altunışık A, Eksilmez H (2020) Age, growth and survival rate in two populations of *Darevskia derjugini* (Nikolsky, 1898) from different altitudes (Squamata: Sauria: Lacertidae). Animal Biology 71(2): 135−149. https://doi.org/10.1163/15707563-bja10025

Altunışık A, Yildiz MZ, Üçeş F, Bozkurt MA, Sömer M (2022) Altitude impacts body size but not age in the Asia Minor Thin-toed Gecko (*Mediodactylus heterocercus*). Canadian Journal of Zoology 100(5): 315−321. https://doi.org/10.1139/cjz-2021-0197

Alvarado-Avilés JC, Venegas-Barrera CS, Alvarado-Díaz J, Hernández-Gallegos O, Ruiz-Gomez MDL, Sánchez-Sánchez H (2020) Potential distribution of *Plestiodon copei* (Squamata: Scincidae), an endemic and threatened lizard of México. South American Journal of Herpetology 18: 56−67. https://doi.org/10.2994/10.2994/SAJH-D-18-00044.1

Amat F, Meiri S (2018) Geographical, climatic and biological constraints on age at sexual maturity in amphibians. Biological Journal of the Linnean Society 123(1): 34–42. https://doi.org/10.1093/biolinnean/blx127

Angilletta MJ, Oufiero CE, Leaché AD (2006) Direct and indirect effects of environmental temperature on the evolution of reproductive strategies: An information-theoretic approach. The American Naturalist 168(4): 123–135. https://doi.org/10.1086/507880

Arribas OJ, Galán P (2005) Reproductive characteristics of the Pyrenean high-mountain lizards: *Iberolacerta aranica* (Arribas, 1993), *I. aurelioi* (Arribas, 1994) and *I. bonnali* (Lantz, 1927). Animal Biology 55: 163−190. https://doi.org/10.1163/1570756053993505

Atkinson D, Sibly RM (1997) Why are organisms usually bigger in colder environments? Making sense of a life history puzzle. Trends in Ecology & Evolution 12(6): 235–239. https://doi.org/10.1016/S0169-5347(97)01058-6

Badyaev AV, Ghalambor CK (2001) Evolution of life histories along elevational gradients: Trade-off between parental care and fecundity. Ecology 82(10): 2948−2960. https://doi.org/10.1890/0012-9658(2001)082[2948:EOLHAE]2.0.CO;2

Ballinger RE (1977) Reproductive strategies: food availability as a source of proximal variation in a lizard. Ecology 58(3): 628−635. https://doi.org/10.2307/1939012

Bañuelos JA, Carbajal RA, Quintero GE (2016) *Plestiodon bilineatus*. Reproduction. Mesoamerican Herpetology 3: 728−729.

Bastiaans E, Méndez F, Rodríguez K, Flores CF, Sinervo B (2013) Female reproductive investment in the mesquite lizard (*Sceloporus grammicus*) species complex (Squamata: Phrynosomatidae). The Southwestern Naturalist 58: 335−344. https://doi.org/10.1894/0038-4909-58.3.335

Bateson ZW, Krenz JD, Sorensen RE (2011) Multiple paternity in the common five-lines skink (*Plestiodon fasciatus*). Journal of Herpetology 45: 504−510. https://doi.org/10.1670/10-295.1

Bestion E, Teyssier A, Richard M, Clobert J, Cote J (2015) Live fast, die young: experimental evidence of population extinction risk due to climate change. PLoS Biology 13: 1−19. https://doi.org/10.1371/journal.pbio.1002281

Bleu J, Galliard JF, Fitze P, Meylan S, Clobert J, Massot M (2013) Reproductive allocation strategies: a long-term study on proximate fac-

tors and temporal adjustments in a viviparous lizard. Oecologia 171: 141–151. https://doi.org/10.1007/s00442-012-2401-1

Bonnet X, Naulleau G, Shine R, Lourdais O (2001) Short-term versus long-term effects of food intake on reproductive output in a viviparous snake, *Vipera aspis*. Oikos 92: 297–308. https://doi.org/10.1034/j.1600-0706.2001.920212.x

Boretto JM, Cabezas-Cartes F, Ibargüengoytía NR (2018) Slow life histories in lizards living in the highlands of the Andes Mountains. Journal of Comparative Physiology B 188(3): 491–503. https://doi.org/10.1007/s00360-017-1136-z

Bustos MG, Méndez FR, Castro R, Villagrán M (2011) Ciclo reproductor de *Sceloporus ochoterenae* en el Estado de Morelos, México. Revista Mexicana de Biodiversidad 82: 589–597. https://doi.org/10.22201/ib.20078706e.2011.2.471

Cabezas-Cartes F, Boretto JM, Ibargüengoytía NR (2018) Effects of climate and latitude on age at maturity and longevity of lizards studied by skeletochronology. Integrative and Comparative Biology 58(6): 1086–1097. https://doi.org/10.1093/icb/icy119

Castro-Franco R, Bustos MG, Méndez FR (2011) Variation in parental investment and relative clutch mass of the spiny-tail iguana, *Ctenosaura pectinata* (Squamata: Iguanidae) in central México. Revista Mexicana de Biodiversidad 82: 199–204. https://doi.org/10.22201/ib.20078706e.2011.1.454

Chávez L (2012) Biología reproductiva de *Plestiodon brevirostris* (Squamata: Scincidae) en el centro-este del estado de Puebla. Bachelor Thesis, Facultad de Estudios Superiores Zaragoza, Universidad Nacional Autónoma de México, Ciudad de México, México.

Congdon JD, Gibbons JW (1987) Morphological constraint on egg size: a challenge to optimal egg size theory? Proceedings of the National Academy of Sciences USA 84: 4145–4147. https://doi.org/10.1073/pnas.84.12.4145

Cruz-Elizalde R, Ramírez-Bautista A (2016) Reproductive cycles and reproductive strategies among populations of the Rose-bellied Lizard *Sceloporus variabilis* (Squamata: Phrynosomatidae) from central Mexico. Ecology and Evolution 6: 1753–1768. https://doi.org/10.1002/ece3.1998

Cruz-Elizalde R, Ramírez-Bautista A (2016) Reproductive cycles and reproductive strategies among populations of the Rose-bellied Lizard *Sceloporus variabilis* (Squamata: Phrynosomatidae) from central Mexico. Ecology and Evolution 6(6): 1753–1768. https://doi.org/10.1002/ece3.1998

De Marco VG (1989) Annual variation in the seasonal shift in egg size and clutch size in *Sceloporus woodi*. Oecologia 80: 525–553. https://doi.org/10.1007/BF00380077

Du W, Ji X, Shine R (2005) Does body-volume constrain reproductive output in lizards? Biology Letters 1: 98–100. https://doi.org/10.1098/rsbl.2004.0268

Du W, Robbins TR, Warner DA, Langkilde T, Shine R (2014) Latitudinal and seasonal variation in reproductive effort of the eastern fence lizard (*Sceloporus undulatus*). Integrative Zoology 9: 360–371. https://doi.org/10.1111/1749-4877.12072

Dunham AE, Miles DB, Reznick DN (1988) Life history patterns in squamate reptiles. In: Gans C, Huey RB (Eds) Biology of the Reptilia. Copeia, New York, 441–522.

Elia M, Lafortezza R, Tarasco E, Colangelo G, Sanesi G (2012) The spatial and temporal effects of fire on insect abundance in Mediterranean forest ecosystems. Forest Ecology and Management 263: 262–267. https://doi.org/10.1016/j.foreco.2011.09.034

Feria-Ortiz M, García UO, Aguilar JL (2007) *Plestiodon brevirostris* (Short-nosed Skink). Reproduction. Herpetological Review 38: 81–82.

Feria-Ortiz M, Nieto A, Salgado IH (2001) Diet and reproductive biology of the viviparous lizard *Sceloporus torquatus* (Squamata: Phrynosomatidae). Journal of Herpetology 35: 104–112. https://doi.org/10.2307/1566029

Ford NB, Seigel RA (1989) Phenotypic plasticity in reproductive traits: evidence from a viviparous snake. Ecology 70(6): 1768–1774. https://doi.org/10.2307/1938110

Forsman A, Shine R (1995) Parallel geographic variation in body shape and reproductive life history within the Australian scincid lizard *Lampropholis delicata*. Functional Ecology 9: 818–828. https://doi.org/10.2307/2389979

García UO (2009) Reporte del tamaño de camada de *Plestiodon copei* (Squamata: Scincidae). Boletín de la Sociedad Herpetológica Mexicana 17: 96–99.

Guillette LJ (1983) Notes concerning reproduction of the montane skink, *Eumeces copei*. Journal of Herpetology 17: 144–148. https://doi.org/10.2307/1563455

Guillette LJ, Casas-Andreu G (1987) The reproductive biology of the high elevation Mexican lizard *Barisia imbricata*. Herpetologica 43: 29–38.

Guo K, Xiang-Mo L, Yan-Qing W, Yan-Fu Q, Xiang J (2022) Measuring annual variation in reproductive output reveals a key role of maternal body condition in determining the size of eggs in snakes. Animals 12(2): 1494. https://doi.org/10.3390/ani12121494

Gutiérrez J, Krenz JD, Ibargüengoytía NR (2010) Effect of altitude on thermal responses of *Liolaemus pictus argentinus* in Argentina. Journal of Thermal Biology 35: 332–337. https://doi.org/10.1016/j.jtherbio.2010.07.001

Hosseinian SS, Yousefi M, Mohammadpour A, Masroor R, Rastegar N (2014) Phenotypic variation in males of the agamid lizard *Paralaudakia caucasia* (Eichwald, 1831) across a wide geographic range. Herpetologica 70: 464–471. https://doi.org/10.1655/HERPETOLOGICA-D-13-00079

Hutchings JA (2021) A primer of life histories: ecology, evolution, and application. Oxford University Press, Glasgow, 226 pp. https://doi.org/10.1093/oso/9780198839873.001.0001

IBM Corporation (2016) IBM SPSS Statistics, Version 24.0. IBM Corp, Armonk, New York.

Ji X, Du WG, Li H, Lin LH (2006) Experimentally reducing clutch size reveals a fixed upper limit to egg size in snakes, evidence from the king ratsnake, *Elaphe carinata*. Comparative Biochemistry and Physiology 144: 474–478. https://doi.org/10.1016/j.cbpa.2006.04.013

Ji X, Du WG, Lin ZH, Luo LG (2007) Measuring temporal variation in reproductive output reveals optimal resource allocation to reproduction in the northern grass lizard, *Takydromus septentrionalis*. Biological Journal of the Linnean Society 91(2): 315–324. https://doi.org/10.1111/j.1095-8312.2007.00791.x

Kubisch E, Piantoni C, Williams J, Scolaro A, Navas CA, Ibargüengoytía NR (2012) Do higher temperatures increase growth in the nocturnal gecko *Homonota darwini* (Gekkota: Phyllodactylidae)? A skeletochronological assessment analyzed at temporal and geographic scales. Journal of Herpetology 46(4): 587–595. https://doi.org/10.1670/10-277

Lagunas AI (2014) Ciclos reproductivos y de lípidos de la lagartija vivípara *Plestiodon indubitus* (Scincidae) al Noroeste del Estado de Morelos. Bachelor Thesis, Facultad de Estudios Superiores Zaragoza, Universidad Nacional Autónoma de México, Ciudad de México, México.

Li J, Zhou R, Liu N (2014) Life-history variation among three populations of the toad-headed lizard *Phrynocephalus vlangalii* along an elevation gradient on the northeastern Tibetan Plateau. The Herpetological Journal 24(1): 17−23.

López ID, Feria M, Martínez M (2008) Tamaño de camada en *Plestiodon brevirostris*. Boletín de la Sociedad Herpetológica Mexicana 16: 15−17.

López-Alcaide S, Cuateta-Bonilla LY, Macip-Ríos R (2020) Reproductive output in spiny lizards (genus *Sceloporus*) with different reproductive mode. A comparative approach. Revista Mexicana de Biodiversidad 91: e913020. https://doi.org/10.22201/ib.20078706e.2020.91.3020

Lovern MB, Adams AL (2008) The effects of diet on plasma and yolk steroids in lizards (*Anolis carolinensis*). Integrative and comparative biology 48(3): 428–436. https://doi.org/10.1093/icb/icn058

Maciel-Mata C (2017) Caracterización de la estructura de nicho ecológico de la lagartija *Sceloporus minor* (Squamata: Phrynosomatidae): evaluación sobre atributos de historia de vida. PhD Thesis, Universidad Autónoma del Estado de Hidalgo, Hidalgo, México.

Mathies T, Andrews RM (1995) Thermal and reproductive biology of high and low elevation populations of the lizard *Sceloporus scalaris*: implications for the evolution of viviparity. Oecologia 104: 101−111. https://doi.org/10.1007/BF00365568

Meiri S, Bauer AM, Chirio L, Colli GR, Das I, Doan TM, Feldman A, Herrera FC, Novosolov M, Pafilis P, Pincheira-Donoso D, Powney G, Torres-Carbajal O, Uetz P, Van Damme R (2013) Are lizards feeling the heat? A tale of ecology and evolution under two temperatures. Global Ecology and Biogeography 22(7): 834−845. https://doi.org/10.1111/geb.12053

Meiri S, Brown JH, Sibly RM (2012) The ecology of lizard reproductive output. Global Ecology and Biogeography 21: 592−602. https://doi.org/10.1111/j.1466-8238.2011.00700.x

Morales ME (2014) Morfometría comparada y etología de la reproducción de *Plestiodon copei* (Squamata: Scincidae). Bachelor Thesis, Facultad de Ciencias, Universidad Autónoma del Estado de México, Estado de México, México.

Morrison C, Hero JM (2003) Geographic variation in life history characteristics of amphibians: a review. Journal of Animal Ecology 72: 270–279. https://doi.org/10.1046/j.1365-2656.2003.00696.x

Muñoz FJ (2015) Esfuerzo reproductor y ecología térmica de dos especies vivíparas del género *Plestiodon* (Squamata: Scincidae) de México. Bachelor Thesis, Centro Universitario de Ciencias Biológicas y Agropecuarias, Universidad de Guadalajara, Guadalajara, México.

Nunes L, Silva I, Pité M, Rego F, Leather S, Serrano A (2006) Carabid (Coleoptera) community changes following prescribed burning and the potential use of carabids as indicators species to evaluate the effects of fire management in Mediterranean regions. Silva Lusitana 14: 85–100.

Olsson M, Shine R (1998) Timing of parturition as a maternal care tactic in an alpine lizard species. Evolution 52(6): 1861−1864. https://doi.org/10.1111/j.1558-5646.1998.tb02265.x

Padilla-Perez DJ, Angilletta Jr MJ (2022) Macroclimatic and maternal effects on the evolution of reproductive traits in lizards. Ecology and Evolution 12(5): e8885. https://doi.org/10.1002/ece3.8885

Padilla-Perez DJ, DeNardo DF, Angilletta Jr MJ (2022) Early transition in foraging mode promoted greater reproductive effort in widely foraging lizards. bioRxiv: 2022.01.22.477347. https://doi.org/10.1101/2022.01.22.477347

Pavón-Vázquez CJ, García-Vázquez UO, Bryson Jr RW, Feria-Ortiz M, Manríquez-Morán NL, Nieto-Montes de Oca A (2018) Integrative species delimitation in practice: revealing cryptic lineages within the short-nosed skink *Plestiodon brevirostris* (Squamata: Scincidae). Molecular Phylogenetics and Evolution 129: 242–257. https://doi.org/10.1016/j.ympev.2018.08.020

Pérez-Mendoza HA, Zúñiga-Vega JJ (2014) A test of the fast–slow continuum model of life-history variation in the lizard *Sceloporus grammicus*. Evolutionary Ecology Research 16(3): 235−248.

Promislow DE, Harvey PH (1990) Living fast and dying young: A comparative analysis of life-history variation among mammals. Journal of Zoology 220(3): 417−437. https://doi.org/10.1111/j.1469-7998.1990.tb04316.x

Qualls FJ, Shine R (1997) Geographic variation in 'costs of reproduction' in the scincid lizard *Lampropholis guichenoti*. Functional Ecology 11: 757−763. https://doi.org/10.1046/j.1365-2435.1997.00150.x

Ramírez-Bautista A, Cruz-Elizalde R, Hernández-Salinas U, Lozano A, Grummer JA (2017) Reproductive trait variation in the *Sceloporus scalaris* species group (Squamata: Phrynosomatidae) from the Transvolcanic Belt, Mexico. Biological Journal of the Linnean Society 122(4): 838−849. https://doi.org/10.1093/biolinnean/blx100

Ramírez-Bautista A, Jiménez-Cruz E, Marshall JC (2004) Comparative life history for populations of the *Sceloporus grammicus* complex (Squamata: Phrynosomatidae). Western North American Naturalist 64(2): 175−183.

Ramírez-Bautista AJ, Arizmendi MC (2004) *Eumeces copei*. Sistemática e Historia Natural de algunos anfibios y reptiles de México. Facultad de Estudios Superiores Iztacala, Universidad de Biología, Tecnología y prototipos (UBIPRO), Universidad Nacional Autónoma de México. Base de datos SNIB-CONABIO. Proyecto W013, México.

Ramírez-Bautista AJ, Barba J, Vitt LJ (1998) Reproductive cycle and brood size of *Eumeces lynxe* from Pinal de Amoles, Queretaro, México. Journal of Herpetology 32: 18−24. https://doi.org/10.2307/1565474

Ramírez-Bautista AJ, Guillette L, Gutierrez G, Uribe Z (1996) Reproductive biology of the lizard "*Eumeces copei*"(Lacertilia: Scincidae) from the Eje Neovolcanico, Mexico. The Southwestern Naturalist 41: 103−110.

Ramírez-Bautista AJ, Hernández U, Cruz-Elizalde R, Lozano A, Rodríguez-Romero FDJ (2016) Sexual dimorphism and reproductive traits over time in *Sceloporus aeneus* (Squamata: Phrynosomatidae), based on a population in the Trans Mexican Volcanic Belt, Mexico. Salamandra 52: 197−203.

Rodríguez-Romero FDJ, Méndez FR, García R, Cruz VS (2002) Comparación del esfuerzo reproductor en dos especies hermanas del género *Sceloporus* (Sauria: Phrynosomatidae) con diferente modo reproductor. Acta Zoológica Mexicana 85: 181−188. https://doi.org/10.21829/azm.2002.85851821

Rodríguez-Romero FDJ, Méndez FR, López LL (2005) Análisis comparado del esfuerzo reproductor en algunos lacertilios mexicanos de ambientes tropical y templado. Revista de la Sociedad Mexicana de Historia Natural 2: 168–177.

Rodríguez-Romero FDJ, Smith GR, Cuellar O, Méndez FR (2004) Reproductive traits of a high elevation viviparous lizard *Sceloporus bicanthalis* (Lacertilia: Phrynosomatidae) from Mexico. Journal of Herpetology 38: 438−443. https://doi.org/10.1670/7-04N

Rohr DH (1997) Demographic and life-history variation in two proximate populations of a viviparous skink separated by a steep altitudinal gradient. Journal of Animal Ecology 66: 567−578. https://doi.org/10.2307/5950

Roitberg ES, Kuranova VN, Bulakhova NA, Orlova VF, Eplanova GV, Zinenko OI, Shamgunova RR, Hofmann S, Yakovlev VA (2013)

Variation of reproductive traits and female body size in the most widely-ranging terrestrial reptile: testing the effects of reproductive mode, lineage, and climate. Evolutionary Biology 40: 420−438. https://doi.org/10.1007/s11692-013-9247-2

Rutschmann A, Miles DB, Clobert J, Richard M (2016) Warmer temperatures attenuate the classic offspring number and reproductive investment trade-off in the common lizard, *Zootoca vivipara*. Biology Letters 12: 20160101. https://doi.org/10.1098/rsbl.2016.0101

Schwarzkopf L (1993) Costs of reproduction in water skinks. Ecology 74: 1970−1981. https://doi.org/10.2307/1940840

Selman RG, Houston DC (1996) The effect of prebreeding diet on reproductive output in zebra finches. Proceedings of the Royal Society: Biological Sciences 263(1376): 1585–1588. https://doi.org/10.1098/rspb.1996.0232

Shine R (2005) Life-history evolution in reptiles. Annual Review of Ecology, Evolution, and Systematics 36: 23−46. https://doi.org/10.1146/annurev.ecolsys.36.102003.152631

Shine R, Schwarzkopf L (1992) The evolution of reproductive effort in lizards and snakes. Evolution 46: 62−75. https://doi.org/10.2307/2409805

Sinervo B, Litch P (1991) Proximate constraints on the evolution of egg size, number, and total clutch mass in lizards. Science 252: 1300−1302. https://doi.org/10.1126/science.252.5010.1300

Sokal RR. Rohlf FJ (2012) Biometry. WH Freeman and Company, New York, 915 pp.

Stearns SC (1992) The evolution of life histories. Oxford University Press, London, 264 pp.

Suárez O, Suárez G, Chávez F, Pérez F, Andrade G, Aguilar L, Hernández O (2018) Masa relativa de la camada en *Phrynosoma orbiculare* (Squamata: Phrynosomatidae) en el Parque Estatal Sierra Morelos, Toluca, Estado de México. Revista Mexicana de Biodiversidad 89: 282−289. https://doi.org/10.22201/ib.20078706e.2018.1.1925

Tinkle DW, Hadley NF (1973) Reproductive effort and winter activity in the viviparous montane lizard *Sceloporus jarrovi*. Copeia 1973: 272−277. https://doi.org/10.2307/1442965

Tinkle DW, Hadley NF (1975) Lizard reproductive effort: caloric estimates and comments on its evolution. Ecology 56: 427–434. https://doi.org/10.2307/1934973

Tinkle DW, Wilbur HM, Tilley SG (1970) Evolutionary strategies in lizard reproduction. Evolution 24: 55–74. https://doi.org/10.2307/2406714

Vitt LJ (1974) Reproductive effort and energy comparisons of adults, eggs, and neonates of *Gerrhonotus coeruleus principis*. Journal of Herpetology 8: 165−168. https://doi.org/10.2307/1562813

Vitt LJ, Congdon JD (1978) Body shape, reproductive effort, and relative clutch mass in lizards: resolution of a paradox. American Naturalist 112: 595−608. https://doi.org/10.1086/283300

Vitt LJ, Cooper WEJR (1986) Skink reproduction and sexual dimorphism: *Eumeces fasciatus* in the southeastern United States, with notes on *Eumeces inexpectatus*. Journal of Herpetology 20: 65−76. https://doi.org/10.2307/1564126

Vitt LJ, Price H (1982) Ecological and evolution determinants of relative clutch mass in lizards. Herpetologica 38: 237−255.

Wang Y, Zeng ZG, Ma L, Li SR, Du WG (2017) Food restriction affects maternal investment but not neonate phenotypes in a viviparous lizard. Zoological Research 38: 81−87. https://doi.org/10.24272/j.issn.2095-8137.2017.011

Wapstra E, O´Reilly JM (2001) Potential "costs of reproduction" in a skink: Inter-and intrapopulational variation. Austral Ecology 26: 179−186. https://doi.org/10.1046/j.1442-9993.2001.01104.x

Wapstra E, Swain R, O'Reilly JM (2001) Geographic variation in age and size at maturity in a small Australian viviparous skink. Copeia 2001(3): 646−655. https://doi.org/10.1643/0045-8511(2001)001[0646:GVIAAS]2.0.CO;2

Warne RW, Charnov EL (2008) Reproductive allometry and the size-number trade-off for lizards. American Naturalist 172: E80−E98. https://doi.org/10.1086/589880

Warner DA, Lovern MB, Shine R (2007) Maternal nutrition affects reproductive output and sex allocation in a lizard with environmental sex determination. Proceedings of the Royal Society B: Biological Sciences 274(1611): 883–890. https://doi.org/10.1098/rspb.2006.0105

Warner DA, Buckelew AM, Pearson PR, Dhawan A (2015) The effect of prey availability on offspring survival depends on maternal food resources. Biological Journal of the Linnean Society 115(2): 437–447. https://doi.org/10.1111/bij.12519

Zamudio KR, Parra-Olea G (2000) Reproductive mode and female reproductive cycles of two endemic Mexican horned lizards (*Phrynosoma taurus* and *Phrynosoma braconnieri*). Copeia 2000(1): 222−229. https://doi.org/10.1643/0045-8511(2000)2000[0222:RMAFRC]2.0.CO;2

Supplementary material 1

Games-Howell post-hoc multiple comparisons

Authors: Hermilo Sánchez-Sánchez, Adriana Jocelyn Morales-Gonzaga, Ken Oyama, J. Carlos Alvarado-Avilés

Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/herpetozoa.36.e97705.suppl1

Supplementary material 2

Mean monthly temperature based on 59 years of records (1951–2010) for four populations of *Plestiodon copei*

Authors: Hermilo Sánchez-Sánchez, Adriana Jocelyn Morales-Gonzaga, Ken Oyama, J. Carlos Alvarado-Avilés

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/herpetozoa.36.e97705.suppl2